



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 15/00, 17/00, A61K 38/08, C07K 7/66		A1	(11) International Publication Number: WO 97/28172 (43) International Publication Date: 7 August 1997 (07.08.97)
(21) International Application Number: PCT/US97/01097 (22) International Filing Date: 17 January 1997 (17.01.97) (30) Priority Data: 08/588,773 19 January 1996 (19.01.96) US (71) Applicant (for all designated States except US): THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HIRSCHMANN, Ralph, F. [US/US]; 740 Palmer Place, Blue Bell, PA 19422 (US). NICOLAOU, Kyriacos, C. [US/US]; 9625 Blackgold Road, LaJolla, CA 92037 (US). PIETRANICO, Sherrie [US/US]; 239 Germantown Road, West Milford, NJ 07480 (US). REISINE, Terry [US/US]; 438 South 44th Street, Philadelphia, PA 19104 (US). SALVINO, Joseph [US/US]; 272 2nd Street, Schwenksville, PA 19473 (US). STRADER, Catherine [US/US]; 119 Moringside Road, Verona, NJ 07044 (US). SPRENGELER, Paul [US/US]; 614 S. 26th Street, Philadelphia, PA 19146 (US). SMITH, Amos, B., III [US/US]; 517 General Lafayette Road, Merion, PA 19066 (US). YAO, Wenqing [CN/US]; 5918 Drexelbrook Drive,		Drexel Hill, PA 19026 (US). CICHY, Maria [US/US]; 201 S. 18th Street #717, Philadelphia, PA 19103 (US). HYNES, John, Jr. [US/US]; Sandy Hill Road, Laverock, PA 19038 (US). (74) Agents: ELDERKIN, Diann, B. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: NON-PEPTIDE PEPTIDOMIMETICS AND RELATED CYCLIC HEXAPEPTIDES			
(57) Abstract Compounds are provided which are crossreactive with peptides such as those which bind G-protein-linked receptors, together with preparative and therapeutic methods therefor.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**NON-PEPTIDE PEPTIDOMIMETICS
AND RELATED CYCLIC HEXAPEPTIDES**

GOVERNMENT SUPPORT

Certain of the inventors have been supported by National Institute of Mental Health Grant 45533 and National Institutes of Health Grant GM-41821.

5 CROSS REFERENCE TO RELATED APPLICATIONS

This patent application is a continuation-in-part of application Serial No. 08/144,660, filed October 28, 1993, which is a continuation-in-part of application Serial No. 07/748,826, filed August 22, 1991. The contents of both of
10 these patent applications are incorporated by reference herein.

FIELD OF THE INVENTION

This invention relates to compounds which bind G-protein-linked receptors. In particular, the invention relates to cyclic hexapeptides which bind G-protein-linked receptors, and
15 to synthetic compounds which mimic or inhibit the biological and/or chemical activity of such peptides.

BACKGROUND OF THE INVENTION

Peptides are implicated in a wide variety of biochemical
20 processes in humans and other mammals. For example, it is known that a number of hormones and neurotransmitters are controlled by receptor-mediated stimulation of one or more of a family of guanine nucleotide-binding regulatory proteins, known as G-proteins. G-proteins activate or inhibit different

- 2 -

effector enzymes, modulating the levels of intracellular second messengers. At least 50 sub-types of G-protein-linked receptors have been identified, among them the α -adrenergic, β -adrenergic, muscarinic, cholinergic, dopamine, histamine, adenosine, serotonin, prostaglandin, leukotriene, thromboxane, prostacyclin, PAF, CAMP, enkephalin, endorphin, cholecystokinin, bombesin, substance K, substance P, neuromedin, bradykinin, FMLP, C5a, C3a, vasopressin, oxytocin, angiotensin, VIP, parathyroid hormone, calcitonin, neurotensin, TRH, somatostatin, rhodopsin, epinephrine, norepinephrine, acetylcholine, S-hydroxytryptamine, thyrotropin, thyrotropin releasing hormone, follicle stimulating, lutropin, choriogonadotropin, thrombin, retinal, and olfactory receptors. Nine or more G-proteins and at least seven effector systems have also been described. All of the G-protein-linked receptors analyzed to date contain from one to three potential sites of asparagine-linked glycosylation. The transmembrane signaling pathway used by G-protein-linked receptors represents one of the major mechanism of signal transduction in cellular systems. It is known, for example, that substance P acts as a vasodilator, a depressant, stimulates salivation, and produces increased capillary permeability. Substance P is a naturally occurring undecapeptide belonging to the tachykinin family of peptides, the latter being so-named because of their prompt contractile action on extravascular smooth muscle tissue. In addition to substance P (neurokinin-1, NK-1), the known mammalian tachykinins include neurokinin A (NK-2) and neurokinin B (NK-2). The tachykinins have been implicated in gastrointestinal (GI) disorders and diseases of the GI tract, such as inflammatory bowel disease, ulcerative colitis and Crohn's disease.

Substance P is known to produce both analgesia and hyperalgesia in animals, depending on dose and pain responsiveness of the animal and plays a role in sensory transmission and pain perception. Substance P also is believed to be involved in the inflammatory response in diseases such as rheumatoid arthritis and osteoarthritis. Other disease

- 3 -

areas where the tachykinins are believed to be involved include allergic conditions, immunoregulation, bronchospasm, reflex or neuronal control of the viscera, and Alzheimer's disease and Downs Syndrome.

5 To date, there have been limited therapeutic applications involving peptides, due in considerable part to lack of oral bioavailability and to proteolytic degradation. Typically, for example, peptides are rapidly degraded *in vivo* by exo- and endopeptidases, resulting in generally very short biological
10 half-lives. Another deficiency of peptides as potential therapeutic agents is their lack of bioavailability via oral administration. Degradation of the peptides by proteolytic enzymes in the gastrointestinal tract is likely an important contributing factor. The problem is, however, more
15 complicated, because it has been recognized that even small, cyclic peptides which are not subject to rapid metabolic inactivation nevertheless exhibit poor oral bioavailability. This likely is due to poor transport across the intestinal membrane and rapid clearance from the blood by hepatic
20 extraction with subsequent excretion into the intestine. These observations suggest that multiple amide bonds may interfere with oral bioavailability.

 The design of peptide mimics which are resistant to degradation by proteolytic enzymes has become of increasing
25 interest to peptide chemists, both for hormone agonist/antagonist and for enzyme inhibitor design. A primary goal has been to reduce the susceptibility of mimics to cleavage and inactivation by peptidases. In one approach, such as disclosed by Sherman and Spatola, *J. Am. Chem. Soc.*, 112,
30 1990, 433, one or more amide bonds have been replaced in an essentially isosteric manner by a variety of chemical functional groups. This stepwise approach has met with some success in that active analogs have been obtained. In some instances, these analogs have been shown to possess longer
35 biological half-lives than their naturally-occurring counterparts. Nevertheless, this approach has limitations. Successful replacement of more than one amide bond has been

- 4 -


rare. Consequently, the resulting analogs have remained susceptible to enzymatic inactivation elsewhere in the molecule. Moreover, this approach does not permit generalizations between chemically unrelated peptides
5 concerning permissible amide mimic substitutions.

In another approach, a variety of uncoded or modified amino acids such as D-amino acids and N-methyl amino acids have been used to modify mammalian peptides. Alternatively, a presumed bioactive conformation has been stabilized by a
10 covalent modification, such as cyclization or by incorporation of γ -lactam or other types of bridges. See, e.g., Veber and Hirschmann, et al., *Proc. Natl. Acad. Sci. USA*, 1978 75 2636 and Thorsett, et al., *Biochem Biophys. Res. Comm.*, 1983 111 166. The primary purpose of such manipulations has not been
15 to avoid metabolism or to enhance oral bioavailability but rather to constrain a bioactive conformation to enhance potency or to induce greater specificity for a receptor subtype.


Another approach, disclosed by Rich, D.H. in *Protease Inhibitors*, Barrett and Selveson, eds., Elsevier (1986), has
20 been to design peptide mimics through the application of the transition state analog concept in enzyme inhibitor design. For example, it is known that the secondary alcohol of statine mimics the tetrahedral transition state of the scissile amide bond of the pepsin substrate. Again, increased potency rather
25 than decreased susceptibility to peptidases or increased bioavailability was the principal objective. Moreover, the transition state analog concept has no apparent relevance to hormone agonist/antagonist design.

Nicolaou and Hirschmann, et al., *Design and synthesis of a peptidomimetic employing β -D-glucose for scaffolding*, in
30 *Peptides*, Rivier and Marshall, eds., ESCOM (1990), disclosed non-peptide somatostatin mimics having structures (1) and (2), wherein Bn is benzyl.

- 5 -



(1)



(2)

These mimics bound somatostatin receptors of AtT-20 cells with IC_{50} of about 9.5×10^{-6} M and about 1×10^{-6} M, respectively, compared with an IC_{50} of about 9.3 nM (9.3×10^{-9} M) for somatostatin itself. Significantly, the mimics failed to bind other G-protein-linked receptors at clinically acceptable concentrations. For example, while it was found that the β -adrenergic receptor, which is also found in AtT-20 cells, bound mimic (1), it required a five fold higher concentration to do so than was required for the somatostatin receptor. The goal of the authors was to increase the specificity of the mimics for the somatostatin receptor, not to develop compounds which would be bound by G-protein-linked receptors. Indeed, the authors suggested increasing the potency of the compounds as a means for enhancing this specificity.

Accordingly, there remains a long-felt need for metabolically stable chemical compounds which exhibit both good bioavailability and the capacity to bind a variety of G-protein-linked receptors.

20 OBJECTS OF THE INVENTION

It is one object of the present invention to provide compositions of matter which mimic or inhibit the biological and/or chemical activity of peptides.

It is another object to provide compositions which are chemically more stable than naturally-occurring peptides, particularly under conditions such as found in the human body.

It is a further object to provide compositions which function as hormone agonists or hormone antagonists.


- 6 -

It is a further object to provide compositions which effectively bind G-protein-linked receptors, especially the substance P receptor.

It is still a further object to provide prophylactic, diagnostic, and therapeutic uses for peptides and peptide analogs.


SUMMARY OF THE INVENTION:

These and other objects are accomplished by the present invention, which provides compounds, known as peptide analogs, which contain no peptide bonds yet which mimic or inhibit the chemical and/or biological activity of peptides. In general, the peptide analogs of the invention have structure (3):




(3)

wherein at least one of R_1 , R_2 , R_3 , R_4 , or R_5 comprises a chemical functional group which causes the compounds to be crossreactive with the peptide of interest. In preferred embodiments, peptide analogs of the invention have the structure (4) and, more preferably, the structure (5):



(4)




(5)

- 7 -

Compounds having these structures have been found to effectively bind a number of G-protein-linked receptors. Indeed, it has even been discovered in accordance with the present invention that compounds having structures (1) and (2) are able to bind G-protein-linked receptors other than the SRIF receptor.

The peptide analogs of the invention can be employed to mediate the chemical and/or biological effects of hormone agonists/antagonists or other peptides. These compounds are believed to possess beneficial properties such as increased half-life, lack of immunogenicity, and the ability to cross the blood-brain barrier; they are believed to be useful for the development of pharmaceutical, therapeutic, and diagnostic techniques.

In another aspect, the present invention provides cyclic hexapeptides having structure:



wherein:

R_{10} is indolyl;

R_{11} is H, isopropyl, phenyl, 4-hydroxyphenyl, 4-methoxyphenyl, or fluorophenyl;

R_{12} is phenyl; and

R_{13} is -OH, -C(O)OH, -H, -indolyl, -phenyl, -CH₂-phenyl, -cyclohexyl, or -naphthyl.

The invention also provides methods for producing a prophylactic or therapeutic response in a mammal by administering to the mammal a pharmaceutically effective amount of one or more compounds. In accordance with preferred embodiments, the present invention provides methods for producing such responses by modulating the activity of at

- 8 -

least one mammalian G-protein-linked receptor by administering an effective amount of one or more such compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1a-c depict synthetic schemes for the imidazol
5 compounds of Example 9.

Figures 2a-d depict synthetic schemes for the ester compounds of Example 10.

Figures 3a-f depict synthetic schemes for the compounds of Example 11.

10 DETAILED DESCRIPTION OF THE INVENTION

It has been found in accordance with the present invention that non-peptide compounds which mimic or inhibit the chemical and/or biological activity of a variety of peptides can be produced by appending to certain core species
15 such as the tetrahydropyranyl ring of structure (3) chemical functional groups which cause the compounds to be at least partially crossreactive with the peptide. As will be recognized, compounds which mimic or inhibit peptides are to varying degrees crossreactive therewith. In accordance with
20 the present invention, crossreactive moieties are those which compete with one another in binding G-protein-linked receptors through one of the many chemical reaction phenomena known in the art such as, for example, complexation, crystallization, or ionic, hydrogen, or covalent bonding. Thus, it is intended
25 that the term "crossreactive" include both agonism and antagonism. Those skilled in the art recognize that a substance which competes with a G-protein in binding to a cell receptor is described as an agonist if the response of the cell is the same as or mimics the action of the peptide
30 ligand. A substance that competes with the G-protein in binding to a receptor is referred to as antagonist if it blocks or inhibits the action of the cell to the action of the G-protein.

There exist a wide variety of useful analytical
35 techniques for elucidating the precise structure of a peptide.


- 9 -

These techniques include amino acid sequencing, x-ray crystallography, mass spectroscopy, nuclear magnetic resonance spectroscopy, computer-assisted molecular modeling, peptide mapping, and combinations thereof. Structural analysis of a peptide generally provides a large body of data which in preferred embodiments comprises the amino acid sequence of the peptide as well as the three-dimensional positioning of its atomic components. It is believed that only certain of these components, which are known both individually and collectively as chemical functionality, participate in any given reaction phenomena. It will be appreciated that the participation of a chemical functional group in peptide reactivity is manifested by the linkage or coordination of the functional group with at least a portion of a complementary reactive moiety such as a hormone receptor. Such linkage or binding may be effected through a covalent, ionic, or hydrogen bond or some weaker atomic coordination effect such as complexation or crystallization.

In accordance with the present invention, peptide chemical functionality which participates in binding is identified by one of the many techniques known in the art. For example, such identification can be effected through a stepwise process wherein one or more peptide analogs are prepared. For example, peptide analogs having structure (3) can be prepared by substitution at certain of the positions R_1 - R_n with chemical functionalities which are crossreactive with functionalities found in the peptide. The activity of the analog in a binding assay is then compared with that of the peptide. The degree to which the binding of the analog corresponds with that of the peptide indicates the degree to which the substituents participate in the binding phenomena. Accordingly, one important criterion in preparing peptide analogs according to the present invention is the respective chemical similarity of the side chains found in the peptide and any potential substitutes therefor appended to the core structure in the analog. In general, it is desired that the chemical functional group in the peptide of interest and its


- 10 -

substitute in at least one of the peptide analogs be somewhat chemically dissimilar. Where the substitute is chemically dissimilar from the peptide side chain, it will generally be easier to elucidate the contribution, if any, of side chain to activity of the peptide. For example, it is believed that the bioactive conformation of somatostatin (also known as somatotropin release inhibiting factor or SRIF) includes a β -turn involving residues 7-10 (Phe⁷-Trp⁸-Lys⁹-Thr¹⁰). These four amino acids have been shown to be necessary and sufficient for receptor recognition and activation, so long as they are held in the proper orientation. Somatostatin accomplishes this proper orientation through its ten remaining amino acids and the cystine bridge contained therein. In a number of active cyclic hexapeptide analogs for somatostatin, proper orientation of the four amino acids is maintained via dipeptide segments. For example, the cyclic hexapeptide L-363,301 (structure (6a)), disclosed by Veber and Hirschmann, et al., *Life Sciences*, 1984, 34, 1371 and the cyclic hexapeptide MK-678 (structure (6b)), disclosed by Veber and Hirschmann, et al., *Nature*, 1981, 292, accomplish the proper orientation via the segments Phe-N-Me-Ala or Phe-Pro, respectively.



(6a)

- 11 -



(6b)

It is believed that the solution conformation of somatostatin involves a type I β -turn for residues 7-10 and that of the significantly more potent D-TRP diastereomer involves a type II' β -turn. While these two turns differ in the Φ and Ψ angles of the amide backbone, they are believed to assume similar orientations of the side chains at the receptor. In the cyclic hexapeptides, the Phe-N-Me-Ala sequence and the Phe-Pro sequence are believed to be part of a type VI β -turn. Of particular significance is the high activity found for a modified retro-enantiomeric cyclic hexapeptide wherein the amide backbone is reversed. This demonstrates that proper side chain topography is important for activity but that the amide backbone may not be.

In accordance with the present invention, peptide analogs having structure (3) were further simplified by including only three adjacent side chains of the four amino acids of the β -turn. These side chains are attached to rigid frameworks devoid of peptide bonds. The frameworks were developed through molecular modeling to orient the side chains appropriately and/or to permit the receptor to induce the proper fit.


While a proper β -turn requires the fourth amino acid of the β -turn -- Thr in somatostatin and several cyclic

- 12 -

hexapeptides and Val in the superactive cyclic hexapeptide -- it is believed that neither the Thr nor the Val side chains are required for binding. This assumption is based on the fact that highly active somatostatin analogs are known which have either Val, Thr, Ser, α -aminobutyric acid, or Gly in the fourth position of the β -turn. Such non-specificity suggests a conformational rather than a binding role for that amino acid of the β -turn.

The phenylalanine residue in the dipeptide segments Phe-N-Me-Ala or Phe-Pro appears to add an important hydrophobic binding element. For this reason, the present synthetic analogs of somatostatin contain a corresponding aromatic residue. Increased hydrophobicity also should prove helpful in improving the duration of action and activity via oral administration of such compounds.

It is now believed that for the L-363,301 hexapeptide, structure (6a), the β -turn is important and the three groups extending from carbons a, b, and c -- benzyl, indole, and alkylamino, respectively -- are necessary for binding. Whereas the substituent at carbon d appears to be required to stabilize the β -turn rather than be required for binding, a benzyl group attached at carbon e of the skeleton is believed to be an important binding ligand which improves the activity of analogs. It has now been discovered that a new class of therapeutic agents can be formulated having activity in a broad spectrum of utilities, especially those related to the G-protein-linked receptors. One member of the class is represented by structure (7).







- 13 -

The calculated bond distances for structure (7) and the cyclic hexapeptide suggest close geometrical similarities. Furthermore, overlaying models of the designed structure (7) and the cyclic hexapeptide (6a) shows close correspondence of the important functionalities, particularly the phenylalanine, tryptophan and lysine residues.


The present invention, however, is not limited to embodiments wherein benzyl, indole, or alkylamino groups participate in binding. Participatory chemical functionality according to the present invention includes any of the wide variety of functional groups known in the art. The side chains of naturally-occurring amino acids provide examples of suitable participatory functionality. Representative participatory chemical functionality which may be contained within groups R_1 - R_5 is set forth in Table 1. For example, one or more of R_1 - R_5 can have the structure $Z-(CH_2)_y-$ or $Z-O-$, where y is from 0 to about 5 and Z is one of the side chains of Table 1.

- 14 -

TABLE 1

CH ₃ -	CH ₃ -CH ₂ -S-CH ₂ -CH ₂ -HO-CH ₂ -CH ₂ -
HO-CH ₂ -	CH ₃ -CH ₂ (OH)-
C ₆ H ₅ -CH ₂ -	HO ₂ C-CH ₂ -NH ₂ C(O)-CH ₂ -
5 HO-C ₆ H ₅ -CH ₂ -	
	HCO ₂ -CH ₂ -CH ₂ -
	NH ₂ C(O)-CH ₂ -CH ₂ -
	(CH ₃) ₂ -CH-
HS-CH ₂ -	(CH ₃) ₂ -CH-CH ₂ -
HO ₂ C-CH(NH ₂)-CH ₂ -S-S-CH ₂ -	CH ₃ -CH ₂ -CH ₂ -
CH ₃ -CH ₂ -	H ₂ N-CH ₂ -CH ₂ -CH ₂ -
CH ₃ -S-CH ₂ -CH ₂ -	H ₂ N-C(NH)-NH-CH ₂ -CH ₂ -CH ₂ -
	H ₂ N-C(O)-NH-CH ₂ -CH ₂ -CH ₂ -
	CH ₃ -CH ₂ -CH(CH ₃)-
	CH ₃ -CH ₂ -CH ₂ -CH ₂ -
	H ₂ N-CH ₃ -CH ₂ -CH ₂ -CH ₂ -

- 10 In accordance with the present invention, non-peptide analogs preferably possess the general structure (3):



(3)

wherein:

- 15 R₁ is -O(CH₂)_nR_A, -OC(O)(CH₂)_nR_A, -(CH₂)_nR_A, or -C(O)(CH₂)_nR_A where R_A is -H, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about

- 15 -

14 carbon atoms and up to about 4 nitrogen atoms, and n is an integer from 0 to about 12;

at least one of R_2 , R_3 , and R_4 , independently, is $-O(CH_2)_mR_B$, $-OC(O)(CH_2)_mR_B$, $-(CH_2)_mR_B$ or $-C(O)(CH_2)_mR_B$ where R_B is $-H$ or aryl, and m is an integer from 0 to about 5; and

R_5 is $-O(CH_2)_pNHR_C$, $-OC(O)(CH_2)_pNHR_C$, $-O(CH_2)_pR_D$, $-OC(O)(CH_2)_pR_D$, $-(CH_2)_pNHR_C$, $-C(O)(CH_2)_pNHR_C$, $-(CH_2)_pR_D$ or

$-C(O)(CH_2)_pR_D$, where:

p is an integer from 0 to about 10;

R_C is $-R_E$ or $-C(O)R_E$;

R_D is $-H$, $-OR_E$, or $-C(O)R_E$;

R_E is $-H$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms; or a pharmaceutically acceptable salt thereof.



It will be understood that the terms "alkyl" and "alkenyl" as employed herein are intended to include cyclic as well as straight chain moieties, including methyl, tert-butyl groups, fluoroethyl, and vinyl groups. Preferred alkyl groups have 1 to about 14 carbon atoms, and preferred alkenyl groups have 2 to about 14 carbon atoms. Aryl groups according to the invention are aromatic and substituted aromatic groups having 6 to about 14 carbon atoms, including phenyl, fluorophenyl, benzyl, imidazolyl, indolyl, and naphthyl groups. In certain embodiments, the chemical structure and stereochemistry of the peptide analogs of the invention roughly correspond to that of β -D-glucose. Hence, the analogs can possess structures (4) and (5), with R_1 - R_5 defined as above.

As will be recognized, the precise identity of R_1 - R_5 depends intimately upon the peptide of interest whose biological and/or chemical activity is to be mimicked or

- 16 -


inhibited. For example, in the case of compounds which are bound by G-protein-linked receptors such as the substance P receptor, R_A should be an aryl functional group, preferably an nitrogen-substituted aryl group such as pyridine or indole.

- 5 More preferably, R_A is a 3-substituted indole. For such compounds, n should be 2 and R_B should be phenyl. The integer m should be zero or, preferably, 1. Also, R_5 should be $-O(CH_2)_pNH_2$ or $-O(CH_2)_pNHR_C$, where p is from about 2 to about 8, preferably 3 to about 6, more preferably 5. R_C can be, for
 10 example, a phenyl, benzyl or nitrogen heterocyclic moiety. Where substitution is possible at more than one position of these and other R_C , it is intended that the present invention include each of the resulting peptide analogs. For example, it is intended that the invention include analogs wherein R_C
 15 is a pyridine or isonicotinic acid residue having one of the following structures:




Preferably, however, R_C is $-CH_3$.

- In general, preferred peptide analogs have structures
 20 (8) - (13).




(8)




(9)


- 17 -




(10)



(11)




(12)



(13)

Also preferred are compounds having formula (3) wherein: (a) R_1 is $O-(CH_2)_2$ -indolyl, R_2 is $O-CH_2$ -fluorophenyl, R_3 and R_4 are O -benzyl, and R_5 is $O-CH_2$ -naphthyl; and (b) R_1 is $O-(CH_2)_2$ -indolyl, R_2 is $O-CH_2$ -naphthyl, R_3 and R_4 are O -benzyl, and R_5 is $O-CH_2$ -fluorophenyl. These peptide analogs are preferred to the extent that they selectively and effectively bind G-protein-linked receptors such as the somatostatin receptor, the β -adrenergic receptor, and the substance P receptor.

In another aspect, the present invention also provides cyclic hexapeptides having structure:



- 18 -

wherein:

R_{10} is indolyl;

R_{11} is H, isopropyl, phenyl, 4-hydroxyphenyl, 4-methoxyphenyl, or fluorophenyl;

5 R_{12} is phenyl; and

R_{13} is -OH, -C(O)OH, -H, -indolyl, -phenyl, -CH₂-phenyl, -cyclohexyl, or -naphthyl.

In preferred embodiments, these cyclic hexapeptides also bind G-protein-linked receptors selectively and effectively.

10 It will be recognized that the degree to which a compound binds a receptor is known as its binding activity or potency. The potency of a compound commonly is expressed as its inhibitory concentration (IC), the concentration at which the compound is able to displace a predetermined portion --
15 typically 50% -- of another compound which is already bound to a particular receptor. In the case of ligand-binding studies, the compound that is displaced is a radioactive agonist or antagonist at the receptor under study. It is preferred in accordance with the present invention that a
20 peptide or peptide analog possess a clinically effective IC₅₀ in at least one mammal, that is, a concentration which is low enough to inhibit binding of radioactive agonist or antagonist of a given G-protein-linked receptor while causing a minimum of unacceptable side effects in the mammal. As will be
25 recognized, clinically effective inhibitory concentrations vary depending on a number of factors, such as the pharmacokinetic characteristics and stability of the compound under study and thus must be determined empirically for each analog and each factor. For example, the clinically effective
30 concentration for the human somatostatin receptor is about 50-500 nM, but for the *in vitro* system the potency is about 1-10 nM. In general, it is desired that the potency of a compound of the invention be as great as possible, preferably greater than or equal to the native hormone.

35 Selectivity or specificity is manifested for a compound of the present invention by its tendency to bind one particular G-protein-linked receptor but not other G-protein-

- 19 -

linked receptors. In an experimental context, selectivity is manifested where a compound is bound by a particular receptor when placed in contact or close proximity with a medium containing at least one other receptor. Typically, 5 specificity is expressed as a ratio of the potency or activity of a compound for two different receptors. Thus, a compound having an IC_{50} of 100 μM for compound A and IC_{50} of 200 μM for compound B can be said be two times more selective for compound A. In general, the selectivity of the peptides and 10 peptide analogs of the present invention should be as great as possible. Selectivities greater than about 50-100 fold are preferred and selectivities greater than about 500 fold even more preferred.

As can be seen, the present invention provides a wide 15 variety of peptides and peptide analogs which effectively and selectively are bound by individual G-protein-linked receptors. Those compounds which bear amino groups are capable of forming salts with various inorganic and organic acids and such salts are also within the scope of this 20 invention. Examples of such acid addition salts include acetate, adipate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, ethanesulfonate, fumarate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, methanesulfonate, 25 lactate, maleate, methanesulfonate, 2-napthalenesulfonate, nitrate, oxalate, pamoate, persulfate, picrate, pivalate, propionate, succinate, sulfate, tartrate, tosylate, and undecanoate. The salts may be formed by conventional means, such as by reacting the free base form of the product with one 30 or more equivalents of the appropriate acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is later removed *in vacuo* or by freeze drying. The salts also may be formed by exchanging the anions of an existing salt for another anion on a suitable ion exchange 35 resin.

The present invention also provides compositions which comprise one or more peptides or peptide analogs. To the

- 20 -

extent that the compositions comprise individual compounds which are bound by certain receptors, the compositions will likely also be bound by the same receptors. The compounds themselves may be present in the compositions in any of a wide
5 variety of forms. For example, two or more peptides or peptide analogs may be merely mixed together or may be more closely associated through complexation, crystallization, or ionic or covalent bonding.

Those skilled in the art will appreciate that a wide
10 variety of prophylactic, diagnostic, and therapeutic treatments may be prepared from the synthetic compounds and compositions of the invention, due in large part to the crossreactivity --that is, agonism or antagonism -- of these moieties with one or more naturally-occurring peptides. For
15 example, by administering an effective amount of a peptide or peptide analog, prophylactic or therapeutic responses can be produced in a human or some other type of mammal. Preferred responses are produced by modulating -- that is, increasing, decreasing or otherwise modifying -- the activity of at least
20 one G-protein-linked receptor. It will be appreciated that the production of prophylactic or therapeutic responses includes the initiation or enhancement of desirable responses, as well as the cessation or suppression of undesirable responses.

25 Certain preferred peptides and peptide analogs of the present invention exhibit significant substance P receptor-binding activity and therefore, are of value in the treatment of a wide variety of clinical conditions which are characterized by the presence of an excess of tachykinin, in
30 particular substance P, activity. These include disorders of the central nervous system such as anxiety, psychosis and schizophrenia; neurodegenerative disorders such as senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; respiratory diseases such as bronchospasm and
35 asthma; inflammatory diseases such as inflammatory bowel disease, osteoarthritis and rheumatoid arthritis; adverse immunological reactions such as rejection of transplanted

- 21 -

tissues; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease and incontinence; disorders of blood flow caused by vasodilation; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions or the transmission of pain in migraine. Hence, these compounds are readily adapted to therapeutic use as substance P antagonists for the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

Compositions for use in the methods of this invention can be in the form of a solid, semisolid or liquid form and can include one or more of peptides or peptide analogs as an active ingredient in a mixture with an organic or inorganic carrier or excipient suitable for external, enteral or parenteral applications. The active ingredient may be compounded, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The carriers which can be used are water, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form, and in addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. The active ingredient is included in the pharmaceutical composition in an amount sufficient to produce the desired effect upon the process or condition of diseases.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates, together with granulation binders

- 22 -

like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may
5 also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with
10 various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water ethanol, propylene glycol, glycerin and various like combinations thereof.

15 For parenteral administration, solutions of said compounds in either sesame or peanut oil in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH>8) if necessary and the liquid diluent first rendered isotonic. These aqueous
20 solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known
25 to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with
30 standard pharmaceutical practice.

A compound of the invention may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles.
35 The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques.

- 23 -

Dosage levels of the compounds within the present invention on the order from about 0.01 mg to about 50 mg per kilogram of body weight per day, preferably from about 0.1 mg to about 10 mg per kilogram body weight per day, are believed
5 to be useful in the treatment of the above-indicated conditions (i.e., from about 0.7 mg to about 3.5 g per patient per day, assuming a 70 kg patient). In addition, the compounds of the present invention may be administered on an intermittent basis; i.e. at semi-weekly, weekly, semi-monthly
10 or monthly intervals.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended
15 anisaldehyde solution (sugars), ninhydrin (primary amines), phosphomolybdic acid (secondary amines), or Erlich's reagent (incoles).

Flash column chromatography for Examples 1-11 was performed using Merck 60-200 mesh silica gel. All yields
20 reflect purified isolated product after flash column chromatography or recrystallization unless otherwise noted.

For Examples 12-17, unless otherwise noted, all solvents and reagents were obtained from commercial sources and used without further purification. Analytic reverse-phase
25 HPLC was carried out employing a LKB system (2152 LC controller, 2150 HPLC pump, 2141 variable wavelength monitor on a C18 Dynamax 300 (0.46-25 cm) column. Semi- or preparative reverse-phase HPLC separations were achieved using a Ranin solvent delivery system equipped with a Dynamax
30 detector (model UV-D) utilizing either C18 Dynamax 300 (21.4 x 250 mm) column or C8 Vydac column (10 x 250 mm). The mobile phase consisted of 0.1% TFA in water (buffer A) and 0.1% TFA in acetonitrile (buffer B). The FAB-mass spectra were obtained on a ZAB-E VG analytical spectrometer. 1H and 13C
35 NMR spectra were obtained with a Bruker AM500 spectrometer. Chemical shifts are reported in δ values relative to tetramethylsilane for proton and solvent for carbon spectra.

- 24 -

Optical rotation were measured on a Perkin-Elmer Model 241 polarimeter.

EXAMPLE 1

Preparation of Analog Having Structure (1), 2-(1H-Indol-3yl)ethyl-6-O-(5-aminopentyl)-2,3,4-tri-O-benzyl- β -D-glucopyranoside

A. 1-Bromo α -D glucose tetraacetate

Hydrobromic acid (30% in acetic acid, 11.85 ml, 55.4 mmol) was added to β -D-glucose pentaacetate (12.01 g, 30.8 mmol) at 0°C. After 10 minutes, the resulting solution was warmed to room temperature and stirred for 4 hours. The reaction mixture was slowly poured, with stirring, into ice water (250 ml) and was stirred until the product solidified. The product was collected by vacuum filtration and washed with cold water. The white solid was dissolved in carbon tetrachloride (60 ml) and washed with H₂O (1 x 20 ml), saturated aqueous NaHCO₃ (3 x 20 ml, until pH = 7), H₂O (1 x 20 ml), dried with CaCl₂, and poured into cold petroleum ether (250 ml). After 30 min, the crystalline product was collected by vacuum filtration to give the target compound as a white solid (10.0 g, 80%).

B. N-phenylsulfonyl tryptophol**(a) 1-O-tert-butyldimethylsilyl-2-3-indolyl)ethanol**

To a solution of tryptophol (5.0 g, 31 mmol) in dimethylformamide (DMF, 30 ml) was added imidazole (4.64 g, 68 mmol) and the reaction cooled to 0°C. To the cooled solution was added tert-butyldimethylsilyl chloride (5.14 g, 34.1 mmol) and the reaction was stirred at room temperature overnight. The reaction was diluted with ethyl acetate (100 ml) and extracted with water (2 x 100 ml). The aqueous layer was extracted with ethyl acetate (1 x 200 ml.) The organic layers were combined and dried over anhydrous sodium sulfate. The solvents were removed under reduced pressure to yield a pale orange oil. Purification by flash column chromatography using 30% ether in petroleum ether yielded the target compound as a colorless oil (8.43 g, 99%).

- 25 -

(b) 1-O-tert-butyldimethylsilyl-2-[3-(1-N-phenylsulfonyl)indolyl]ethanol

Sodium hydride (1.91 g, 60% oil dispersion) was placed in a flame dried flask under argon. Dry DMF (64 ml) was added and the suspension cooled to 0°C. A solution of 1-O-tert-butyldimethylsilyl-2-3-indolyl]ethanol (8.43 g, 30.6 mmol) in dry DMF (30 ml) was added to the suspension and the reaction stirred to room temperature for 30 minutes. After cooling to 0°C, benzenesulfonyl chloride (5.30 ml, 39.7 mmol) was added dropwise. The reaction was stirred at room temperature overnight. A solution of ammonium chloride (100 ml) was added and the reaction was extracted with ether (3 x 200 ml). The organic layers were combined, extracted with saturated sodium chloride, and dried over anhydrous sodium sulfate. Removal of the solvents under reduced pressure yield a pale yellow oil. Purification by flash column chromatography using 30% ether in petroleum ether yielded the target compound as a colorless oil (7.37 g, 79%).

(c) N-phenylsulfonyl tryptophol

To a solution of 1-O-tert-butyldimethylsilyl-2-[3-(1-N-phenylsulfonyl)indolyl]ethanol (6.6 g, 21.9 mmol) in tetrahydrofuran (THF, 100 ml) was added tetrabutylammonium fluoride (21 ml, 1 M in THF) and the solution stirred at room temperature overnight. The reaction was diluted with ethyl acetate (100 ml) and extracted with water (2 x 100 ml). The organic layer was re-extracted with saturated sodium chloride solution, dried over anhydrous sodium sulfate and the solvents removed under reduced pressure to yield a pale yellow oil. Purification by flash column chromatography using 40% ethyl acetate in petroleum ether yielded the target compound as a pale yellow oil which crystallized upon standing (4.00 g, 84%).

C. 2-(1-Phenylsulfonyl-3-yl)ethyl-2,3,4,6-tetra-O-acetyl- β -D-glucopyranoside

To a suspension of crushed, flame dried 4 Å sieves (0.89 g) and silver (I) oxide (412 mg, 17.8 mmol) in 9 ml of dry hexane at room temperature, was added a solution of the

- 26 -

above N-phenyl sulfonyl tryptophol (537 mg, 1.78 mmol) in 3 ml of dry benzene followed by a solution of 1-bromo α -D glucose tetraacetate (804 mg, 1.95 mmol) in 3 ml of dry benzene. The reaction vessel was covered with aluminum foil and allowed to stir for 2 days at room temperature. Thin layer chromatography (TLC, 5% ether in methylene chloride) revealed product and some unchanged starting material. Silver (I) oxide (206 mg, 8.9 mmol) was added followed by 1 ml of dry benzene to loosen the suspension. The reaction was allowed to stir at room temperature an additional 2 days. The reaction suspension was filtered through celite. Concentration and crystallization from ethyl acetate/petroleum ether afforded 580 mg of the β -isomer of the target compound as a white solid. Concentration of the filtrate and flash chromatography (silica, 5% ether in methylene chloride) afforded a mixture of the β -isomer along with the α -isomer and the corresponding ortho ester. Flash chromatography (silica, 70% ether in petroleum ether) on the mixture afforded an additional 134 mg of the β -isomer, bringing the yield to 64% (716 mg).

D. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl- β -D-glucopyranoside

Sodium methoxide (221 mg, 4.09 mmol) was added to a suspension of 2-(1-phenylsulfonyl-3-yl)ethyl-2,3,4,6-tetra-O-acetyl- β -D-glucopyranoside (3.22 g, 5.12 mmol) in 26 ml of methanol at room temperature. After 20 minutes, the resulting solution was diluted with 26 ml of methanol and neutralized by addition of amberlyst H⁺ resin. The resin was quickly removed by filtration to avoid formation of the methyl glucoside. Concentration of the filtrate and flash chromatography (silica, 5:1:1 methylene chloride, methanol, acetone) afforded the target compound (2.09 g, 88%) as a white foam.

E. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-6-O-tert-butyl-diphenylsilyl- β -D-glucopyranoside

To a stirred solution of 2-(1-Phenylsulfonyl-indol-3-yl)ethyl- β -D-glucopyranoside (7.11 g, 15.4 mmol) in 51 ml of dry DMF was added at room temperature, imidazole (2.93 g, 43.1

- 27 -

mmol) followed by tert-butyldiphenylsilyl chloride (5.58 g, 21.6 mmol). The solution was maintained at 50°C for 24 hours. After removal of the DMF under reduced pressure, the reaction mixture was diluted with 250 ml of ethyl acetate and washed
5 with H₂O (1 x 100 ml), saturated aqueous NaCl (1 x 100 ml), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 5% methanol in dichloromethane) provided pure target compound (9.15 g, 85%) as a white foam.

10 **F. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-2,3,4-tri-O-benzyl-6-O-tert-butyldiphenyl-silyl-β-D-glucopyranoside**

To a stirred suspension of sodium hydride (323 mg, 60% oil dispersion, 808 mmol) in 5 ml of dry THF at 0°C was added a solution of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-6-O-tert-
15 butyldiphenylsilyl-β-D-glucopyranoside (1.62 g, 2.31 mmol) in 7 ml dry THF. After stirring 1 hour at room temperature, benzyl bromide (1.09 ml, 9.24 mmol) was added dropwise to the reaction mixture at 0°C followed by tetrabutylammonium iodide (85 mg, 0.23 mmol). After stirring 3 days at room
20 temperature, the suspension was treated with 3 ml of saturated aqueous ammonium chloride at 0°C. The resulting solution was diluted with 80 ml of ether and washed with saturated aqueous NH₄Cl (1 x 30 ml), saturated aqueous NaCl (1 x 30 ml) and dried over magnesium sulfate. Concentration and flash
25 chromatography (silica, 20% ether in petroleum ether) afforded the target compound (1.66 g, 74%) as a white foam.

G. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-2,3,4 tri-O-benzyl-β-D-glucopyranoside

To a stirred solution of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-2,3,4-tri-O-benzyl-6-O-tert-butyldiphenyl-silyl-β-D-
30 glucopyranoside (1.55 g, 1.60 mmol) in 8 ml of dry THF at room temperature was added tetrabutylammonium fluoride (1 M in THF, 2.4 ml, 2.4 mmol). After stirring 7 hours, the solution was diluted with 70 ml of ethyl acetate and washed with H₂O (1 x
35 30 ml) and saturated aqueous NaCl (1 x 30 ml) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 30% ethyl acetate in petroleum ether) afforded the

- 28 -

target compound (1.10 g, 94%) as a clear oil: R_f 0.50 (40% ethyl acetate in petroleum ether); 1H NMR (500 MHz, $CDCl_3$) δ 7.84 (d, J = 8.3 Hz, 1H), 7.82 (d, J = 7.9 Hz, 2H), 7.53 (s, 1H), 7.48-7.17 (m, 21H), 4.92 (d, J = 11.0 Hz, 1H), 4.86 (d, J = 10.9 Hz, 1H), 4.81 (d, J = 11.0 Hz, 1H), 4.74 (d, J = 11.0 Hz, 1H), 4.62 (d, J = 11.0 Hz, 1H), 4.48 (d, J = 7.8 Hz, 1H), 4.20 (ddd, J = 9.4, 7.0, 7.0 Hz, 1H), 3.91-3.86 (m, 2H), 3.73 (dd, J = 3.5, 11.9 Hz, 1H), 3.63 (ddd, J = 9.0, 9.0, 18.0 Hz, 2H), 3.40 (ap. t, J = 8.0 Hz, 1H), 3.35 (ddd, J = 9.4, 4.2, 2.6 Hz, 1H), 3.04-2.93 (m, 2H), 2.06 (s, 1H); ^{13}C NMR (500 MHz, $CDCl_3$) δ 138.48, 138.21, 138.13, 137.95, 135.09, 133.60, 130.92, 129.10, 128.40, 128.30, 128.25, 128.22, 127.98, 127.90, 127.82, 127.76, 127.55, 126.58, 124.72, 123.57, 123.12, 119.61, 119.31, 113.66, 103.59, 84.39, 82.25, 77.37, 75.56, 75.16, 74.99, 74.75, 68.60, 61.77, 25.57; IR (thin film) 3480 (w), 3065 (w), 3035 (w), 2920 (m), 2878 (m), 1498 (w), 1450 (s), 1365 (s), 1280 (w), 1220 (m), 1176 (s), 1123 (s), 1090 (s), 1073 (s), 1030 (s), 750 (s), 700 (s) cm^{-1} ; UV-Vis (c = 9.21×10^{-5} , acetonitrile) λ_{max} 254.0 (ϵ = 2.81×10^3), 211.6 (ϵ = 3.19×10^4) nm; HRMS m/e calculated for $C_{43}H_{43}NO_8S$ ($M+H$): 734.2774, found 734.2743; $[\alpha]_D^{20}$ -13.3° (c = 0.135, acetonitrile); Analysis calculated for $C_{43}H_{43}NO_8S$: C, 70.37; H, 5.91; found: C, 70.30; H, 6.08.

H. 2-(1-phenylsulfonyl-indol-3yl)ethyl-2,3,4-tri-O-benzyl-6-O-trifluoromethylsulfonyl- β -D-glucopyranoside

To a stirred solution of 2-(1-phenylsulfonyl-indol-3yl)ethyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside (196 mg, 0.27 mmol) in 2.7 mL of dry dichloromethane at -78°C was added 2,6-di-tert-butyl-4-methyl pyridine (880 mg, 0.427 mmol) followed by triflic anhydride (58 μ l, 0.347 mmol). After stirring 15 minutes at -78°C, the mixture was warmed to room temperature over 20 minutes, and then poured into saturated aqueous $NaHCO_3$ (20 mL) and extracted with ethyl acetate (60 mL). The organic layer was washed with saturated aqueous $NaHCO_3$ (3 x 20 mL), saturated aqueous $NaCl$ (1 x 20 mL) and dried over magnesium

- 29 -

sulfate. Concentration provided the crude triflate target compound, which used in the next step without purification.

I. N-trifluoroacetyl-5-amino pentanol

To a solution of 5-amino pentanol (1 g, 9.69 mmol) in
5 methanol (25 ml, 0.4 M) at 0°C was added triethylamine (2 ml, 1.5 equiv, 10 mmol) followed by very slow dropwise addition of trifluoroacetic anhydride (1.8 ml, 1.3 equiv, 12.5 mmol). The reaction mixture was warmed to room temperature and stirred overnight. TLC (5% CH₃OH/CH₂Cl₂) stained with
10 ninhydrin revealed starting material; TLC stained with PMA revealed product. The reaction mixture was cooled to 0°C and triethylamine (1.3 ml, 1 equiv. 9.69 mmol) was added followed by trifluoroacetic anhydride (1 ml, 0.8 equiv.). The reaction mixture was warmed to room temperature and stirred an
15 additional night. Concentration and flash chromatography (silica, 60% EtOAc/petroleum ether) afforded the target compound (1.7 g, 85%).

J. 2-(1-Phenylsulfonyl-indol-3yl)ethyl-2,3,4 tri-O-benzyl-6-O-(N-trifluoroacetyl-5-aminopentyl)-β-D-glucopyranoside

To a stirred suspension of sodium hydride (123 mg, 0.307 mmol, 60% oil dispersion) in 17 mL of dry THF at 0°C was added a solution of N-trifluoroacetyl-5-amino pentanol (265 mg, 1.3 mmol) in 10 mL of dry THF. After stirring 10 minutes
25 at 0°C, the suspension was warmed to room temperature, stirred for 1 hours, and cooled to 0°C. A solution of the above 2-(1-phenylsulfonyl-indol-3yl)ethyl-2,3,4-tri-O-benzyl-6-O-trifluoromethylsulfonyl-β-D-glucopyranoside (theoretically 0.27 mmol) in 16 ml of dry dichloromethane was added slowly
30 dropwise. The reaction mixture was stirred at 0°C for 30 minutes and then warmed to room temperature. After stirring 24 hours, TLC (2% methanol in dichloromethane) revealed diprotected target compound and a minor amount of monoprotected product. The reaction mixture was cooled to 0°C
35 and quenched with 10 mL of saturated aqueous ammonium chloride. The resulting mixture was diluted with ethyl acetate (150 mL) and washed with H₂O (1 x 50 mL), saturated

- 30 -

aqueous NaCl (1 x 50 mL) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 2% methanol in dichloromethane) yielded a mixture of diprotected target compound and monoprotected product which was used as a mixture
5 in the next step.

K. Structure (1), 2-(1-Phenylsulfonyl-indol-3yl)ethyl-6-O-(5-aminopentyl)-2,3,4-tri-O-benzyl- β -D-glucopyranoside

To a stirred solution of the mixture of step J, above,
10 (theoretically 0.27 mmol) in 6 mL of ethanol at room temperature was added a solution of 5M NaOH (2 mL, 10 mmol). The solution was heated to reflux for 2 hours. The solvents were removed under reduced pressure. The reaction mixture was diluted with ethyl acetate (40 mL) and washed with H₂O (1 x 15
15 mL), saturated aqueous NaCl (1 x 15 mL) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 5% methanol in dichloromethane provided structure (1) (150 mg, 83% for 3 steps) as an oil: R_f 0.26 (7% methanol in dichloromethane); ¹H NMR (500 MHz, CDCl₃) δ 7.98 (s, 1H), 7.59
20 (d, J = 7.9 Hz, 1H), 7.33-7.04 (m, 19H), 4.90 (d, J = 10.9 Hz, 1H), 4.85 (d, J = 11.1 Hz, 1H), 4.80 (d, J = 11.0 Hz, 1H), 4.77 (d, J = 10.9 Hz, 1H), 4.64 (d, J = 11.0 Hz, 1H), 4.60 (d, J = 11.1 Hz, 1H), 4.48 (d, J = 7.8 Hz, 1H), 4.21 (ddd, J =
9.4, 6.7, 6.7 Hz, 1H), 3.89 (ddd, J = 9.4, 7.3, 7.3 Hz, 1H),
25 3.64 (dd, J = 9.0, 9.0 Hz, 1H), 3.56 (t, J = 6.4 Hz, 2H), 3.51-3.47 (m, 1H), 3.42 (t, J = 9.2 Hz, 2H), 3.11 (t, 7.0 Hz, 2H), 2.96 (dd, J = 12.3, 2.6 Hz 1H), 2.66 (dd, J = 12.3, 7.8 Hz, 1H), 2.62-2.54 (m, 2H), 1.93 (s, 2H), 1.54-1.44 (m, 4H),
30 1.38-1.32 (m, 2H); ¹³C NMR (500 MHz, CDCl₃) δ 138.57, 138.49, 138.14, 136.17, 128.43, 128.36, 128.29, 128.02, 127.88, 127.82, 127.60, 127.56, 127.50, 122.14, 121.96, 119.30, 118.68, 112.60, 111.13, 103.67, 84.61, 82.45, 79.70, 77.20, 75.68, 74.99, 74.73, 73.82, 70.25, 62.63, 50.52, 49.59, 32.36, 29.28, 25.86, 23.31; IR (thin film) 3420(w), 3300(w), 3063(w),
35 3033(w), 2938(m), 2860(m), 1495(w), 1455(m), 1360(m), 1210(w), 1072(s), 1026(m), 910(w), 538(s), 495(s) cm⁻¹; UV-Vis (c=1.14x10⁻⁴, acetonitrile) λ_{max} 289.6 (e=4.17 x 10³), 280.8

- 31 -

($\epsilon=4.97 \times 10^3$), 220.0 ($\epsilon=2.4 \times 10^4$) nm; HRMS m/e calc'd $C_{42}H_{50}N_2O_6$ (M + H): 679.373, found 679.370; $[\alpha]D^{20}+3.2^\circ$ (c=0.31, acetonitrile).

EXAMPLE 2

5 **Preparation of Analog Having Structure (7), 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-6-O-(5-acetamidopentyl)-2,3,4-tri-O-benzyl- β -D-glucopyranoside**

To a solution of 5-amino pentanol (0.75 g, 7.27 mmol) in methanol (15 ml, 0.5 M) at 0°C was added triethylamine
10 (1.62 ml, 1.6 equiv, 11.6 mmol) followed by acetic anhydride (0.891 ml, 1.3 equiv, 9.45 mmol). The reaction mixture was warmed to room temperature and stirred overnight. TLC (8% CH_3OH/CH_2Cl_2) stained with ninhydrin revealed starting material. Triethylamine (1.6 ml, 1.6 equiv, 11.6 mmol) was
15 added to room temperature followed by acetic anhydride (0.9 ml, 1.3 equiv, 9.45 mmol) and the reaction mixture was stirred an additional night. Concentration and flash chromatography (silica, 7% $CH_3OH/EtOAc$) afforded N- CH_3CO -5-amino-pentanol (1 g, 100%).

20 Sodium hydride (0.108 g, 60% suspension in oil, 0.307 mmol, 2.2 equiv. compared to N- CH_3CO -5-amino-pentanol) was quickly weigh into a flame dried flask under argon. THF (20 ml, 0.01 M compared to moles of the triflate was added and the resulting suspension was cooled to 0°C. A solution of N-
25 CH_3CO -5-amino-pentanol (0.108 g, 0.22 moles, 5 equiv) in 5 ml of THF was added dropwise and then warmed to room temperature for 1 hour. The resulting suspension was cooled to 0°C and a solution of the 2-(1-phenylsulfonyl-indol-3-yl)ethyl-2,3,4 tri-O-benzyl- β -D-glucopyranoside triflate prepared in Example
30 1H (assumed 0.245 mmol) in CH_2Cl_2 (15 ml, CH_2Cl_2 :THF=3:5) was added slowly dropwise and stirred for 1 hour. The reaction mixture was warmed to room temperature and stirred overnight. TLC (3% CH_3OH/CH_2Cl_2) revealed no starting material and a major and minor product very close in R_f . Both were collected since
35 the minor product is deprotected indole and the mixture is transformed to the same product in the next step. The

- 32 -

reaction mixture was cooled to 0°C and quenched with aqueous saturated ammonium chloride. The reaction mixture was poured into EtOAc and washed 1 x H₂O and 1 x aqueous saturated NaCl. The organic layer was dried with MgSO₄ and filtered.

5 Concentration and flash chromatography (silica, 3% CH₃OH/CH₂Cl₂) yielded the major and minor product which was used as a mixture in the next step.

To a solution of the above mixture (assumed 0.245 mmol) in ethanol (4 mL, 0.05 M) at room temperature was added 2 mL

10 of 5 M NaOH and the cloudy reaction mixture was heated to reflux for 2 hours. The reaction solvent was concentrated, diluted with EtOAc, and washed 1 x H₂O and 1 x aqueous saturated NaCl. The organic layer was dried with MgSO₄ and filtered. Concentration and flash chromatography (silica, 4%

15 CH₃OH/CH₂Cl₂) yielded structure (7), 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-6-O-(5-acetamidopentyl)-2,3,4-tri-O-benzyl-β-D-glucopyranoside.

EXAMPLE 3

Preparation of Analog Having Structure (2), 2-(1H-indol-3-yl)ethyl-6-O-(5-aminopentyl)-2,4-di-O-deoxy-β-D-glucopyranoside;

20

A. Methyl 2-O-benzoyl-4,6-O-isopropylidene-α-D-glucopyranoside

To a stirred solution of methyl 2-4,6-O-isopropylidene-α-D-glucopyranoside (28.8 g, 123 mmol) in 410 mL of

25 dichloromethane at 0°C was added triethylamine (25.7 mL, 185 mmol) followed by benzoic anhydride (30.73 g, 135 mmol). The solution was warmed to room temperature and stirred for 24 hours. The solvent was removed under reduced pressure and the

30 residue was extracted with ethyl acetate (500 mL) and washed with H₂O (1 x 200 mL), a saturated salt solution (1 x 200 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 25% ethyl acetate in petroleum ether) provided the target compound (33.4 g, 80%) as a white form.

- 33 -

B. Methyl 2-0-benzoyl-3-0-(methylthio)thiocarbonyl-4,6-0-isopropylidene- α -D glucopyranoside

To a stirred solution of methyl 2-0-benzoyl-4,6-0-isopropylidene- α -D-glucopyranoside (1 g, 2.95 mmol) in 10 mL of dry THF at -78° was added sodium bis(trimethyl silyl)amide (1 M solution in THF, 3.54 mL, 3.4 mmol) followed immediately by carbon disulfide (248 μ l, 4.13 mmol). After stirring the solution for 15 minutes at -78°C, methyl iodide (550 μ l, 11.8 mmol) was added. The solution was stirred at -78°C an additional 10 minutes and then brought to room temperature. After stirring 30 minutes, the reaction was quenched with 2 mL of H₂O, diluted with 60 mL of ether, washed with H₂O (1 x 30 mL), a saturated solution of NaCl (1 x 30 mL) and dried over magnesium sulfate. Removal of the solvent yielded a crude xanthate (1.52 g crude). A 1.28 g aliquot of the crude xanthate was used in the next step without further purification. The remaining 0.24 g of target compound was purified by flash chromatography (silica, 20% ether in petroleum ether) to yield a white solid.

C. Methyl 2-0-benzoyl-3-deoxy-4,6-0-isopropylidene- α -D-glucopyranoside

To a solution of crude methyl 2-0-benzoyl-3-0-(methylthio)thiocarbonyl-4,6-0-isopropylidene- α -D glucopyranoside (1.28 g, 2.48 mmol theoretically) in 10 mL of dry toluene at room temperature was added 2,2'-azobisisobutyro-nitrile (AIBN, 40 mg) followed by tributyl tin hydride (2 mL, 7.48 mmol). The reaction was heated to reflux for 2 hours. The toluene was removed under reduced pressure. The resulting oil was dissolved in 60 mL of acetonitrile and washed with petroleum ether (3 x 20 mL). Concentration of the acetonitrile and flash chromatography (silica, 10% ethyl acetate in petroleum ether) yielded pure target compound (585 mg, 73% from 3b) as a clear oil.

- 34 -

D. Methyl 3-deoxy- α -D-glucopyranoside

To a stirred suspension of methyl 2-O-benzoyl-3-deoxy-4,6-O-isopropylidene- α -D-glucopyranoside (520 mg, 1.61 mmol) in 8 mL of methanol at room temperature was added sodium methoxide (70 mg, 1.29 mmol). After stirring 2h, the benzoyl group had been completely removed as evidenced by TLC. Amberlyst H⁺ resin was added and the mixture stirred for 1 hours until the generation of the free triol was completed as evidenced by TLC. After filtration, the solvents were removed under reduced pressure. Flash chromatography (silica, 10% methanol in methylene chloride) yielded pure target compound (286 mg, 100%) as an oil.

E. Methyl 2,4,6-tri-O-acetyl-3-deoxy- α -D-glucopyranoside

To a stirred solution of methyl 3-deoxy- α -D-glucopyranoside (535 mg, 3.0 mmol) in 10 mL of methylene chloride at 0° C was added triethylamine (2.92 mL, 21.0 mmol), acetic anhydride (1.41 mL, 15.0 mmol) and dimethyl amino pyridine, one at a time (37 mg, 0.30 mmol). The solution was warmed to room temperature. After stirring 7 hours, the solution was diluted with mL of ethyl acetate and washed with H₂O (1 x 30 mL), a saturated solution of NaCl (1 x 30 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 40% ether in petroleum ether) provided pure target compound (820 mg, 90%) as a clear oil.

F. 1,2,4,6-tetra-O-acetyl-3-deoxy- α -D-glucopyranoside

To a stirred solution of methyl 2,4,6-tri-O-acetyl-3-deoxy- α -D-glucopyranoside (127 mg, 0.41 mmol) in 3 mL of acetic anhydride at 0°C was added boron trifluoride etherate (15 μ l, 0.12 mmol). The solution was warmed to room temperature, stirred for 1.25 hours, and poured into 30 mL of an ice cold saturated solution of NaHCO₃ and extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with saturated aqueous NaHCO₃ and extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with saturated aqueous NaHCO₃ (2 x 40 mL), saturated aqueous

- 35 -

NaCl (1 x 40 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 30% ethyl acetate in petroleum ether) provided the target compound (133 mg, 96%) as an oil.

5 **G. Bromo 2,4,6-tri-O-acetyl-3-deoxy- α -D-glucopyranoside**

Hydrobromic acid (30% in acetic acid solution, 3 mL, 14.0 mmol) was added to 1,2,4,6-tetra-O-acetyl-3-deoxy- α -D-glucopyranoside (750 mg, 2.26 mmol) at 0°C. After 10 minutes, the resulting solution was warmed to room temperature for 30 minutes. The solution was then diluted with ether (20 mL) and poured into a mixture of ice and a saturated solution of NaHCO₃ (25 mL). An additional 30 mL of ether was added and the layers were separated. The organic layer was washed with saturated aqueous NaHCO₃ (3 x 25 mL), H₂O (1 x 25 mL, saturated aqueous NaCl (1 x 25 mL), and dried over magnesium sulfate. Removal of the solvent provided crude target compound, which was used in the next step without further purification.

20 **H. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-2,4,6-tri-O-acetyl-3-deoxy- β -D-glucopyranoside**

To a stirred suspension of flame dried 4A Sieves (1.33 g) in 11 mL of dry hexane at room temperature was added a solution of N-benzenesulfonyltryptophol (1.20 g, 4.0 mmol) in 4 mL of dry benzene. Next, a solution of the above bromo 2,4,6-tri-O-acetyl-3-deoxy- α -D-glucopyranoside (theoretically 2.26 mmol) in 4 mL of dry benzene was added, followed by silver(I)oxide (523 mg, 2.26 mmol). The reaction vessel was covered with aluminum foil and the suspension stirred for 3 days. After filtration through celite, concentration of the filtrate under reduced pressure and flash chromatography (silica, 10:1 methylene chloride:ether) provided pure target compound (781 mg, 60%) as a white foam.

35 **I. 2-(1-Phenylsulfonyl-indol-3-yl)ethyl-3-deoxy- β -D-glucopyranoside**

To a stirred suspension of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-2,4,6-tri-O-acetyl-3-deoxy- β -D-glucopyranoside (735 mg, 1.28 mmol) in 6.4 mL of methanol was added sodium

- 36 -

methoxide (55.2 mg, 1.02 mmol) at room temperature. After 90 minutes, the resulting solution was diluted with 6.4 mL of methanol (6.4 mL) and neutralized by addition of amberlyst H⁺ resin. The resin was quickly removed by filtration to avoid
5 formation of the methyl glucoside. Concentration of the filtrate and flash chromatography (silica, 12:1:1 methylene chloride, acetone, methanol) afforded pure target compound (498 mg, 87%) as a white solid.

J. 2-(1-Phenylsulfonyl-indol-3yl)ethyl-3-deoxy-6-0-
10 tert-butyldiphenylsilyl- β -D-glucopyranoside

To a stirred solution of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-3-deoxy- β -D-glucopyranoside (779 mg, 1.74 mmol) in 17 mL of dry DMF was added imidazole (260 mg, 3.83 mmol) followed by tert-butyldiphenylsilyl chloride (541 μ l, 2.09
15 mmol) at room temperature. The solution was stirred at 50°C for 24 hours. The reaction mixture was diluted with 250 mL of ethyl acetate and washed with H₂O (2 x 100 mL), saturated aqueous NaCl (1 x 100 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 3% methanol
20 in methylene chloride) provided pure target compound (1.04 g, 87%) as a white foam.

K. 2-(1-Phenylsulfonyl-indol-3yl)ethyl-3-deoxy-2,4-
di-0-benzyl-6-0-tert-butyldiphenylsilyl- β -D-
glucopyranoside

To a stirred suspension of sodium hydride (4.63 mmol, 185 mg, 60% oil dispersion) in 5 mL of dry THF at 0°C was added a solution of 2-(1-phenylsulfonyl-indol-3yl)ethyl-3-deoxy-6-0-tert-butyldiphenylsilyl- β -D-glucopyranoside (1.27 g, 1.85 mmol) in 10 mL of dry THF. After 10 minutes, the
30 mixture was warmed to room temperature. After stirring 1 hours, the suspension was cooled to 0°C and benzyl bromide (5.55 mmol, 660 μ l) was added followed by tetrabutylammonium iodide (68 mg, 0.185 mmol). The mixture was warmed to room temperature and stirred for 3 days. The reaction was then
35 quenched with 3 mL of aqueous saturated ammonium chloride at 0°C. The resulting solution was diluted with 80 mL of ether and washed with H₂O (2 x 30 mL), saturated aqueous NaCl (1 x

- 37 -

30 mL), and dried over magnesium sulfate. Concentration under reduced pressure and flash chromatography (silica, 25% ether in petroleum ether) provided pure target compound (760 mg, 47%) as a white foam.

5 L. 2-(1-phenylsulfonyl-indol-3yl)ethyl-3-deoxy-2,4-di-o-benzyl- β -D-glucopyranoside

To a stirred solution of 2-(1-phenylsulfonyl-indol-3yl)ethyl-3-deoxy-2,4-di-o-benzyl-6-O-tert-butyldiphenylsilyl- β -D-glucopyranoside (675 mg, 0.780 mmol) in 10 mL of dry THF
10 was added tetrabutylammonium fluoride (1 M solution in THF, 1.17 mmol, 1.17 mL) at room temperature. After stirring 2 hours, the solution was diluted with 80 mL of ethyl acetate and washed with H₂O (1 x 30 mL), saturated aqueous NaCl (1 x 30 mL), and dried over magnesium sulfate. Concentration and
15 flash chromatography (silica, 60% ether in petroleum ether) afforded pure target compound (445 mg, 91%) as an oil.

M. 2-(1-Phenylsulfonyl-indol-3yl)ethyl-3-deoxy-2,4-di-O-benzyl-6-O-trifluoromethylsulfonyl- β -D-glucopyranoside

20 To a stirred solution of 2-(1-phenylsulfonyl-indol-3yl)ethyl-3-deoxy-2,4-di-o-benzyl- β -D-glucopyranoside (360 mg, 0.575 mmol) in 3 mL of dichloromethane at -78°C was added, 2.6 di-tert-butyl-4-methylpyridine (189 mg, 0.92 mmol) followed by triflic anhydride (126 μ l, 0.748 mmol). After stirring 20
25 minutes at -78°C, the mixture was allowed to warm to room temperature for 20 minutes. The suspension was poured into aqueous saturated NaHCO₃ (15 mL) and extracted with ethyl acetate (1 x 35 mL). The organic layer was washed with saturated aqueous NaHCO₃ (3 x 15 mL), saturated aqueous NaCl (1
30 x 15 mL) and dried over magnesium sulfate. Concentration afforded crude target compound as an oil which was used in the next step without further purification.

N. 2-(1-Phenylsulfonyl-3-yl)ethyl-2,4-di-O-benzyl-3-deoxy-6-O-(N-trifluoroacetyl-5-aminopentyl)- β -D-glucopyranoside

35 To a stirred suspension of sodium hydride (8.63 mmol, 345 mg, 60% dispersion in oil) in 20 mL of dry THF at 0°C was

- 38 -

added a solution of N-trifluoro acetyl 5-amino pentanol (687 mg, 3.45 mmol) in 16 mL of dry THF. After stirring 10 minutes at 0°C, the suspension was allowed to warm to room temperature and stir for 90 minutes. The reaction mixture was then cooled to 0°C and a solution of crude triflate of step M (theoretically 0.575 mmol) in 22 mL of dry dichloromethane was added. The suspension was stirred for 30 minutes at 0°C and then warmed to room temperature. After stirring for an additional 24 hours, the reaction was quenched at 0°C with 10 mL of saturated aqueous ammonium chloride. The resulting mixture was diluted with ethyl acetate (200 mL) and washed with H₂O (1 x 75 mL), saturated aqueous NaCl (1 x 75 mL) and dried over magnesium sulfate. Concentration and flash chromatography (silica, eluted column 5 times with 1% methanol in methylene chloride to 2% methanol in methylene chloride) afforded the target compound (392 mg) as a white foam which was used without further purification in the next step.

O. Structure (2), 2-(1H-indol-3-yl)ethyl-6-O-(5-aminopentyl)-2,4-di-O-deoxy-β-D-glucopyranoside.

To a stirred solution of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-3-deoxy-2,4-di-O-benzyl-6-O-trifluoromethylsulfonyl-β-D-glucopyranoside (392 mg, theoretically 0.575 mmol) in 6 mL of ethanol at room temperature was added a solution of 5 M NaOH (1 mL, 5 mmol). The solution was allowed to reflux for 2 hours. The solvents were removed under reduced pressure, and the reaction mixture was diluted with dichloromethane (75 mL) and washed with aqueous HCl (25 mL, 5 mmol). The water layer was re-extracted with dichloromethane (2 x 75 mL). The combined organic layers were washed with saturated aqueous NaCl (2 x 25 ML) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 8% methanol in dichloromethane) afforded the pure product, structure (7) (172 mg, 52% for 3 steps) as an oil. R_f 0.22 (8% methanol in dichloromethane); ¹H NMR (500 MHz, CDCl₃, δ 8.44 (s, 1H), 7.57 (d, J = 7.7 Hz, 1H), 7.31-7.23 (m, 10H), 7.17-7.14 (m, 1H), 7.11-7.07 (m, 1H), 7.04 (d, J = 2.0 Hz, 1H), 4.71 (d, J = 11.8 Hz, 1H), 4.57 (d, J = 11.7 Hz, 1H), 4.56 (d, J = 11.9 Hz, 1H),

- 39 -

4.46(d, $J = 7.5$ Hz, 1H), d, $J = 11.5$ Hz, 1H, 4.20 (ddd, $J = 13.8, 9.4, 6.8$ Hz, 1H), 3.87 (ddd, $J = 14.9, 9.3, 7.4$ Hz, 1H), 3.55-3.50 (m, 3H), 3.32-3.26 (M, 2H), 3.11 (t, $J = 7.2$ Hz, 2H), 3.02 (dd, $J = 12.4, 2.9$ Hz, 1H), 2.68 (dd, $J = 12.4, 8.1$ Hz, 1H), 2.67-2.57 (m, 2H), 2.50 (ddd, $J = 12.3, 4.8, 4.8$ Hz, 1H), 2.20 (s, 3H), 1.57-2.44 (m, 5H), 1.36-1.30 (m, 2H); ^{13}C NMR (500 MHz, CDCl_3) δ 138.61, 137.92, 136.14, 128.41, 128.27, 127.79, 127.70, 127.53, 127.49, 122.18, 121.84, 119.18, 118.67, 112.56, 111.12, 105.22, 105.18, 76.53, 75.14, 74.28, 72.69, 70.99, 69.91, 62.45, 50.69, 49.49, 34.86, 32.28, 29.16, 25.80, 23.27; IR (thin film) 3325(m), 3065(w), 3035(w), 3015(w), 2940(s), 2870(s), 1500(w), 1458(m), 1354(w), 1220(w), 1076(s), 1030(m), 745(s), 700(s), cm^{-1} ; UV-Vis ($c = 6.5 \times 10^{-5}$, acetonitrile) λ_{max} 281.2 ($\epsilon = 6.2 \times 10^3$), 218.8 ($\epsilon = 3.62 \times 10^4$) nm; HRMS m/e calc'd for $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_5$ ($M + H$): 573.3315, found 573.3314; $[\alpha]_{\text{D}}^{20} + 16.7^\circ$ ($c = 0.15$, acetonitrile).

EXAMPLE 4

Preparation of Analog Having Structure (13), Methyl 2,3,4-tri-O-benzyl-6-O-(N-trifluoroacetyl-5-aminopentyl)- β -D-glucopyranoside

A. Methyl 6-O-tert-butyldiphenylsilyl- β -D-glucopyranoside

To a stirred solution of methyl β -D-glucopyranoside (5 g, 25.7 mmol) in 51 mL of dry DMF was added at room temperature imidazole (5.46 g, 80.2 mmol) followed by tert-butyldiphenyl-silyl chloride (11.3 mL, 43.4 mmol). The solution was heated to 50°C for 24 hours and the DMF was removed under reduced pressure. The reaction mixture was diluted with 200 mL of ethyl acetate and washed with H_2O (1 x 100 mL), saturated aqueous NaCl (1 x 100 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 4% methanol in dichloromethane) provided pure target compound (9.82 g, 88%) as a white foam.

- 40 -

B. Methyl 6-O-tert-butyldiphenylsilyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside

To a stirred suspension of sodium hydride (1.67 g, 41.6 mmol) in 100 mL of dry THF was added at 0°C a solution of methyl 6-O-tert-butyldiphenylsilyl- β -D-glucopyranoside (4.0 g, 9.25 mmol) in 50 mL of dry THF. After 5 minutes, the suspension was warmed to room temperature and stirred for 1 hour. Benzyl bromide (5.50 mL, 46.2 mmol) was added at room temperature followed by tetrabutylammonium iodide (341 mg, 0.93 mmol). The suspension was warmed to 50°C and stirred for 4 days. After quenching with 40 mL of saturated aqueous ammonium chloride, the resulting mixture was diluted with ether (600 mL) and washed with H₂O (2 x 200 mL), saturated aqueous NaCl (1 x 200 mL), and dried over magnesium sulfate. Concentration and flash chromatography (silica, 10% ether in petroleum ether) provided pure target compound (4.48 g, 69%) as a clear oil.

C. Methyl 2,3,4-tri-O-benzyl- β -D-glucopyranoside

To a stirred solution of methyl 6-O-tert-butyldiphenylsilyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside (2.81 g, 3.98 mmol) in dry THF (40 mL, 0.1 M) at room temperature was added tetrabutyl ammonium fluoride (4.37 mL, 4.37 mmol, 1 M solution in THF). After stirring for 3 hours, the reaction solution was diluted with ethyl acetate (300 mL) and washed with water (1 x 100 mL) and saturated aqueous NaCl (1 x 100 mL), and dried with magnesium sulfate. Concentration and flash chromatography (silica, 50% ether in petroleum ether) provided pure target compound (1.62 g, 88%) as a white solid.

D. Methyl 2,3,4-tri-O-benzyl-6-O-trifluoromethylsulfonyl- β -D-glucopyranoside

To a stirred solution of methyl 6-O-tert-butyldiphenylsilyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside (800 mg, 1.71 mmol) in 8.55 mL of dry dichloromethane at -78°C was added 2,6-di-tert-butyl-4-methyl pyridine (632 mg, 3.08 mmol) followed by triflic anhydride (345 μ L, 2.05 mmol). After stirring 15 minutes at -78°C, the mixture was warmed to room

- 41 -

temperature over 20 minutes, and then poured into a solution of saturated aqueous NaHCO_3 (20 mL) and extracted with ethyl acetate (50 mL). The organic layer was washed with saturated NaHCO_3 (3 x 20 mL), saturated aqueous NaCl (1 x 20 mL), and
5 dried over magnesium sulfate. Concentration provided crude target compound, which was used in the next step without further purification.

D. Structure (13), Methyl 2,3,4-tri-O-benzyl-6-O-(N-trifluoroacetyl-5-aminopentyl)- β -D-glucopyranoside
10

To a stirred suspension of sodium hydride (855 mg, 21.4 mmol, 60% oil dispersion) in 60 mL of dry THF at 0°C was added a solution of N-trifluoroacetyl-5-aminopentanol (1.7 g, 8.6 mmol) in 35 mL of dry THF. After stirring 10 minutes at 0°C ,
15 the suspension was warmed to room temperature, stirred for 1 hour, and cooled to 0°C . A solution of the above crude methyl 2,3,4-tri-O-benzyl-6-O-trifluoromethylsulfonyl- β -D-glucopyranoside (theoretically 1.71 mmol) in 57 mL of dry dichloromethane was added. The reaction mixture was stirred
20 at 0°C for 30 minutes and then warmed to room temperature. After stirring 24 hours, the reaction was cooled to 0°C and quenched with 40 mL of saturated aqueous ammonium chloride. The resulting solution was diluted with ethyl acetate (400 mL) and washed with H_2O (1 x 150 mL), saturated aqueous NaCl (1 x
25 150 mL) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 30% ethyl acetate in petroleum ether) provided the analog having structure (13), methyl 2,3,4-tri-O-benzyl-6-O-(N-trifluoroacetyl-5-aminopentyl)- β -D-glucopyranoside, (799 mg) as a white solid which was used
30 without further purification.

EXAMPLE 5

Preparation of Analog Having Structure (8), Methyl 6-O-(5-aminopentyl)-2,3,4-tri-O-benzyl- β -D-glucopyranoside,

To a stirred solution of methyl 2,3,4-tri-O-benzyl-6-O-
35 (N-trifluoroacetyl-5-aminopentyl)- β -D-glucopyranoside (799 mg, theoretically 1.71 mmol, structure (13) from Example 4) in 10

- 42 -

mL of ethanol at room temperature was added a solution of 5M (3 mL, 15 mmol). The solution was heated to reflux for 2 hours. The solvents were removed under reduced pressure. The reaction mixture was diluted with dichloromethane (70 mL) and washed with aqueous HCl (25 mL, 15 mmol). The water layer was re-extracted with dichloromethane (3 x 50 mL), and the combined organic layers were washed with saturated aqueous NaCl (1 x 75 mL) and dried over magnesium sulfate. Concentration and crystallization from ethyl acetate/petroleum ether provided pure analog having structure (8), methyl 6-O-(5-aminopentyl)-2,3,4-tri-O-benzyl- β -D-glucopyranoside, (675 mg, 72% from methyl 2,3,4-tri-O-benzyl-6-O-trifluoromethylsulfonyl- β -D-glucopyranoside) as a white solid: m.p. 95-95.5°C; R_f 0.19 (6% methanol in dichloromethane); ^1H NMR (500 MHz, CDCl_3) δ 7.35-7.24 (m, 15H), 4.92 (d, J = 7.5 Hz, 1H), 4.90 (d, J = 7.6 Hz, 1H), 4.85 (d, J = 11.0 Hz, 1H), 4.78 (d, J = 11.0 Hz, 1H), 4.70 (d, J = 11.0 Hz, 1H), 4.60 (d, J = 11.0 Hz, 1H), 4.32 (7.8, 1H), 3.66-3.59 (m, 3H), 3.56 (s, 3H), 3.48-3.36 (m, 3H), 2.94 (dd, J = 12.5, 2.1 Hz, 1H), 2.68 (dd, J = 12.0, 6.8 Hz, 1H), 2.64-2.53 (m, 2H), 1.71 (s, 2H), 1.59-1.53 (m, 2H), 1.51-1.45 (m, 2H), 1.42-1.36 (m, 2H), ^{13}C NMR (500 MHz, CDCl_3) δ 138.55, 138.47, 138.17, 128.39, 128.33, 128.03, 127.95, 127.85, 127.77, 127.60, 127.57, 104.72, 84.56, 82.45, 79.74, 75.66, 75.02, 74.74, 74.16, 62.62, 57.20, 50.69, 49.72, 32.49, 29.65, 23.37; IR (thin film) 3280(m), 3095(w), 3065(w), 3035(w), 2935(s), 2915(s), 2860(s), 1496(w), 1454(m), 1404(w), 1393(w), 1358(m), 1214(m), 1115(s), 1072(s), 1037(m), 1027(m), 1009(m), 911(w), 826(s), 747(s), 696(s) cm^{-1} ; HRMS m/e calc'd for $\text{C}_{33}\text{H}_{43}\text{O}_6\text{N}$ (M+H): 550.3168, found 550.3179; $[\alpha]_D^{20}$ +9.3° (c =0.15, acetonitrile).

EXAMPLE 6

Preparation of Analog Having Structure (12), 2-(1H-Indol-3yl)ethyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside

To a stirred solution of 2-(1-phenylsulfonyl-indol-3-yl)ethyl-2,3,4-tri-O-benzyl- β -D-glucopyranoside (100 mg, 0.136 mmol, prepared in Example 1, step G) in 3 ml of ethanol

- 43 -

at room temperature was added a solution of 5M NaOH (1 mL, 5 mmol). The reaction mixture was refluxed for 2h and the solvents were removed under reduced pressure. The resulting residue was diluted with dichloromethane (70 mL) and washed with aqueous HCl (24 mL, 5 mmol). The water layer was reextracted with dichloromethane (2 x 70 mL). The organic layers were combined and washed with saturated aqueous NaCl (1 x 50 mL) and dried over magnesium sulfate. Concentration and flash chromatography (silica, 25% ethyl acetate in petroleum ether) provided structure (12) (68 mg, 85%) as an oil: R_f 0.42 (40% ethyl acetate in petroleum ether); 1H NMR (500 MHz, $CDCl_3$) δ 7.83 (s, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.33-7.24 (m, 15H), 7.20-7.17 (m, 2H), 7.11 (t, J = 7.8 Hz, 1H), 7.01 (d, J = 1.8 Hz, 1H), 4.91 (d, J = 10.9 Hz, 1H), 4.85 (d, J = 10.9, 1H), 4.80 (d, J = 10.9 Hz, 1H), 4.79 (d, J = 11.0 Hz, 1H), 4.64 (d, J = 11.0 Hz, 1H), 4.63 (d, J = 11.0 Hz, 1H), 4.49 (d, J = 7.8 Hz, 1H), 4.22 (ddd, J = 9.4, 6.7, 6.7 Hz, 1H), 3.90-3.82 (m, 2H), 3.72-3.67 (m, 1H), 3.65 (ap. t, J = 9.1 Hz, 1H), 3.56 (ap. t, J = 9.3 Hz, 1H), 3.42 (ap. t, J = 8.1 Hz, 1H), 3.35 (ddd, J = 9.5, 4.3, 2.8 Hz, 1H), 3.11 (t, J = 7.0 Hz, 2H), 1.87 (dd, J = 7.6, 5.9 Hz, 1H); ^{13}C NMR (500 MHz, $CDCl_3$) δ 138.52, 138.44, 137.98, 136.17, 128.46, 128.36, 128.29, 128.05, 128.00, 127.89, 127.86, 127.60, 127.57, 127.45, 122.09, 122.01, 119.34, 118.68, 112.60, 111.13, 103.69, 84.49, 77.57, 75.64, 75.04, 75.01, 74.75, 70.25, 62.04, 25.86; IR (thin film) 3575(sh), 3435(m), 3085(sh), 3065(w), 3035(w), 2925(m), 2880(m), 1500(w), 1455(m), 1360(w), 1310(w), 1150(sh), 1085(s), 1030(s), 920(w), 810(w), 740(s), 700(s) cm^{-1} ; UV-Vis ($c=2.89 \times 10^{-4}$, acetonitrile) I_{max} 289.6 ($\epsilon=3.56 \times 10^3$), 281.2 ($\epsilon=4.24 \times 10^3$), 222.4 ($\epsilon=1.01 \times 10^4$) nm; HR MS m/e calc'd for $C_{37}H_{39}O_6N(M + NH_4)$: 611.3121, found 611.3043; $[\alpha]_D^{20}$ -2.5° ($c=1.37$, acetonitrile).

- 44 -

EXAMPLE 7

Preparation of Analog Having Structure (10), 2-(1H-Indol-3-yl)ethyl-6-O-aminopentyl)-2,3-di-O-benzyl-4-deoxy- β -D-glucopyranoside

5 **A. Methyl 2,3,6-tri-O-benzoyl-4-(methylthio)thiocarbonyl- α -D-glucopyranoside**

To a solution of the methyl 2,3,6-tri-O-benzoyl-4-O- α -D-glucopyranoside (5.00 g, 9.87 mmol) in 100 mL of dry THF at -78°C was added carbon disulfide (0.45 mL, 7.48 mmol) followed
10 by sodium bis(trimethylsilyl)amide (10.5 mL, 51.8 mmol). The solution was stirred at -78°C for 20 minutes. Methyl iodide (2.10 mL, 33.7 mmol) was added, the solution was stirred for 5 minutes at -78°C and then at room temperature for 45 minutes. The reaction was quenched by the addition of water
15 (5 mL) and the mixture was by extracted with ethyl acetate (2 X 100 mL). The organic layer was washed with a saturated solution of sodium chloride, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to yield a pale yellow oil (5.70g, 97%). The crude xanthate was
20 used without purification in the next step. An analytical sample was purified by flash column chromatography using 20% ethyl acetate in petroleum ether to yield the target compound as white crystals.

25 **B. Methyl 2,3,6-tri-O-benzoyl-4-deoxy- α -D-glucopyranoside**

To a solution of the crude methyl 2,3,6-tri-O-benzoyl-4-(methylthio)thiocarbonyl- α -D-glucopyranoside (5.70 g, 9.55 mmol) in 120 mL of dry toluene was added AIBN (50 mg). Tributyl tin hydride (6.68 mL, 24.8 mmol) was added and the
30 reaction was heated to reflux for 4 hours. The toluene was removed under reduced pressure. Acetonitrile (200 mL) was added and the mixture extracted with petroleum ether (5 x 100 mL) to remove all tin salts. After drying over anhydrous sodium sulfate, the solvent was removed under reduced pressure
35 to yield a clear colorless oil which solidified on standing. Purification by flash column chromatography using 20% ethyl

- 45 -

acetate in petroleum ether as the eluant gave the target compound as a white solid.

C. 1-O-Acetyl-2,3,6-tri-O-benzoyl-4-deoxy- α -D-glucopyranose

5 To a solution of methyl glycoside methyl 2,3,6-tri-O-benzoyl-4-deoxy- α -D-glucopyranoside (0.50 g, 1.1 mmol) in acetic anhydride (3.0 mL, 32 mmol) at 0°C was added boron trifluoride etherate (0.1 mL). The solution was stirred at room temperature for 4 hours, diluted with ethyl acetate and
10 poured in an ice-cold solution of saturated sodium bicarbonate. Extraction with ethyl acetate (2 X 100 mL) was followed by washing with a saturated solution of sodium chloride. The organic layer was dried over anhydrous sodium sulfate and evaporated under reduced pressure to yield the
15 product as a colorless oil which crystallized upon standing to give the target compound as white needles (0.45g, 85%).

D. 1-Bromo-2,3,6-tri-O-benzoyl-4-deoxy- α -D-glucopyranose

To a stirred solution of 1-O-acetyl-2,3,6-tri-O-benzoyl-4-deoxy- α -D-glucopyranose (0.137 g, 0.29 mmol) in 3.0
20 mL of dry dichloromethane at 0°C was added 30% hydrogen bromide in acetic acid (0.07 mL, 0.33 mmol). The solution was stirred under argon at room temperature for 4 hours, diluted with ethyl acetate (100 mL) and extracted with a saturated
25 solution of sodium bicarbonate. The organic layer was washed with a saturated solution of sodium chloride, dried over anhydrous sodium sulfate and evaporated under reduced pressure to yield the target compound as a colorless oil which solidified upon standing. Crystallization from ether and
30 petroleum ether gave the target compound as white crystals (0.15 g, 100%).

E. 2-(1-Phenylsulfonyl)-indol-3-yl-ethyl-2,3,6-tri-O-benzoyl-4-deoxy- β -D-glucopyranoside

To a mixture of activated powdered 4Å molecular sieves
35 (0.83 g), the protected tryptophol prepared in Example 1, step B (0.37 g, 1.23 mmol) and silver (I) oxide (0.83 g, 3.58 mmol) in a flask wrapped with aluminum foil was added a solution of

- 46 -

1-bromo-2,3,6,-tri-O-benzoyl-4-deoxy- α -D-glucopyranose (0.40 g, 0.814 mmol) in 16.7 mL of 40% hexane in benzene. The mixture was stirred under argon for two days, filtered through celite, washed with ethyl acetate and the solvent was removed to yield a colorless oil. Purification by flash column chromatography using 50% ether in petroleum ether gave the target compound as a colorless solid (0.50 g, 81%).

F. 2-(1-Phenylsulfonyl)-indol-3-yl-ethyl-4-deoxy- β -D-glucopyranoside

To a solution of 2-(1-phenylsulfonyl)-indol-3-yl-ethyl-2,3,6-tri-O-benzoyl-4-deoxy- β -D-glucopyranoside (120 mg, 0.158 mmol) in 20 mL of methanol was added sodium methoxide (0.027 g, 0.507 mmol). The solution was stirred under argon overnight. Amberlyst H+ resin was added and the reaction stirred until neutral to wet pH paper. The resin was removed by filtration and washed with methanol. The filtrate was concentrated under reduced pressure to yield a tan solid. Purification by flash column chromatography using 10% methanol in dichloromethane gave the target compound as a white solid (65 mg, 91%).

G. 2-(1-Phenylsulfonyl)-indol-3-yl-ethyl-6-O-tert-butyldiphenylsilyl-4-deoxy- β -D-glucopyranoside

To a solution of diol 2-(1-phenylsulfonyl)-indol-3-yl-ethyl-4-deoxy- β -D-glucopyranoside (0.24 g, 0.536 mmol) in 6 mL of dry DMF was added imidazole (73 mg, 1.07 mmol) followed by tert-butyldiphenylsilyl chloride (0.17 mL, 0.643 mmol). The solution was heated under argon in an oil bath at 70°C for 48 hours. The reaction was quenched by addition of methanol (5 mL). The solvents were removed under reduced pressure. The residue was extracted with ethyl acetate (2 x 200 mL), washed with a saturated solution of sodium chloride and dried over anhydrous sodium sulfate. Removal of the solvent under reduced pressure gave a pale yellow oil. Purification by flash column chromatography using 3% methanol in dichloromethane gave the target compound as a colorless oil (0.36 g, 97%).

- 47 -

H. 2-(1-Phenylsulfonyl)-indol-3-yl-ethyl-2,3,-di-O-
tert-butylldiphenylsilyl-4-deoxy- β -D-
glucopyranoside

To a stirred suspension of sodium hydride (73.0 mg.
5 3.04 mmol, 60% oil dispersion) in 2.7 mL of dry THF at 0°C was
added a solution of diol 2-(1-phenylsulfonyl)-indol-3-yl-
ethyl-6-O-tert-butylldiphenylsilyl-4-deoxy- β -D-glucopyranoside
(0.50 g, 0.729 mmol) in dry THF (6.8 mL). The reaction
mixture was stirred at room temperature for 30 minutes. The
10 mixture was cooled to 0°C and benzyl bromide (0.26 mL, 2.18
mmol) was added dropwise. After stirring at room temperature
for 3 days, the reaction was quenched by addition of ammonium
chloride (10 mL) followed by extraction with ether (2 X 100
mL). The organic layer was washed with a saturated solution
15 of sodium chloride, dried over anhydrous sodium sulfate and
evaporated under reduced pressure to yield a pale yellow oil.
Purification by flash column chromatography using 33% ether
in petroleum ether afforded the target compound as a colorless
oil (0.73 g, 76%).

20 I. 2-(1-Phenylsulfonyl)-indol-3-yl-ethyl-2,3,-di-O-
benzyl-4-deoxy- β -D-glucopyranoside

To a solution of the 2-(1-phenylsulfonyl)-indol-3-yl-
ethyl-2,3,-di-O-benzyl-6-O-tert-butylldiphenylsilyl-4-deoxy- β -
D-glucopyranoside (0.37 g, 0.427 mmol) in 10.5 mL of dry THF
25 was added tetrabutylammonium fluoride (1.33 mL, 1M in THF,
1.33 mmol). The solution was stirred under argon for 3 hours,
diluted with ethyl acetate (100 mL) and washed with water (100
mL). The organic layer was washed with a saturated solution
of sodium chloride and dried over anhydrous sodium sulfate.
30 Removal of the solvents under reduced pressure yielded a pale
yellow oil. Purification by flash column chromatography using
33% petroleum ether in ethyl acetate yielded the target
compound as a colorless oil (0.43 g, 85%).

- 48 -

J. 2-indol-3-yl-ethyl-2,3,-di-O-benzyl-4-deoxy- β -D-glucopyranoside

To a solution of the 2-(1-phenylsulfonyl)-indol-3-yl-ethyl-2,3,-di-O-benzyl-4-deoxy- β -D-glucopyranoside (140 mg, 0.223 mmol) in 6.0 mL of ethanol was added 5 M NaOH (2 mL) and the solution heated to reflux for hours. The solvents were removed under reduced pressure and the residue taken up in water (100 mL) and extracted with ethyl acetate (3 X 100 mL). The organic phase was washed with a saturated solution of sodium chloride, dried with anhydrous sodium sulfate and concentrated to yield a colorless oil. Purification by flash column chromatography using 3% methanol in dichloromethane yielded the target compound as a colorless oil (100 mg, 92%).

K. 5-Phthalimido-1-pentanol

To a solution of 5-amino-1-pentanol (5.00 g, 48.5 mmol) in benzene (150 mL) was added N-carboethoxyphthalimide (11.0 g, 50.2 mmol) and the solution was stirred at room temperature for 5 h). The solvents were removed under reduced pressure to yield a yellow oil. Purification by flash column chromatography using 25% ethyl acetate in petroleum ether yielded the target compound as a clear colorless oil (9.6 mg, 84%).

L. 5-Phthalimido-1-O-trifluoromethanesulfonylpentanol

To a solution of 5-phthalimido-1-pentanol (39.1 mg, 0.168 mmol) in dry dichloromethane (1.5 mL) was added 2,6-di-tert-butyl-4-methylpyridine (34.5 mg, 0.168 mmol) followed by triflic anhydride (28.3 μ g, 0.168 mmol). The solution was stirred at room temperature for 10 minutes. The reaction was poured into water (25 mL) and extracted with dichloromethane (2 x 50 mL). The organic layer was washed with a saturated sodium chloride solution and dried with anhydrous sodium sulfate. The solvents were removed under reduced pressure to yield a pale yellow solid which was used immediately without further purification.

- 49 -

M. 2-(1-Phenylsulfonyl-3-yl)ethyl-2,3-di-O-benzyl-4-deoxy-6-O(phthalimidopentyl)- β -D-glucopyranoside

To a solution of 5-phthalimido-1-O-trifluoromethanesulfonylpentanol (theoretically 0.168 mmol) in dry dichloromethane (1.5 mL) was 2,6-di-tert-butyl-4-methylpyridine (34.5 mg, 0.168 mmol). The solution was cooled to 0°C and to it was added a solution of 2-indol-3-yl-ethyl-2,3-di-O-benzyl-4-deoxy- β -D-glucopyranoside (18.4 mg, 0.029 mmol, from step J, above) in dry dichloromethane (1.5 mL). The solution was stirred for 30 minutes at 0°C and then sodium hydride (7.0 mg, 0.29 mmol, 60% dispersion in oil) was added. Stirring was continued at 0°C for 1 hour and then at room temperature for 24 hours. The reaction was poured into water (50 mL) and extracted with dichloromethane (2 x 100 mL). The organic layers were combined and washed with a saturated sodium chloride solution followed by drying with anhydrous sodium sulfate. The solvents were removed under reduced pressure to yield a pale yellow oil. Purification by flash column chromatography using 20% ethyl acetate in petroleum ether yielded the target compound as a clear colorless oil (19.4 mg, 80%).

N. Structure (10), 2-(1H-Indol-3-yl)ethyl-6-O-aminopentyl)-2,3-di-O-benzyl-4-deoxy- β -D-glucopyranoside

To a solution of 2-(1-phenylsulfonyl-3-yl)ethyl-2,3-di-O-benzyl-4-deoxy-6-O(phthalimidopentyl)- β -D-glucopyranoside (150 mg, 0.178 mmol) in methanol (8 mL) was added sodium methoxide (40 mg, 0.740 mmol). The solution was heated to reflux for 24 hours. The reaction was poured into water (100 mL) and extracted with dichloromethane (2 x 100 mL). The organic layers were combined and washed with a saturated solution of sodium chloride and dried with anhydrous sodium sulfate. Concentration of the solvents under reduced pressure yielded a pale yellow oil. Purification by flash column chromatography using 10% methanol in dichloromethane yielded structure (10) as a colorless oil (72.0 mg, 71%) R_f 0.32 (10% methanol in dichloromethane); ^1H NMR (500 MHz, CDCl_3) δ 7.74

- 50 -

(brm, 1H), 7.48 (d, $J = 7.8$ Hz, 1H), 7.36-6.93 (m, 15H), 4.62-4.49 (m, 4H), 4.32 (d, $J = 7.7$ Hz, 1H), 4.11 (dt, $J = 9.4, 6.7$ Hz, 1H), 3.78 (dt, 9.2, 7.4 Hz, 1H), 3.52 (m, 4H), 3.26 (m, 2H), 3.22 (t, $J = 7.2$ Hz, 1H), 3.13 (t, $J = 7.8$ Hz, 1H), 3.00 (t, $J = 7.0$ Hz, 2H), 2.00 (ddd, $J = 6.7, 5.2, 1.4$ Hz, 1H), 1.29 (m, 9H); ^{13}C NMR (500 MHz, CDCl_3) δ 140.11, 138.10, 130.75, 130.59, 129.31, 128.92, 128.84, 128.57, 128.44, 122.24, 119.40, 112.82, 112.31, 105.01, 84.13, 79.55, 75.76, 74.12, 73.12, 72.53, 72.18, 71.29, 41.05, 34.54, 30.38, 29.90, 27.07, 24.72, IR (CHCl_3) 3350, 3060, 2930, 2860, 1630, 1520, 1450, 1400, 1270, 1100, 740, 700; UV ($c=1.57 \times 10^{-4}\text{M}$, acetonitrile) λ_{max} 280.0 ($\epsilon=1.41 \times 10^3$), 224.8 ($\epsilon=1.66 \times 10^3$) nm; HRMS m/e calc'd for $\text{C}_{35}\text{H}_{45}\text{N}_2\text{O}_5$ ($M+H$): 573.3328, found 573.3301; $[\alpha]_D^{20} + 3.89^\circ$ ($c=1.8$, acetonitrile).

15 EXAMPLE 8

Preparation of Analog Having Structure (11), 2-Indol-3-yl-ethyl-2,3,-di-O-benzyl-4-deoxy- β -D-glucopyranoside

To a solution of 2-(1-phenylsulfonyl)-indol-3-ylethyl-2,3,-di-O-benzyl-4-deoxy- β -D-glucopyranoside (140 mg, 0.223 mmol, from Example 7, step I, above) in 6.0 mL of ethanol was added 5M NaOH (2 mL) and the solution heated to reflux for 2 hours. The solvents were removed under reduced pressure and the residue taken up in water (100 mL) and extracted with ethyl acetate (3 X 100 mL). The organic phase was washed with a saturated solution of sodium chloride, dried with anhydrous sodium sulfate and concentrated to yield a colorless oil. Purification by flash column chromatography using 3% methanol in dichloromethane yielded the analog having structure (11) (2-indol-3-yl-ethyl-2,3,-di-O-benzyl-4-deoxy- β -D-glucopyranoside) as a colorless oil (100 mg, 92%). R_f 0.59 (10% methanol in dichloromethane); ^1H NMR (500 MHz, CDCl_3) δ 7.86 (br s, 1H), 7.59 (d, $J = 7.8$ Hz, 1H), 7.31-6.99 (m, 14H), 4.78-4.66 (m, 4H), 4.41 (d, $J = 7.7$ Hz, 1H), 4.22 (dt, $J = 9.4, 7.4$ Hz, 1H), 3.61-3.56 (m, 3H), 3.49-3.45 (m, 1H), 3.32 (t, $J = 7.9$ Hz, 1H), 3.11 (t, $J = 6.9$ Hz, 2H), 2.03 (br s, 1H), 1.95 (ddd, $J = 12.8, 5.3, 1.8$ Hz, 1H), 1.49 (q, $J = 11.7$

- 51 -

Hz, 1H); ^{13}C NMR (500 MHz, CDCl_3) δ 138.72, 138.48, 136.12, 128.33, 128.20, 127.97, 127.60, 127.56, 127.46, 122.15, 121.92, 119.27, 118.66, 112.57, 111.10, 103.87, 82.81, 78.10, 74.86, 72.23, 72.13, 70.18, 65.17, 32.69, 25.84; UV-Vis ($c = 1.85 \times 10^{-4}$, acetonitrile) λ_{max} 281.2 ($\epsilon = 614.13$), 220.0 ($\epsilon = 864.86$) nm; HRMS m/e calculated for $\text{C}_{30}\text{H}_{34}\text{NO}_5$ ($M+H$): 488.2436, found 488.2483; $[\alpha]_{\text{D}^{20}} +5.55^\circ$ ($c = 1.8$, acetonitrile).

EXAMPLE 9

Preparation of Imidazol Compounds. To distinguish the compounds described in this example an "I" precedes each compound number. The chemical structures and synthetic schemes for the compounds in this example are presented in Figure 1.

A. Phthalimido-protected amine (-)-I-21.

5-Phthalimidopentyl triflate I-20 was prepared as follows: A stirred solution of 5-phthalimido-1-pentanol (1.32 g, 4.67 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.960 g, 4.67 mmol) in dry dichloromethane (10 ml) was treated with triflic anhydride (0.784 ml, 4.67 mmol). After 10 min at room temperature, the mixture was diluted with water (100 ml) and extracted with dichloromethane (2 x 200 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction. Sodium hydride (60% dispersion in oil, 0.20 g, 5.06 mmol) was added to a solution of alcohol I-19 (1.27 g, 3.89 mmol), 5-phthalimidopentyl triflate I-20 (4.67 mmol), and 15-crown-5 (20 mg, 2.3 mol %), in methylene chloride (100 ml) at 0°C . After stirring for 24 h at room temperature, the mixture was poured into water. The aqueous layer was extracted with methylene chloride (3 x 50 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (3% ether/methylene chloride) provided I-21 (1.82 g, 86% yield) as a colorless oil: $[\alpha]_{\text{D}^{25}} -8.2^\circ$ (c 0.70, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.80 (m, 2 H), 7.68 (m, 2 H), 7.25-7.34 (m, 10

- 52 -

H), 6.38, (dd, $J = 6.1, 1.2$ Hz, 1 H), 4.84 (m, 2 H), 4.66 (d, $J = 11.4$ Hz, 1 H), 4.63 (d, $J = 11.7$ Hz, 1 H), 4.55 (d, $J = 11.7$ Hz, 1 H), 4.19 (m, 1 H), 4.00 (m, 1 H), 3.81 (dd, $J = 8.7, 6.2$ Hz, 1 H), 3.64-3.74 (m, 4 H), 3.40-3.50 (m, 2 H),
5 1.60-1.70 (m, 4 H), 1.40 (m, 2 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 168.4, 144.8, 138.4, 138.3, 133.9, 132.2, 128.4, 127.9, 127.8, 127.6, 123.2, 99.9, 76.8, 75.8, 74.5, 73.8, 71.4, 70.5, 69.2, 37.9, 29.2, 28.5, 23.5; high resolution mass spectrum (Cl, NH_3) m/z 541.2483 (M^+ ; calcd for $\text{C}_{33}\text{H}_{35}\text{NO}_6$: 541.2464).

10 B. Alcohol (-)-I-23.

A solution of dimethyldioxirane in acetone (1.2 equiv, ca. 0.05 M) was added dropwise to glycal I-21 (1.53 g, 2.80 mmol) in dichloromethane (26 ml) at 0°C . The mixture was stirred at 0°C for 1 h and concentrated in vacuo.

15 To a solution of the crude epoxide and I-22 (1.15 g, 3.82 mmol) in THF (12 ml) at -78°C was added ZnCl_2 (1.0 M in ether, 5.6 ml, 5.6 mmol) and the mixture was stirred at -78°C for 1 h. The solution was then slowly warmed to room temperature and stirred for 18 h. The mixture was poured into
20 saturated aqueous sodium bicarbonate (50 ml) and extracted with ethyl acetate (3 x 50 ml) and the combined extracts were washed with water, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography (45% ethyl acetate/hexane) yielded I-23 (1.05 g, 44% yield) as a
25 colorless oil: $[\alpha]_{\text{D}}^{25} -8.1^\circ$ (c 1.8 CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.96 (dd, $J = 8.1, 0.6$ Hz, 1 H), 7.85 (dd, $J = 8.2, 0.9$ Hz, 2 H), 7.78 (m, 2 H), 7.66 (m, 2 H), 7.20-7.50 (m, 17 H), 4.89 (d, $J = 11.3$ Hz, 1 H), 4.86 (d, $J = 11.0$ Hz, 1 H), 4.83 (d, $J = 11.4$ Hz, 1 H), 4.60 (d, $J = 10.9$ Hz, 1 H), 4.24
30 (d, $J = 7.6$ Hz, 1 H), 4.20 (dt, $J = 9.5, 6.4$ Hz, 1 H), 3.76 (dt, $J = 9.5, 7.2$ Hz, 1 H), 3.37-3.68 (m, 10 H), 2.98 (m, 2 H), 2.13 (br s, 1 H), 1.57-1.68 (m, 4 H), 1.38 (m, 2 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 168.4, 138.6, 138.2, 135.1, 133.8, 133.7, 132.1, 131.0, 129.1, 128.4, 127.9, 127.8, 127.7, 126.7,
35 124.7, 123.5, 123.1, 119.7, 119.4, 113.7, 102.8, 84.4, 76.5, 75.1, 71.5, 69.6, 68.7, 37.8, 29.2, 28.4, 25.4, 23.5; high

- 53 -

resolution mass spectrum (Cl, NH₃) m/z 662.2774 (M^+ ; calcd for C₃₅H₄₂SO₇: 662.2775).

C. Dibenzyl ether (-)-I-24.

A solution of I-23 (0.455 g, 0.530 mmol) in THF (10 ml) was cooled to -78°C and treated with carbon disulfide (27 ml, 0.583 mmol) followed by sodium bis(trimethylsilyl)amide (0.6 M in toluene, 0.953 ml, 0.572 mmol). After 20 min, methyl iodide (59 ml, 0.640 mmol) was added and the solution was stirred for 5 min at -78°C and then at room temperature for 45 min. The reaction mixture was quenched with water (50 ml) and extracted with ethyl acetate (3 x 50 ml). The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording the crude xanthate as a pale yellow oil (0.462 g, 92% yield) which was used without purification in the next step.

To a solution of the crude xanthate (0.462 g, 0.487 mmol) and AIBN (10 mg) in toluene (8 ml) was added tributyltin hydride (0.214 ml, 0.795 mmol) and the reaction mixture heated at reflux for 4 h, cooled, and concentrated in vacuo. The residue was taken up in acetonitrile (30 ml) and washed with petroleum ether (5 x 10 ml), dried over sodium sulfate, filtered, and concentrated in vacuo to an oil. Flash chromatography (20% ethyl acetate/petroleum ether) yielded I-24 (0.296 g, 72% yield) as a colorless oil; $[\alpha]_D^{25}$ -10° (c 1.1 CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.96 (d, J = 7.5 Hz, 1 H), 7.84 (m, 2 H), 7.79 (m, 2 H), 7.66 (m, 2 H), 7.20-7.41 (m, 15 H), 4.91 (d, J = 11.0 Hz, 1 H), 4.60 (m, 2 H), 4.66 (d, J = 11.7 Hz, 1 H), 4.41 (dd, J = 9.7, 1.8 Hz, 1 H), 4.15 (dt, J = 9.5, 6.6 Hz, 1 H), 3.59-3.71 (m, 6 H), 3.47 (m, 2 H), 3.40 (m, 1 H), 2.94 (t, J = 6.6 Hz, 2 H), 2.57 (ddd, J = 14.2, 5.0, 3.2 Hz, 1 H), 1.57-1.68 (m, 5 H), 1.38 (m, 2 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.5, 25.5, 28.4, 29.2, 36.7, 37.9, 68.1, 70.0, 71.4, 75.0, 75.2, 78.2, 79.3, 99.9, 113.6, 119.6, 123.1, 123.5, 124.7, 126.7, 127.7, 128.0, 128.4, 129.2, 131.1, 132.1, 133.6, 133.8, 135.1, 138.3, 138.5, 168.4; high resolution mass spectrum (Cl, NH₃) m/z 814.3287 (M^+ ; calcd for C₄₄H₅₀SO₈N₂: 814.3289).

- 54 -

D. Amine (-)-I-15.

A solution of hydrazine (0.2 M in MeOH, 6 ml) was added to I-24 (0.034 g, 0.043 mmol). After stirring for 16 h, the reaction mixture was concentrated in vacuo, the residue was dissolved in ethanol (4 ml), and 5N NaOH (0.90 ml) added. The mixture was heated at reflux for 4 h, cooled, and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (11% methanol/methylene chloride) afforded I-15 (11 mg, 44%) as a pale yellow oil; $[\alpha]_D^{25} -15^\circ$ (c 0.62, CHCl₃); IR (CHCl₃) 3490 (m), 3345 (br, m), 3020 (m), 2945 (s), 2882 (s), 1625 (w), 1500 (w), 1459 (m), 1370 (m), 1230 (w), 1100 (s), 695 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.80 (br s, 1 H), 7.49 (d, J = 7.9 Hz, 1 H), 7.19-7.31 (m, 11 H), 7.10 (t, J = 7.1 Hz, 1 H), 7.00 (t, J = 8.0 Hz, 1 H), 6.97 (s, 1 H), 4.83 (d, J = 11.1 Hz, 1 H), 4.59 (d, J = 11.7 Hz, 1 H), 4.51 (d, J = 11.0 Hz, 1 H), 4.50 (d, J = 11.7, 1 H), 4.39 (d, J = 9.7 Hz, 1 H), 4.00 (apparent q, J = 7.3 Hz, 1 H), 3.67 (apparent q, J = 7.3 Hz, 1 H), 3.60 (d, J = 9.0 Hz, 1 H), 3.56 (m, 1 H), 3.46 (dd, J = 10.8, 5.3 Hz), 3.31 (m, 4 H), 2.98 (t, J = 7.2 Hz, 2 H), 2.50 (t, J = 7.3 Hz, 2 H), 2.28 (m, 2 H), 1.57 (q, J = 10 Hz, 1 H), 1.42 (m, 4 H), 1.19 (m, 2 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 138.3, 138.2, 136.2, 128.4, 128.0, 127.7, 127.5, 122.3, 121.8, 119.1, 118.7, 112.0, 111.4, 99.9, 79.3, 78.2, 74.9, 71.4, 71.0, 69.9, 69.8, 39.7, 36.7, 28.8, 27.6, 25.7, 23.1; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.3371 [(M + H)⁺; calcd for C₃₅H₄₄N₂O₅: 573.3328].

E. Azide (-)-I-27.

5-Azidodopentyl triflate I-26 was prepared as follows: A stirred solution of 5-azido-1-pentanol (0.14 g, 1.08 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.22 g, 1.08 mmol) in dry dichloromethane (5 ml) was treated with triflic anhydride (0.19 ml, 1.08 mmol). After 10 min at room temperature, the mixture was diluted with water (100 ml) and extracted with dichloromethane (2 x 200 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and

- 55 -

concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction.

Sodium hydride (60% dispersion in oil, 0.053 g, 2.30 mmol) was added to a solution of alcohol I-25 (0.353 g, 1.08 mmol), 5-azidopentyl triflate 26 (1.08 mmol), and 15-crown-5 (10 mg), in methylene chloride (10 ml) at 0°C. After stirring for 24 h at room temperature, the mixture was poured into water. The aqueous layer was extracted with methylene chloride (3 x 50 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (20% ethyl acetate/petroleum ether) provided I-27 (260 mg, 60%) as a colorless oil; $[\alpha]_D^{25}$ -8.5° (c 0.89, CHCl₃); IR (CHCl₃) 3090 (w), 3062 (w), 3030 (w), 3005 (w), 2940 (m), 2865 (m), 2100 (s), 1650 (m), 1495 (w), 1455 (m), 1355 (w), 1235 (m), 1210 (m), 1100 (br, s), 1070 (s), 1028 (s), 705 (w), 691 (m), cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.42 (m, 2 H), 1.61 (m, 4 H), 3.24 (t, J = 6.9 Hz, 2 H), 3.47 (m, 4 H), 3.70 (dd, J = 10.8, 2.7 Hz, 1 H), 3.76 (dd, J = 10.8, 5.1 Hz, 1 H), 3.84 (dd, J = 8.7, 6.2 Hz, 1 H), 4.03 (m, 1 H), 4.21 (ddd, J = 6.2, 2.5, 1.5 Hz, 1 H), 4.56 (d, J = 11.6 Hz, 1 H), 4.64 (d, J = 11.6 Hz, 1 H), 4.64 (d, J = 11.6 Hz, 1 H), 4.69 (d, J = 11.4 Hz, 1 H), 4.88 (m, 2 H), 6.42 (m, 2 H), 7.27-7.38 (m, 10 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.4, 28.7, 29.2, 51.3, 69.2, 70.5, 71.3, 73.8, 74.5, 76.8, 99.9, 127.6, 127.8, 128.4, 138.3, 144.7; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 472.2031 [(M + Cl)⁺; calcd for C₂₅H₃₁O₄N₃Cl :8471.2003].

F. Amide (-)-I-28.

To a solution of sugar I-27 (0.117 g, 0.268 mmol) in THF (5 ml) was added H₂O (0.217 ml, 12.1 mmol) and PPh₃ (0.176 g, 0.671 mmol) and the reaction mixture was heated to 55°C for 10 h, cooled, and concentrated in vacuo. Flash chromatography (15% methanol/methylene chloride) provided the amine as a colorless oil (82 mg, 77%); $[\alpha]_D^{25}$ -7.2° (c 0.25, CHCl₃); IR (CHCl₃) 3500-2600 (br, w), 3090 (w), 3060 (w), 3003 (m), 2933 (s), 2864 (s), 1650 (m), 1495 (w), 1452 (m), 1355 (w), 1235 (m), 1220 (m), 1100 (br, s), 1025 (m), 850 (br, w), 690 (m)

- 56 -

cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.36 (m, 2 H), 1.43 (quin., J = 7.4 Hz, 1 H), 1.59 (quin., J = 6.7 Hz, 1 H), 2.65 (t, J = 6.9 Hz, 2 H), 3.46 (m, 2 H), 3.69 (dd, J = 10.8, 2.7 Hz, 1 H), 3.73 (dd, J = 10.8, 5.1 Hz, 1 H), 3.82 (dd, J = 8.7, 6.3 Hz, 1 H), 4.01 (m, 1 H), 4.22 (m, 1 H), 4.55 (d, J = 11.7 Hz, 1 H), 4.63 (d, J = 11.7 Hz, 1 H), 4.67 (d, J = 11.4 Hz, 1 H), 4.85 (m, 2 H), 6.40 (d, J = 6.2 Hz, 1 H), 7.26-7.36 (m, 10 H); ¹³C NMR (125.8 MHz, CDCl₃) δ 22.4, 28.4, 32.5, 41.0, 68.1, 69.4, 70.5, 72.7, 73.5, 74.8, 75.7, 98.9, 126.6, 126.7, 127.3, 127.3, 137.2, 137.3, 143.7.

To a solution of the amine (0.077 g, 0.19 mmol) in CH₂Cl₂ (2.5 ml) at 0°C was added Et₃N (0.040 ml, 0.29 mmol) and Ac₂O (0.020 ml, 0.21 mmol). After stirring for one minute, the mixture was poured into water. The aqueous layer was extracted with methylene chloride (3 x 20 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (8% methanol/methylene chloride) provided I-28 (80 mg, 94%) as a colorless oil; [α]_D²⁵ -8.2° (c 0.38, CHCl₃); IR (CHCl₃) 3450 (w), 3090 (w), 3062 (w), 3004 (m), 2940 (m), 2865 (m), 1665 (s), 1520 (br, m), 1455 (m), 1367 (br, m), 1237 (m), 1208 (m), 1102 (br, s), 1025 (m), 690 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.37 (m, 2 H), 1.48 (quin., J = 7.5 Hz, 2 H), 1.59 (m, 2 H), 1.92 (s, 3 H), 3.20 (m, 2 H), 3.45 (m, 2 H), 3.68 (dd, J = 10.9, 2.6 Hz, 1 H), 3.73 (dd, J = 10.9, 5.1 Hz, 1 H), 3.81 (dd, J = 8.7, 6.3 Hz, 1 H), 4.00 (m, 1 H), 4.20 (m, 1 H), 4.55 (d, J = 11.6 Hz, 1 H), 4.63 (d, J = 11.6 Hz, 1 H), 4.67 (d, J = 11.3 Hz, 1 H), 4.87 (m, 2 H), 5.45 (br s, 1 H), 6.39 (dd, J = 6.2, 1.3 Hz, 1 H), 7.27-7.35 (m, 10 H); ¹³C NMR (125.8 MHz, CDCl₃) δ 23.3, 23.6, 29.3, 29.3, 39.5, 69.2, 70.6, 71.4, 73.8, 74.6, 75.9, 77.6, 100.0, 127.7, 127.8, 127.8, 128.5, 138.3, 144.7, 170.0; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 488.2537 [(M + Cl)⁺; calcd for C₂₇H₃₅O₅NCl : 488.2515].

35 G. α-Amide (+)-I-29.

To a solution of amide I-28 (0.022 g, 0.051 mmol) and tryptophol (0.041 g, 0.26 mmol) in acetonitrile (1 ml) was

- 57 -

added CSA (1 mg). After stirring for 24 h at room temperature the mixture was added to saturated sodium bicarbonate and extracted with methylene chloride (3 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (ethyl acetate) provided I-29 α (4.2 mg, 14%) as a colorless oil; $[\alpha]_D^{25} +55.0^\circ$ (c 0.40, CHCl₃); IR (CHCl₃) 3485 (m), 3460 (m), 3300 (br, w), 3015 (m), 2950 (m), 2875 (m), 1670 (s), 1525 (w), 1460 (m), 1370 (w), 1130 (m), 1105 (br, s), 1030 (m), 980 (w), 695 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.31 (m, 2H), 1.43 (quin., J = 7.4 Hz, 2 H), 1.54 (m, 2 H), 1.67 (dt, J = 12.4, 3.3 Hz, 1 H), 1.91 (s, 3 H), 2.29 (dd, J = 12.7, 5.0 Hz, 1 H), 3.02 (m, 2 H), 3.14 (m, 2 H), 3.35 (m, 1 H), 3.40-3.59 (m, 4 H), 3.65 (m, 2 H), 3.89 (q, J = 7.5 Hz, 1 H), 4.21 (m, 1 H), 4.58 (d, J = 11.1 Hz, 1 H), 4.63 (d, J = 11.5 Hz, 1 H), 4.67 (d, J = 11.5 Hz, 1 H), 4.90 (d, J = 11.1 Hz, 1 H), 4.97 (d, J = 11.7 Hz, 1 H), 5.37 (br s, 1 H), 6.98 (s, 1 H), 7.09 (t, J = 7.1 Hz, 1 H), 7.17 (t, J = 7.1 Hz, 1 H), 7.27-7.37 (m, 11 H), 7.59 (d, J = 7.9 Hz, 1 H), 8.30 (br s, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.2, 23.6, 25.6, 29.3, 35.5, 39.6, 67.4, 69.8, 70.5, 71.3, 71.7, 74.8, 97.2, 111.1, 112.8, 118.8, 119.1, 121.8, 121.9, 127.5, 127.6, 127.8, 128.4, 136.2, 138.7, 170.1; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 615.3407 [(M + H)⁺; calcd for C₃₇H₄₇O₆N₂ : 615.3434].

25 H. β -Amide (-)-I-29.

(1.7 mg, 6%) as a colorless oil; $[\alpha]_D^{25} -13.0^\circ$ (c 0.16, CHCl₃); IR (CHCl₃) 3480 (w), 3010 (m), 2940 (m), 2877 (m), 1670 (s), 1532 (w), 1458 (m), 1369 (m), 1270 (w), 1100 (br s), 1011 (w), 695 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.22 (m, 2 H), 1.45 (m, 2 H), 1.53-1.69 (m, 3 H), 1.92 (s, 3 H), 2.34 (m, 1 H), 3.08 (t, J = 7.5 Hz, 2 H), 3.14 (m, 2 H), 3.38 (m, 1 H), 3.40-3.51 (m, 3 H), 3.57-3.67 (m, 2 H), 3.69 (dd, J = 10.8, 1.8 Hz, 1 H), 3.75 (m, 1 H), 4.13 (dt, J = 9.6, 2.0 Hz, 1 H), 4.46 (dd, J = 9.7, 1.8 Hz, 1 H), 4.58 (d, J = 11.7 Hz, 1 H), 4.61 (d, J = 11.1 Hz, 1 H), 4.66 (d, J = 11.6 Hz, 1 H), 4.92 (d, J = 11.0 Hz, 1 H), 5.39 (br s, 1 H), 7.05 (s, 1 H), 7.09 (m, 1 H), 7.17 (m, 1 H), 7.26-7.36 (m, 11 H), 7.59 (d, J = 7.8

- 58 -

Hz, 1 H), 8.37 (br s, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 23.6, 23.7, 25.7, 29.3, 36.8, 39.7, 69.6, 70.1, 71.4, 74.9, 75.2, 76.8, 78.4, 79.4, 99.9, 111.2, 112.5, 118.7, 119.1, 121.9, 122.1, 127.7, 127.7, 127.9, 128.4, 138.3, 170.2; high
5 resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 615.3410 $[(M + H)^+]$; calcd for $\text{C}_{37}\text{H}_{47}\text{O}_6\text{N}_2$: 615.3434].

I. Acetal (-)-I-31.

To a solution of the triol I-30 (9.43 g, 21.1 mmol) dissolved in DMF (35 ml) was added α,α -dimethoxytoluene (3.42
10 ml, 22.8 mmol) and $p\text{TsOH}$ (100 mg) and the mixture was heated to 45°C under aspirator pressure for 5 h. After cooling, the mixture was added to H_2O (300 ml) and saturated sodium bicarbonate (10 ml) and extracted with ethyl acetate (3 x 150 ml). The combined extracts were washed with water, dried over
15 magnesium sulfate and concentrated in vacuo. Flash chromatography (40% ethyl acetate/petroleum ether) provided I-31 (10.0 g, 89% yield) as a colorless oil: $[\alpha]_D^{25} -8.2^\circ$ (c 0.70, CHCl_3); 3590 (br w), 3080 (w), 3010 (w), 2920 (w), 2880 (w), 1450 (m), 1375 (m), 1330 (w), 1280 (w), 1182 (m), 1175
20 (s), 1130 (m), 1120 (m), 1100 (s), 1085 (s), 1070 (s), 1018 (m), 1000 (m), 905 (w), 680 (w), 595 (m), 565 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.70 (q, $J = 11.7$ Hz, 1 H), 2.05 (br s, 1 H), 2.43 (dt, $J = 9.2, 4.6$ Hz, 1 H), 3.00 (m, 2 H), 3.44 (m, 1 H), 3.59 (m, 2 H), 3.79 (m, 2 H), 4.24 (dt, $J = 9.5, 6.4$ Hz,
25 1 H), 4.31 (m, 2 H), 5.52 (s, 1 H), 7.25 (m, 2 H), 7.36 (m, 4 H), 7.40-7.54 (m, 6 H), 7.87 (m, 2 H), 7.99 (d, $J = 8.1$ Hz, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 25.5, 34.9, 68.7, 68.8, 68.9, 70.5, 76.0, 101.7, 105.2, 113.6, 119.2, 119.5, 123.0, 123.3, 124.6, 126.5, 128.3, 128.8, 129.0, 131.0, 133.6, 135.0,
30 137.1; high resolution mass spectrum (Cl , CH_4) m/z 536.1722 $[(M + H)^+]$; calcd for $\text{C}_{29}\text{H}_{30}\text{SO}$, N: 536.1743]. J.Acetal (-)-I-32.

To a solution of the acetal I-31 (1.84 g, 3.44 mmol) dissolved in DMF (4 ml) was added imidazole (0.52 g, 7.57 mmol) followed by TIPSCl (0.81 ml, 3.78 mmol). After stirring
35 for 24 h, the mixture was added to H_2O (200 ml) and extracted with ether (3 x 100 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in

- 59 -

vacuo. Flash chromatography (15% ethyl acetate/petroleum ether) provided I-32 (2.12 g, 90% yield) as a colorless oil: $[\alpha]_D^{25} -27.8^\circ$ (c 0.95, CHCl_3); IR (CHCl_3) 3080 (w), 3040 (w), 3020 (w), 2960 (s), 2905 (s), 2880 (s), 1467 (m), 1453 (m), 1335 (w), 1285 (w), 1190 (m), 1179 (s), 1135 (s), 1130 (s), 1095 (s), 1000 (br m), 885 (m), 810 (m), 720 (w), 670 (br, m), 600 (m), 572 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.05 (m, 21 H), 1.77 (q, $J = 11.2$ Hz, 1 H), 2.44 (m, 1 H), 3.00 (t, $J = 7.7$ Hz, 2 H), 3.42 (m, 1 H), 3.56 (m, 1 H), 3.73-3.85 (m, 3 H), 4.12 (m, 1 H), 4.29 (dd, $J = 10.5, 4.9$ Hz, 1 H), 4.38 (d, $J = 7.3$ Hz, 1 H), 5.50 (s, 1 H), 7.23 (m, 1 H), 7.25 (s, 1 H), 7.28-7.38 (m, 3 H), 7.43 (m, 3 H), 7.46-7.54 (m, 4 H), 7.86 (m, 2 H), 7.97 (dt, $J = 8.3, 0.8$ Hz); ^{13}C NMR (62.9 MHz, CDCl_3) δ 12.4, 17.7, 18.0, 25.6, 38.0, 68.9, 69.2, 70.0, 70.2, 75.9, 101.7, 106.0, 113.7, 119.4, 123.1, 123.5, 124.8, 126.2, 126.7, 128.4, 129.1, 129.2, 131.0, 133.7, 135.2, 137.4, 138.4; high resolution mass spectrum (CI, CH_4) m/z 691.3041 (M^+ ; calcd for $\text{C}_{38}\text{H}_{49}\text{SiSO}$, N: 691.2998).

K. Alcohol (-)-I-33.

To a solution of the acetal I-32 (1.45 g, 2.10 mmol) dissolved in CH_2Cl_2 (30 ml) was added DIBAL (1.0 M toluene; 21.0 ml, 21.0 mmol) at 0°C . After stirring for 4 h the mixture was quenched with Rochelle's salt (100 ml) and water (100 ml) and extracted with ethyl acetate (3 x 100 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (20% ethyl acetate/petroleum ether) provided I-33 (1.31 g, 90% yield) as a colorless oil: $[\alpha]_D^{25} -11.6^\circ$ (c 1.12, CHCl_3); IR (CHCl_3) 3080 (w), 3040 (w), 3018 (w), 2960 (s), 2880 (s), 1455 (s), 1375 (s), 1285 (w), 1185 (m), 1179 (s), 1138 (s), 1135 (s), 1090 (s), 1040 (m), 1030 (m), 1020 (m), 885 (m), 810 (w), 680 (m), 600 (m), 570 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.07 (m, 21 H), 1.58 (q, $J = 11.4$ Hz, 1 H), 2.10 (br s, 1 H), 2.45 (dt, $J = 12.3, 4.8$ Hz, 1 H), 2.98 (m, 2 H), 3.42 (m, 1 H), 3.52 (m, 1 H), 3.59 (m, 1 H), 3.86 (m, 2 H), 4.13 (dt, $J = 9.2, 7.7$ Hz, 1 H), 4.32 (d, $J = 7.3$ Hz, 1 H), 4.54 (d, $J = 1.5$ Hz, 1 H), 4.64 (d, $J = 11.5$ Hz, 1 H), 7.24 (m, 1 H), 7.28 (s,

- 60 -

1 H), 7.24-7.39 (m, 5 H), 7.43 (m, 2 H), 7.48 (d, $J = 7.8$ Hz, 1 H), 7.51 (m, 2 H), 7.87 (m, 2 H), 8.00 (d, $J = 8.3$ Hz, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 12.6, 18.2, 25.8, 38.2, 62.6, 68.5, 69.5, 71.8, 72.2, 78.2, 104.8, 113.9, 19.3, 119.6, 123.5, 123.9, 124.8, 127.0, 128.0, 128.5, 129.8, 131.2, 133.9, 135.4, 138.1, 138.5; high resolution mass spectrum (Cl , NH_3) m/z 693.3167 (M^+ ; calcd for $\text{C}_{38}\text{H}_{51}\text{SiSO}_7$, N: 693.3155).

L. Azide (-)-I-35.

6-Azidohexyl triflate I-34 was prepared as follows:
A stirred solution of 6-azido-1-hexanol (0.17 g, 1.17 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.24 g, 1.17 mmol) in dry dichloromethane (10 ml) was treated with triflic anhydride (0.19 ml, 1.17 mmol). After 10 min at room temperature, the mixture was diluted with water (50 ml) and extracted with dichloromethane (3 x 25 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction. To a solution of alcohol I-33 (0.54 g, 0.78 mmol) in dry CH_2Cl_2 (30 ml) at 0°C was added NaH (60%, 0.050 g, 1.17 mmol) and 15-crown-5 (5 mg). After stirring for 20 minutes, triflate 34 (0.32 g, 1.17 mmol) as a solution in CH_2Cl_2 (2 ml) was added via cannula. The mixture was stirred for an additional 24 h, quenched with water (30 ml) and the layers were separated. The aqueous layer was further extracted with CH_2Cl_2 (3 x 20 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (12% ethyl acetate/petroleum ether) provided I-35 (0.57 g, 89% yield) as a colorless oil: $[\alpha]_{\text{D}^{25}} -14.6^\circ$ (c 1.22, CHCl_3); IR (CHCl_3) 3075 (w), 3017 (w), 2955 (s), 2880 (s), 2105 (s), 1450 (m), 1375 (m), 1275 (br, w), 1180 (s), 1125 (s), 1097 (s), 1070 (s), 975 (w), 885 (w), 810 (w), 670 (br, w), 600 (m), 570 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.03 (s, 21 H), 1.31 (m, 4 H), 1.51 (m, 5 H), 2.40 (dt, $J = 12.3, 4.7$ Hz, 1 H), 2.98 (t, $J = 7.2$ Hz, 2 H), 3.15 (t, $J = 6.9$ Hz, 2 H), 3.40 (m, 4 H), 3.56 (m, 2 H), 3.77 (m, 2 H), 4.09 (m, 1 H), 4.24 (d, $J = 7.3$ Hz, 1 H), 4.48 (d, $J = 11.6$ Hz, 1 H), 4.59 (d, $J = 11.6$ Hz, 1 H),

- 61 -

7.22 (t, $J = 7.6$ Hz, 1 H), 7.25 (s, 1 H), 7.30 (m, 5 H), 7.40 (m, 3 H), 7.47 (m, 2 H), 7.84 (d, $J = 7.9$ Hz, 2 H), 7.96 (d, $J = 8.4$ Hz, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 12.4, 18.0, 25.6, 25.7, 26.6, 28.8, 29.5, 38.3, 51.4, 68.5, 69.4, 70.1, 71.4, 71.5, 72.3, 78.1, 105.6, 113.7, 119.4, 119.7, 123.1, 123.4, 124.7, 126.8, 127.8, 128.4, 129.3, 131.1, 133.6, 135.2, 138.2, 138.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 853.3835[(M + Cl) $^+$]; calcd for $\text{C}_{44}\text{H}_{62}\text{SiSO}_7$, N_4Cl : 853.3797).

10 **M. Alcohol (-)-I-36.**

A solution of azide I-35 (0.18 g 0.22 mmol) in THF (3 ml) was cooled to 0°C and TBAF (0.26 ml, 1.00 M, 0.26 mmol) was added dropwise. The mixture was stirred for 2 h, added to water and extracted with EtOAc (3 x 20 ml). The combined
15 extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (12% ethyl acetate/petroleum ether) yielded the alcohol as a colorless oil (0.14 g, 99%); $[\alpha]_D^{25}$ -8.8° (c 1.1, CHCl_3); IR (CHCl_3) 3002 (w), 2940 (m), 2870 (m), 2100 (s), 1450 (s), 1370 (s), 1280
20 (w), 1172 (s), 1130 (s), 1120 (s), 1100 (s), 1088 (s), 1070 (s), 970 (w), 600 (m), 570 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.38 (m, 4 H), 1.59 (m, 5 H), 2.10 (br s, 1 H), 2.47 (dt, $J = 12.5$, 4.5 Hz, 1 H), 3.01 (m, 2 H), 3.23 (t, $J = 6.9$ Hz, 2 H), 3.45 (m, 3 H), 3.59 (m, 5 H), 3.70 (dd, $J = 12.3$, 4.3 Hz, 1 H), 3.77 (dt, $J = 9.5$, 7.0 Hz, 1 H), 4.21 (dt, $J = 9.6$, 6.5
25 Hz, 1 H), 4.29 (d, $J = 6.9$ Hz, 1 H), 4.51 (d, $J = 11.5$ Hz, 1 H), 4.63 (d, $J = 11.6$ Hz, 1 H), 7.26 (t, $J = 7.7$ Hz, 1 H), 7.28 (s, 1 H), 7.35 (m, 5 H), 7.45 (m, 3 H), 7.53 (m, 2 H), 7.89 (m, 2 H), 8.00 (d, $J = 7.9$ Hz, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 25.5, 25.7, 26.5, 28.7, 29.4, 33.8, 51.3, 68.3, 68.6, 70.0, 71.3, 71.4, 72.1, 77.7, 104.5, 113.8, 119.4, 119.9, 123.1, 123.5, 124.8, 126.7, 127.7, 127.8, 128.4, 129.2, 131.0, 133.7, 135.2, 137.9, 138.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 662.2811(M^+); calcd for
30 $\text{C}_{35}\text{H}_{42}\text{SO}_7$, N_4 : 662.2774).

- 62 -

N. Mmt-Chloromethylimidazole (I-37).

To a solution of the chloromethylimidazole (0.20 g, 1.30 mmol) and MmtCl (0.82 g, 2.65 mmol) in dichloromethane at 0°C was rapidly added Hunig's base (0.51 ml, 2.91 mmol).
5 After stirring for 0.5 h the mixture was added to water and the layers were separated. The aqueous layer was further extracted with dichloromethane (2 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (30% ethyl
10 acetate/petroleum ether) yielded I-37 as a colorless oil (0.24 g, 47%) which was used immediately in the next reaction; IR (CHCl₃) 3095 (w), 3060 (w), 3005 (m), 2960 (m), 2840 (w), 1610 (m), 1586 (w), 1510 (s), 1487 (m), 1463 (m), 1445 (m), 1300 (w), 1255 (s), 1180 (m), 1155 (m), 1120 (m), 1085 (w), 1031
15 (m), 990 (w), 905 (w), 825 (m), 695 (m); ¹H NMR (500 MHz, CDCl₃) δ 3.79 (s, 3 H), 4.56 (s, 2 H), 6.84 (m, 3 H), 7.05 (d, J = 8.8 Hz, 2 H), 7.10 (m, 4 H), 7.32 (m, 6 H), 7.39 (br s, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 39.9, 55.2, 75.1, 113.3, 120.2, 128.0, 129.6, 131.1, 134.1, 137.4, 139.2, 147.4, 159.1.

20 O. Imidazole-Azide (+)-I-38.

To a solution of alcohol I-36 (0.20 g, 0.31 mmol) in dry THF (4 ml) at 0°C was added NaHMDS (0.6 M toluene, 0.56 ml, 0.34 mmol). After 10 minutes, chloro-imidazole I-37 (0.24 g, 0.62 mmol) as a solution in THF (5 ml) was added via
25 cannula. After stirring for 48 h at room temperature, the mixture was added to water and extracted with CH₂Cl₂ (3 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (toluene/ethyl acetate/methanol, 7.7:2.0:0.3)
30 provided I-38 (0.071 g, 23% yield) as a colorless oil: [α]_D²⁵ +1.4° (c 0.86, CHCl₃); IR (CHCl₃) 3009, (m), 2965 (m), 2880 (m), 2110 (s), 1610 (w), 1510 (m), 1455 (m), 1375 (m), 1260 (m), 1180 (s), 1135 (s), 1125 (s), 1090 (s), 1075 (s), 1040 (m), 830 (w), 700 (w), 600 (m), 570 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.32 (m, 4 H), 1.55 (m, 5 H), 2.58 (m, 1 H), 2.92 (t, J = 7.2 Hz, 2 H), 3.17 (t, J = 6.9 Hz, 2 H), 3.35 (m, 1 H),
35 3.42 (m, 3 H), 3.46 (dt, J = 9.5, 6.5 Hz, 1 H), 3.55 (dd, J

- 63 -

= 10.7, 5.1 Hz, 1 H), 3.75 (m, 2 H), 3.76 (s superimposed on a m, 3 H), 4.11 (dt, $J = 9.6, 7.1$ Hz, 1 H), 4.37 (d, $J = 7.6$ Hz, 1 H), 4.40 (d, $J = 11.4$ Hz, 1 H), 4.46 (d, $J = 12.1$ Hz, 1 H), 4.58 (d, $J = 11.4$ Hz, 1 H), 4.65 (d, $J = 12.2$ Hz, 1 H),
5 6.76 (s, 1 H), 6.80 (m, 2 H), 7.04 (m, 2 H), 7.11 (m, 4 H), 7.15 (m, 1 H), 7.21-7.37 (m, 15 H), 7.45 (m, 3 H), 7.81 (m, 2 H), 7.93 (d, $J = 8.3$ Hz, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 25.6, 25.7, 26.5, 28.7, 29.5, 34.9, 51.3, 55.2, 66.9, 68.3, 70.0, 71.1, 71.5, 72.4, 74.9, 75.2, 78.0, 105.1, 113.2, 113.6,
10 119.5, 120.0, 123.0, 123.6, 126.7, 127.7, 127.9, 128.0, 128.3, 129.0, 129.6, 131.0, 131.1, 133.6, 134.5, 135.1, 138.1, 138.5, 139.0, 142.3, 159.0; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 1015.4496[(M + H) $^+$; calcd for $\text{C}_{59}\text{H}_{63}\text{SO}_8$ N_6 : 1015.4496].

15 P. Amine (+)-I-39.

To a solution of azide I-38 (0.071 g, 0.070 mmol) in THF (5 ml) was added H_2O (0.059 ml, 3.30 mmol) and PPh_3 (0.046 g, 0.17 mmol) and the reaction mixture heated to 55°C for 10 h, cooled, and concentrated in vacuo. Flash chromatography
20 (15% methanol/methylene chloride) provided I-39 as a colorless oil (62 mg, 90%); $[\alpha]_{\text{D}}^{25} +1.8^\circ$ (c 1.24, CHCl_3); IR (CHCl_3) 3300 (br, w), 3080 (w), 3005 (w), 2940 (m), 2880 (m), 1605 (w), 1510 (w), 1450 (m), 1375 (m), 1290 (w), 1255 (w), 1175 (s), 1130 (s), 1120 (s), 1095 (s), 1085 (s), 830 (w), 595 (m), 565
25 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.29 (m, 4 H), 1.42 (m, 2 H), 1.54 (m, 3 H), 2.56 (m, 1 H), 2.67 (t, $J = 7.1$ Hz, 2 H), 2.92 (t, $J = 7.1$ Hz, 2 H), 3.23 (br s, 2 H), 3.33 (m, 1 H), 3.42 (m, 4 H), 3.56 (dd, $J = 10.7, 4.5$ Hz, 1 H), 3.70 (d, $J = 10.6$ Hz, 1 H), 3.76 (m, 1 H), 3.76 (s superimposed on a m, 3 H), 4.13 (dt, $J = 9.5, 7.1$ Hz, 1 H), 4.37 (d, $J = 7.5$ Hz, 1 H), 4.40 (d, $J = 11.1$ Hz, 1 H), 4.46 (d, $J = 12.1$ Hz, 1 H), 4.57 (d, $J = 11.4$ Hz, 1 H), 4.64 (d, $J = 12.1$ Hz, 1 H), 6.76 (s, 1 H), 6.80 (m, 2 H), 7.04 (m, 2 H), 7.15 (t, $J = 7.8$ Hz, 1 H), 7.20-7.46 (m, 18 H), 7.81 (dd, $J = 8.2, 0.9$ Hz, 2 H),
30 7.91 (d, $J = 8.3$ Hz, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 25.5, 25.9, 26.5, 29.5, 29.7, 34.9, 55.2, 66.8, 58.3, 70.0, 71.1, 71.6, 72.3, 74.9, 75.2, 78.0, 105.1, 113.2, 113.6, 119.5,

- 64 -

119.9, 120.1, 123.1, 124.6, 126.7, 127.7, 127.7, 127.9, 128.0, 128.4, 129.1, 129.7, 131.1, 131.1, 133.6, 134.4, 135.0, 138.1, 138.3, 138.4, 139.0, 142.7, 159.1; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 989.4483[(M + H)⁺];
5 calcd for C₅₉H₆₅SO₈N₄: 989.4522].

Q. Free imidazole (+)-I-16.

To a solution of amine I-39 (0.020 g, 0.020 mmol) in EtOH (3 ml) was added 5M NaOH (0.50 ml) and the mixture was heated at reflux for 4 h. After cooling, the mixture was
10 diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (15% methanol/methylene chloride) afforded the amine (11 mg, 63%) as a pale yellow oil; $[\alpha]_D^{25}$
15 +10.1° (c 0.54, CHCl₃); IR (CHCl₃) 3480 (w), 3500-2700 (br, w), 3060 (w), 3005 (m), 2955 (s), 2860 (m), 1605 (w), 1505 (m), 1450 (m), 1290 (w), 1255 (m), 1180 (w), 1155 (w), 1128 (s), 1075 (br, s), 1030 (s), 820 (w), 690 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.25-1.45 (m, 6 H), 1.55 (m, 3 H), 2.33 (br s, 2 H), 2.59 (m, 1 H), 2.65 (t, J = 7.1 Hz, 2 H), 3.08 (m, 2 H),
20 3.38 (m, 1 H), 3.40-3.56 (m, 6 H), 3.58 (dd, J = 10.8, 5.2 Hz, 1 H), 3.75 (d, J = 9.6 Hz, 1 H), 3.81 (s superimposed on a m, 3 H), 3.81 (m, 1 H), 4.24 (m, 2 H), 4.45 (m, 2 H), 4.49 (d, J = 11.9 Hz, 1 H), 4.60 (d, J = 11.4 Hz, 1 H), 4.66 (d, J =
25 12.0 Hz, 1 H), 6.75 (s, 1 H), 6.85 (m, 2 H), 7.05-7.18 (m, 8 H), 7.25 (d, J = 8.2 Hz, 1 H), 7.27-7.38 (m, 12 H), 7.42 (s, 1 H), 7.57 (d, J = 7.9 Hz, 1 H), 8.41 (br s, 1 H); ¹³C NMR (125.8 MHz, CDCl₃) δ 24.6, 24.9, 25.6, 28.6, 31.8, 33.9, 40.6, 54.2, 65.8, 68.5, 69.0, 70.5, 71.4, 74.1, 76.9, 104.0, 110.0,
30 111.6, 112.2, 117.6, 119.0, 120.6, 121.3, 126.6, 126.9, 126.9, 127.3, 128.6, 130.1, 133.4, 135.0, 137.0, 137.5, 137.9, 141.7, 158.0; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 849.4672[(M + H)⁺]; calcd for C₅₃H₆₁O₆N₄: 849.4591].

To a solution of the amine (0.023 g, 0.027 mmol) in dry
35 CH₂Cl₂ (2 ml) was added TFA (3.5 ml, 0.045 mmol). After stirring for 5 minutes, the mixture was added to brine (20 ml) that had been adjusted to pH 8.0 with aqueous sodium

- 65 -

bicarbonate and extracted with methylene chloride (3 x 15 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Purification by RP HPLC (water/acetonitrile) afforded I-16
5 (9.7 mg, 63%) as a pale yellow oil; $[\alpha]_D^{25} +11.2^\circ$ (c 0.42, CH₃OH); ¹H NMR (500 MHz, CD₃OD) δ 1.26-1.42 (m, 5 H), 1.52 (m, 4 H), 2.44 (m, 1 H), 2.78 (t, $J = 6.6$ Hz, 2 H), 3.16 (m, 1 H), 3.36-3.48 (m, 4 H), 3.52 (dd, $J = 10.9, 4.8$ Hz, 1 H), 3.64 (dd, $J = 11.1, 1.5$ Hz, 1 H), 3.79 (dt, $J = 9.4, 7.3$ Hz, 1 H),
10 4.15 (dt, $J = 9.4, 6.1$ Hz, 1 H), 4.35 (m, 2 H), 4.39 (d, $J = 11.7$ Hz, 1 H), 4.42 (d, $J = 13.0$ Hz, 1 H), 4.53 (d, $J = 11.6$ Hz, 1 H), 6.92 (m, 2 H), 6.99 (m, 1 H), 7.02 (s, 1 H), 7.25 (m, 6 H), 7.49 (d, $J = 7.9$ Hz, 1 H), 8.64 (s, 1 H); ¹³C NMR (62.9 MHz, CD₃OD) δ 26.8, 26.9, 27.2, 28.5, 30.5, 35.8, 40.6,
15 62.8, 71.0, 72.3, 72.5, 73.2, 76.8, 79.1, 106.2, 112.2, 113.0, 117.9, 119.5, 119.6, 122.3, 123.8, 128.8, 128.9, 129.4, 132.5, 135.3, 138.0, 139.6; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 577.3421[(M + H)⁺; calcd for C₃₃H₄₅O₅N₄: 577.3390].

20 **R. Amide (+)-I-40.**

To a solution of amine I-39 (0.043 g, 0.043 mmol) in methylene chloride (1 ml) and methanol (2 ml) was added acetic anhydride (4.4 ml, 0.043 mmol). After 2 h, two additional equivalents of acetic anhydride (8.8 ml) were added and
25 stirring was continued for a total of 24 h. The mixture was diluted with methylene chloride (15 ml) and washed sequentially with saturated sodium bicarbonate and water. The organic layer was dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (15%
30 methanol/methylene chloride) afforded two inseparable components which were used uncharacterized in the following reaction.

To a solution of the above sugars in ethanol (4 ml) was added 5N NaOH (0.200 ml) and mixture was heated at reflux for
35 2 h. After cooling, the mixture was diluted with water, adjusted to pH 8.0 with HCl, and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with

- 66 -

brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Purification by RP HPLC (water/acetonitrile) afforded I-40 (10 mg, 38%) as a colorless oil; $[\alpha]_D^{25} +13.4^\circ$ (c 0.62, C_2H_5OH); 1H NMR (500 MHz, CD_3OD) δ 1.21-1.35 (m, 4 H), 1.36 (m, 3 H), 1.48 (m, 2 H), 1.83 (s, 3 H), 2.40 (dt, $J = 12.2, 4.7$ Hz, 1 H), 2.99 (t, $J = 6.7$ Hz, 2 H), 3.03 (t, $J = 7.1$ Hz, 2 H), 3.15 (m, 1 H), 3.35-3.46 (m, 4 H), 3.51 (dd, $J = 10.9, 4.9$ Hz, 1 H), 3.63 (dd, $J = 11.0, 1.7$ Hz, 1 H), 3.79 (dt, $J = 9.4, 7.3$ Hz, 1 H), 4.13 (dt, $J = 9.3, 6.1$ Hz, 1 H), 4.33 (d, $J = 7.5$ Hz, 1 H), 4.37 (d, $J = 13.0$ Hz, 1 H), 4.39 (d, $J = 11.7$ Hz, 1 H), 4.42 (d, $J = 13.0$ Hz, 1 H), 4.52 (d, $J = 11.6$ Hz, 1 H), 6.92 (m, 2 H), 6.99 (dt, $J = 7.1, 1.0$ Hz, 1 H), 7.02 (s, 1 H), 7.22 (m, 6 H), 7.48 (d, $J = 7.9$ Hz, 1 H), 8.63 (s, 1 H); ^{13}C NMR (125.8 MHz, CD_3OD) δ 22.5, 26.9, 27.0, 30.3, 30.6, 35.8, 40.5, 62.7, 70.8, 70.9, 72.4, 72.6, 73.3, 76.8, 79.2, 106.2, 112.2, 113.1, 118.0, 119.5, 119.6, 122.3, 123.8, 128.8, 128.9, 129.4, 132.5, 135.3, 138.0, 139.7, 173.2; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 619.3521[(M + H).; calcd for $C_{35}H_{47}O_6N_4$: 619.3495].

20 **S. Azide (-)-44.**

1-Iodo-6-azido-2-hexyne I-43 was prepared as follows: To a stirred solution of 6-azido-2-hexyn-1-ol (0.10 g, 0.72 mmol), imidazole (0.059 g, 0.86 mmol), and triphenylphosphine (0.23 g, 0.86 mmol) in Et_2O/CH_3CN (2 ml; 5:3) at $0^\circ C$, was added iodine (0.23 g, 0.86 mmol). After 5 min at room temperature, the mixture was diluted with ether (10 ml) and washed successively with saturated $Na_2S_2O_3$ and $CuSO_4$. The ether layer was dried over magnesium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction.

To a solution of alcohol I-33 (0.13 g, 0.19 mmol) and iodide I-43 (0.13 g, 0.52 mmol) in dry THF (2 ml) at $0^\circ C$ was added NaH (60 %, 0.012 g, 0.30 mmol). After stirring for 6 h the mixture was poured into water (30 ml) and extracted with Et_2O (3 x 15 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (15% ethyl acetate/petroleum ether)

- 67 -

provided I-44 (107 mg, 70% yield) as a colorless oil: $[\alpha]_D^{25}$ -15.1° (c 0.72, CHCl₃); IR (CHCl₃) 3075 (w), 3039 (w), 3018 (w), 2958 (s), 2876 (s), 2108 (s), 1452 (m), 1371 (br, m), 1175 (s), 1135 (s), 1122 (s), 1100 (s), 1060 (m), 1020 (w), 882 (w), 810 (w), 670 (br, w), 595 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.03 (s, 21 H), 1.53 (m, 1 H), 1.69 (m, 2 H), 2.24 (tt, J = 7.0, 1.9 Hz, 2 H), 2.40 (m, 1 H), 2.98 (t, J = 7.0 Hz, 2 H), 3.30 (t, J = 6.6 Hz, 2 H), 3.46 (m, 2 H), 3.58 (m, 1 H), 3.72-3.80 (m, 3 H), 4.11 (m, 1 H), 4.14-4.22 (m, 2 H), 4.24 (d, J = 7.3 Hz, 1 H), 4.53 (d, J = 11.5 Hz, 1 H), 4.60 (d, J = 11.6 Hz, 1 H), 7.21 (t, J = 7.4 Hz, 1 H), 7.25 (s, 1 H), 7.26-7.33 (m, 5 H), 7.39 (m, 3 H), 7.48 (m, 2 H), 7.85 (d, J = 7.5 Hz, 2 H), 7.96 (d, J = 7.9 Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 12.4, 16.1, 18.0, 25.6, 27.7, 38.3, 50.2, 59.1, 68.5, 68.7, 69.4, 71.5, 71.9, 77.9, 85.1, 105.6, 113.7, 119.4, 119.7, 123.1, 123.4, 124.7, 126.8, 127.8, 128.4, 129.2, 131.1, 133.6, 135.2, 138.2, 138.4.

T. Alcohol (-)-I-45.

A solution of azide I-44 (0.20 g 0.24 mmol) in THF (5 ml) was cooled to 0°C and TBAF (0.29 ml, 1.0 M, 0.29 mmol) was added dropwise. The mixture was stirred for 2 h, added to water and extracted with Et₂O (3 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (40% ethyl acetate/petroleum ether) yielded the alcohol as a colorless oil (0.16 g, 100%); $[\alpha]_D^{25}$ -12.8° (c 0.39, CHCl₃); IR (CHCl₃) 3050 (w), 3039 (w), 3020 (w), 2945 (m), 2888 (m), 2117 (s), 1455 (s), 1375 (s), 1280 (br, m), 1185 (s), 1140 (s), 1130 (s), 1105 (s), 1093 (s), 1075 (s), 1056 (s), 600 (m), 575 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.52 (m, 1 H), 1.60 (quin., J = 6.8 Hz, 2 H), 2.19 (d, J = 3.1 Hz, 1 H), 2.26 (tt, J = 7.0, 2.0 Hz, 2 H), 2.45 (m, 1 H), 2.98 (m, 2 H), 3.22 (t, J = 6.6 Hz, 2 H), 3.43 (m, 1 H), 3.75 (m, 3 H), 4.11-4.23 (m, 3 H), 4.25 (d, J = 7.1 Hz, 1 H), 4.52 (d, J = 11.5 Hz, 1 H), 4.61 (d, J = 11.5 Hz, 1 H), 7.23 (m, 1 H), 7.27-7.36 (m, 2 H), 7.43 (m, 3 H), 7.50 (m, 2 H), 7.86 (m, 2 H), 7.98 (m, 1 H); ¹³C NMR (125.8 MHz, CDCl₃) δ 16.1, 25.5, 27.7, 34.2, 50.1, 59.1, 68.3,

- 68 -

68.6, 68.6, 71.4, 71.8, 77.7, 85.2, 104.7, 113.8, 119.4, 123.2, 123.5, 124.8, 126.7, 127.7, 127.8, 128.4, 129.2, 131.1, 133.7, 135.2, 138.0, 139.0, 139.2; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 658.2482 (M^+ ; calcd for $C_{15}H_{18}SO_7$, N_4 : 658.2461].

U. Imidazole-Azide (-)-I-46.

To a solution of alcohol I-45 (0.16 g, 0.24 mmol) and chloro-imidazole I-37 (0.27 g, 0.69 mmol) at 0°C in dry THF (4 ml) was added NaH (60%, 0.015 g, 0.36 mmol). After stirring for 12 h at room temperature, the mixture was added to water and extracted with Et_2O (3 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (60% ethyl acetate/petroleum ether) yielded I-46 as a colorless oil (0.13 g, 54%); $[\alpha]_D^{25}$ -0.4° (c 1.14, $CHCl_3$); IR ($CHCl_3$) 3025 (w), 3017 (w), 3010 (m), 2980 (m), 2959 (m), 2880 (m), 2108 (s), 1613 (w), 1590 (w), 1516 (m), 1452 (s), 1385 (s), 1360 (s), 1290 (s), 1280 (s), 1238 (s), 1225 (s), 1100 (s), 1075 (s), 1050 (s), 830 (m), 700 (m), 600 (m), 572 (m) cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 1.51 (m, 1 H), 1.70 (quin., J = 6.8 Hz, 2 H), 2.25 (tt, J = 7.0, 2.1 Hz, 2 H), 2.59 (m, 1 H), 2.92 (t, J = 7.0 Hz, 2 H), 3.29-3.39 (m, 3 H), 3.45 (m, 2 H), 3.69-3.80 (m, 3 H), 3.28 (s superimposed on a m, 3 H), 4.10-4.20 (m, 3 H), 4.37 (d, J = 7.6 Hz, 1 H), 4.44 (d, J = 11.4 Hz, 1 H), 4.46 (d, J = 12.2 Hz, 1 H), 4.59 (d, J = 11.5 Hz, 1 H), 4.64 (d, J = 12.2 Hz, 1 H), 6.79 (m, 3 H), 7.04 (m, 2 H), 7.10 (m, 4 H), 7.15 (m, 1 H), 7.20-7.49 (m, 18 H), 7.82 (m, 2 H), 7.91 (d, J = 8.2 Hz, 1 H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 13.1, 15.0, 24.5, 26.7, 33.9, 49.1, 54.2, 58.1, 65.9, 67.3, 67.6, 70.1, 71.0, 73.8, 74.0, 76.2, 76.7, 84.0, 104.1, 112.2, 112.6, 118.4, 118.8, 122.0, 122.5, 125.5, 125.6, 126.6, 126.8, 126.9, 127.3, 128.1, 128.6, 130.0, 130.1, 132.5, 133.4, 134.1, 137.0, 137.2, 137.4, 137.9, 141.7, 158.0; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 1011.4100 [$(M + H)^+$; calcd for $C_{59}H_{59}SO_8$, N_6 : 1011.4115].

- 69 -

V. Free imidazole (+)-42.

To a solution of azide I-45 (0.11 g, 0.11 mmol) in THF (5 ml) was added H₂O (0.095 ml, 5.27 mmol) and PPh₃ (0.073 g, 0.28 mmol) and the reaction mixture heated to 55°C for 6 h, cooled, and concentrated in vacuo. Flash chromatography (10% methanol/methylene chloride) provided the amine as a colorless oil (103 mg, 93%); $[\alpha]_D^{25} +2.2^\circ$ (c 0.87, CHCl₃); IR (CHCl₃) 3070 (w), 3010 (m), 2960 (m), 2942 (m), 2878 (m), 1612 (m), 1590 (w), 1515 (m), 1452 (m), 1374 (m), 1259 (m), 1179 (s), 1145 (s), 1120 (s), 1090 (s), 1070 (m), 1050 (m), 827 (w), 700 (w), 597 (w), 569 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.52 (m, 1 H), 1.76 (m, 2 H), 2.23 (m, 2 H), 2.55 (m, 1 H), 2.91 (m, 4 H), 3.38 (m, 1 H), 3.47 (m, 2 H), 3.69 (dd, $J = 10.5, 4.8$ Hz, 1 H), 3.77 (m, 5 H), 4.11 (m, 3 H), 4.40 (d, $J = 7.7$ Hz, 1 H), 4.43 (d, $J = 11.4$ Hz, 1 H), 4.44 (d, $J = 12.2$ Hz, 1 H), 4.59 (d, $J = 11.2$ Hz, 1 H), 4.60 (d, $J = 12.2$ Hz, 1 H), 5.60 (br s, 1 H), 6.71 (s, 1 H), 6.78 (m, 2 H), 7.00 (m, 2 H), 7.08 (m, 4 H), 7.12 (m, 1 H), 7.20 (m, 1 H), 7.22-7.38 (m, 14 H), 7.45 (m, 3 H), 7.81 (dd, $J = 8.4, 1.0$ Hz, 2 H), 7.91 (d, $J = 7.6$ Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 16.2, 25.6, 28.0, 34.8, 39.7, 55.3, 59.1, 66.7, 68.4, 68.7, 71.2, 72.2, 75.0, 75.2, 77.4, 77.5, 77.7, 85.1, 105.0, 113.3, 113.6, 119.6, 119.9, 120.1, 123.1, 123.7, 124.6, 126.7, 127.8, 128.0, 128.4, 129.2, 129.7, 131.2, 133.6, 134.4, 135.1, 138.1, 138.3, 138.3, 139.0, 142.6, 159.1; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 985.4254 [(M + H)⁺; calcd for C₂₉H₆₁SO₈ N₄: 985.4210].

To a solution of the amine (0.085 g, 0.087 mmol) in EtOH (3 ml) was added 5M NaOH (0.50 ml) and mixture was heated at reflux for 4 h. After cooling, the mixture was diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (15% methanol/methylene chloride) afforded the amine (39 mg, 56%) as a colorless oil; $[\alpha]_D^{25} +3.1^\circ$ (c 1.95, CHCl₃); IR (CHCl₃) 3480 (w), 3520-2500 (br, w), 3020 (s), 2960 (s), 2940 (s), 2880 (s), 1609 (m), 1590 (w), 1513 (s), 1493

- 70 -

(m), 1459 (m), 1447 (m), 1355 (m), 1340 (m), 1302 (m), 1257 (s), 1185 (m), 1156 (m), 1130 (s), 1090 (s), 1037 (s), 1010 (m), 910 (w), 825 (m), 695 (m), 660 (w), 582 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.50 (q, $J = 11.4$ Hz, 1 H), 1.61 (m, 1 H), 1.69 (m, 1 H), 2.11 (m, 2 H), 2.52 (m, 1 H), 2.79 (m, 2 H), 3.01 (m, 2 H), 3.36-3.49 (m, 3 H), 3.65 (dd, $J = 10.8$, 5.0 Hz, 1 H), 3.77 (m, 5 H), 4.05-4.17 (m, 3 H), 4.40 (m, 2 H), 4.47 (d, $J = 11.9$ Hz, 1 H), 4.54 (d, $J = 11.4$ Hz, 1 H), 4.60 (d, $J = 12.0$ Hz, 1 H), 6.72 (br s, 1 H), 6.80 (apparent d, $J = 9.0$ Hz, 2 H), 6.98-7.13 (m, 9 H), 7.23-7.33 (m, 12 H), 7.40 (br s, 1 H), 7.50 (d, $J = 7.1$ Hz, 1 H), 8.75 (br s, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 16.1, 25.8, 26.0, 34.8, 38.9, 55.3, 59.1, 66.3, 68.7, 70.0, 71.2, 72.2, 75.1, 77.7, 84.7, 105.1, 111.5, 112.2, 113.3, 118.6, 119.0, 120.2, 121.7, 122.8, 128.1, 128.4, 129.7, 131.2, 134.2, 136.2, 138.0, 138.8, 142.5, 159.2; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 845.4261 [(M + H) $^+$; calcd for $\text{C}_{53}\text{H}_{57}\text{SO}_6$ N_4 : 845.4278].

To a solution of the amine (0.040 g, 0.047 mmol) in dry CH_2Cl_2 (2 ml) was added TFA (24 ml, 0.31 mmol). After stirring for 5 minutes, the mixture was added to brine (20 ml) that had been adjusted to pH 8.0 with aqueous sodium bicarbonate and extracted with methylene chloride (3 x 15 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Purification by RP HPLC (water/acetonitrile) afforded I-42 (12.3 mg, 45%) as a pale yellow oil; $[\alpha]_D^{25} +0.9^\circ$ (c 0.56, CH_3OH); ^1H NMR (500 MHz, CD_3OD) δ 1.42 (m, 1 H), 1.78 (apparent quin., $J = 7.0$ Hz, 2 H), 2.30 (tt, $J = 7.0$, 2.1 Hz, 2 H), 2.47 (m, 1 H), 2.95 (t, $J = 7.6$ Hz, 2 H), 3.06 (t, $J = 6.7$ Hz, 2 H), 3.20 (m, 2 H), 3.47 (m, 2 H), 3.69 (dd, $J = 10.7$, 4.4 Hz, 1 H), 3.76 (dd, $J = 11.0$, 1.4 Hz, 1 H), 3.86 (dt, $J = 9.4$, 7.3 Hz, 1 H), 4.18 (m, 3 H), 4.40 (m, 2 H), 4.49 (m, 2 H), 4.60 (d, $J = 11.6$ Hz, 1 H), 6.98 (m, 2 H), 7.06 (m, 1 H), 7.09 (s, 1 H), 7.26-7.34 (m, 7 H), 7.55 (d, 7.8 Hz, 1 H), 8.70 (br s, 1 H); ^{13}C NMR (62.9 MHz, CD_3OD) δ 16.6, 26.9, 27.5, 35.8, 39.8, 59.7, 62.7, 69.6, 71.0, 72.3, 73.1, 76.8, 78.4, 79.0, 85.6, 106.2, 112.2, 113.1, 117.9, 119.5, 119.6, 122.3, 124.0, 128.8,

- 71 -

128.9, 129.4, 132.5, 135.4, 139.7, 142.2; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.3062 [(M + H)⁺; calcd for C₃₃H₄₁SO₅ N₄: 845.4278].

5 W. Saturated amine I-16 from Pd/CaCO₃ reduction of acetylene-azide I-46.

To a solution of azide I-46 (8 mg) in ethanol (1.7 ml) was added Pd/CaCO₃ (1 mg). The system was evacuated and back flushed with H₂ gas four times and then allowed to stir under an atmosphere of H₂ gas for 2.5 h. The mixture was filtered
10 through celite, the celite was washed with Et₂O (20 ml), and the filtrate was concentrated in vacuo. Flash chromatography (20% methanol/methylene chloride) afforded I-16 (5.1 mg, 64%) as a pale yellow oil which was identical in all respects with material obtained by other methods.

15 X. Benzyl ether (-)-I-50.

To a solution of the alcohol I-33 (0.044 g, 0.063 mmol) and benzyl bromide (8.3 ml, 0.070) in dichloromethane (2 ml) at 0°C was added NaH (60%, 3.0 mg, 0.070 mmol) and 15-crown-5 (1 ml). After stirring for 5 h, the mixture was added to H₂O
20 (200 ml) and extracted with dichloromethane (3 x 10 ml). The combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (10% ethyl acetate/petroleum ether) provided I-50 (0.035 g, 70% yield) as a colorless oil: $[\alpha]_D^{25}$ -10.2° (c 3.0, CHCl₃); IR
25 (CHCl₃) 3065 (w), 3010 (m), 2950 (s), 2877 (s), 1610 (w), 1496 (w), 1465 (m), 1452 (s), 1370 (s), 1270 (w), 1205 (m), 1175 (s), 1125 (s), 1098 (s), 1070 (s), 880 (m), 725 (br, s), 665 (s), 595 (m), 569 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.04 (s, 21 H), 1.51 (apparent q, J = 11.1 Hz, 1 H), 2.41 (dt, J =
30 12.3, 4.8 Hz, 1 H), 3.00 (t, J = 7.9 Hz, 3 H), 3.43-3.52 (m, 2 H), 3.60 (m, 1 H), 3.66 (dd, 10.7, 5.1 Hz, 1 H), 3.78 (m, 2 H), 4.12 (m, 1 H), 4.26 (d, J = 7.3 Hz, 1 H), 4.43 (d, J =
11.5 Hz, 1 H), 4.53 (d, J = 7.0 Hz, 1 H), 4.57 (d, J = 6.2 Hz, 1 H), 4.59 (d, J = 12.3 Hz, 1 H), 7.18-7.32 (m, 12 H),
35 7.36-7.41 (m, 3 H), 7.47 (m, 2 H), 7.83 (m, 2 H), 7.97 (apparent d, J = 8.4 Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 12.4, 18.0, 25.7, 38.3, 68.5, 69.4, 71.4, 72.2, 73.5, 78.1,

- 72 -

105.6, 113.7, 119.4, 119.7, 123.1, 123.4, 124.7, 126.7, 127.5, 127.7, 128.3, 128.4, 129.2, 131.1, 133.6, 135.2, 138.1, 138.4; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 783.3662 (M^+ ; calcd for $C_{15}H_{17}SiSO$, N: 783.3625).

5 **Y. Alcohol (-)-I-51.**

A solution of benzyl ether I-50 (0.080 g 0.10 mmol) in THF (2 ml) was cooled to 0°C and TBAF (0.11 ml, 1.0 M, 0.11 mmol) was added dropwise. The mixture was stirred for 2 h, added to water and extracted with EtOAc (3 x 20 ml). The
10 combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (40% ethyl acetate/petroleum ether) yielded alcohol I-51 as a colorless oil (0.070 g, 100%); $[\alpha]_D^{25}$ -7.7° (c 0.27, $CHCl_3$); IR ($CHCl_3$) 3080 (w), 3040 (w), 3010 (m), 2955 (m), 2880 (m),
15 1450 (m), 1370 (m), 1280 (w), 1173 (s), 1120 (s), 1100 (s), 1060 (s), 690 (w), 680 (w) cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 1.53 (apparent q, J = 10.2 Hz, 1 H), 2.30 (br s, 1 H), 2.45 (dt, J = 12.4, 4.6 Hz, 1 H), 2.99 (m, 2 H), 3.46 (m, 1 H), 3.55 (m, 1 H), 3.59 (m, 1 H), 3.65 (dd, J = 10.5, 5.0 Hz, 1 H),
20 3.74 (m, 2 H), 4.20 (dt, J = 9.5, 6.5 Hz, 1 H), 4.27 (d, J = 6.9 Hz, 1 H), 4.43 (d, J = 11.5 Hz, 1 H), 4.49-4.58 (m, 3 H), 7.21-7.33 (m, 12 H), 7.39 (m, 2 H), 7.43 (s, 1 H), 7.49 (d, 2 H), 7.85 (m, 2 H), 7.99 (d, J = 8.4 Hz, 1 H); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 24.5, 32.8, 67.2, 67.5, 68.3, 70.2, 71.0, 72.3,
25 76.7, 103.5, 112.7, 118.4, 118.8, 122.1, 122.4, 123.7, 125.6, 126.5, 126.6, 126.7, 126.7, 127.3, 127.4, 128.1, 130.0, 132.6, 134.1, 136.8, 137.2; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 628.2335 [$(M + H)^+$; calcd for $C_{36}H_{38}SO$, N: 628.2368].

30 **Z. Azide (-)-I-53.**

1-Iodo-5-azido-2-pentyne I-52 was prepared as follows: To a stirred solution of 5-azido-2-pentyn-1-ol (0.13 g, 1.00 mmol), imidazole (0.085 g, 1.25 mmol), and triphenylphosphine (0.32 g, 1.25 mmol) in Et_2O/CH_3CN (2 ml; 5:3) at 0°C, was
35 added iodine (0.32 g, 1.25 mmol). After 5 min at room temperature, the mixture was diluted with ether (10 ml) and washed successively with saturated $Na_2S_2O_3$ and $CuSO_4$. The

- 73 -

ether layer was dried over magnesium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction.

To a solution of alcohol I-51 (0.073 g, 0.12 mmol) and
5 iodide I-52 (0.24 g, 1.00 mmol) in dry dichloromethane (2 ml)
at 0°C was added NaH (60 %, 6.0 mg, 0.15 mmol). After
stirring for 6 h the mixture was poured into water (30 ml) and
extracted with CH₂Cl₂ (3 x 15 ml). The combined extracts were
washed with water, dried over magnesium sulfate and
10 concentrated in vacuo. Flash chromatography (30% ethyl
acetate/petroleum ether) provided I-53 (64 mg, 75% yield) as
a colorless oil: $[\alpha]_D^{25}$ -8.9° (c 0.63, CHCl₃); IR (CHCl₃) 3070
(w), 3028 (w), 3010 (m), 2940 (m), 2870 (m), 2110 (s), 1450
(s), 1378 (s), 1270 (m), 1250 (s), 1178 (s), 1133 (s), 1120
15 (s), 1090 (s), 1072 (s), 1045 (s), 690 (w), 595 (m) cm⁻¹; ¹H
NMR (500 MHz, CDCl₃) δ 1.48 (q, J = 12.0 Hz, 1 H), 2.50 (tt,
J = 6.9, 2.1 Hz, 2 H), 2.55 (dt, J = 12.2, 4.7 Hz, 1 H), 3.00
(t, J = 6.9 Hz, 2 H), 3.39 (m, 3 H), 3.50 (m, 2 H), 3.66 (dd,
J = 10.7, 5.0 Hz, 1 H), 3.78 (m, 2 H), 4.19 (m, 1 H), 4.24
20 (tq, J = 15.2, 2.2 Hz, 2 H), 4.34 (d, J = 7.6 Hz, 1 H), 4.40
(d, J = 11.4 Hz, 1 H), 4.58 (m, 3 H), 7.19-7.31 (m, 12 H),
7.39 (m, 2 H), 7.48 (m, 4 H), 7.85 (m, 2 H), 7.98 (d, J = 8.3
Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 19.9, 25.6, 34.8, 49.7,
58.3, 68.3, 69.2, 71.3, 72.1, 73.4, 74.3, 78.0, 78.5, 82.4,
25 104.9, 113.7, 123.1, 123.6, 124.7, 126.7, 127.5, 127.7, 128.3,
128.4, 129.2, 131.0, 133.6, 135.1, 137.9, 138.3; high
resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z
735.2827 [(M + H)⁺; calcd for C₄₁H₄₃SO, N₄: 735.2852].

AA. Amine (-)-I-54.

30 To a solution of azide I-53 (0.021 g, 0.027 mmol) in
THF (1.5 ml) was added H₂O (0.012 ml, 0.69 mmol) and PPh₃
(0.014 g, 0.055 mmol) and the reaction mixture was heated to
55°C for 4 h, cooled, and concentrated in vacuo. Flash
chromatography (6% methanol/methylene chloride) provided I-54
35 as a colorless oil (16.2 mg, 83%); $[\alpha]_D^{25}$ -9.0° (c 0.81,
CHCl₃); IR (CHCl₃) 3070 (w), 3038 (w), 3017 (w), 2940 (m),
2878 (w), 1451 (m), 1370 (br, m), 1210 (s), 1187 (m), 1179

- 74 -

(m), 1122 (m), 1090 (m), 1072 (m), 930 (w), 750 (br, s), 665 (s), 595 (m), 569 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.48 (q, $J = 12.0$ Hz, 1 H), 2.00 (br s, 2 H), 2.36 (br s, 2 H), 2.55 (dt, $J = 12.3, 4.7$ Hz, 1 H), 2.83 (br s, 2 H), 3.00 (t, $J = 6.7$ Hz, 2 H), 3.39 (m, 1 H), 3.50 (m, 2 H), 3.66 (dd, $J = 10.8, 5.0$ Hz, 1 H), 4.19 (m, 1 H), 4.25 (tq, $J = 15.3, 2.1$ Hz, 2 H), 4.35 (d, $J = 7.6$ Hz, 1 H), 4.40 (d, $J = 11.4$ Hz, 1 H), 4.56 (m, 3 H), 7.20-7.33 (m, 12 H), 7.38 (m, 3 H), 7.48 (m, 4 H), 7.85 (m, 2 H), 7.97 (d, $J = 8.4$ Hz, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 25.6, 34.8, 58.4, 68.3, 69.2, 71.3, 72.1, 73.4, 74.3, 77.9, 80.0, 84.2, 104.9, 113.7, 119.4, 119.8, 123.7, 124.7, 126.7, 127.5, 127.7, 127.7, 127.8, 128.3, 128.4, 129.1, 131.0, 133.6, 135.1, 137.9, 138.3, 138.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 709.2980 [(M + H) $^+$]; calcd for $\text{C}_{41}\text{H}_{45}\text{SO}_7$, N_2 : 709.2947].

AB. Amine (+)-I-48.

To a solution of amine I-54 (0.012 g, 0.017 mmol) in MeOH (1.5 ml) was added 5M KOH (0.30 ml) and the mixture was heated at reflux for 8 h. After cooling, the mixture was diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (8% methanol/methylene chloride) afforded I-48 (7.1 mg, 73%) as a pale yellow oil; $[\alpha]_{\text{D}}^{25} +13.5^\circ$ (c 0.31, CHCl_3); IR (CHCl_3) 3480 (m), 3010 (s), 2930 (s), 2879 (s), 2861 (s), 1460 (m), 1270 (w), 1140 (w), 1105 (m), 1079 (s), 861 (w), 690 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.49 (q, $J = 11.9$ Hz, 1 H), 2.38 (br s, 2 H), 2.51 (dt, $J = 12.3, 4.6$ Hz, 2 H), 2.66 (br s, 2 H), 2.84 (br s, 2 H), 3.09 (t, $J = 6.7$ Hz, 2 H), 3.33 (m, 1 H), 3.45-3.55 (m, 2 H), 3.66 (dd, $J = 10.7, 4.9$ Hz, 1 H), 3.76 (dd, $J = 10.9, 1.8$ Hz, 1 H), 3.83 (dt, $J = 9.4, 7.2$ Hz, 1 H), 4.17 (dt, $J = 15.0, 2.0$ Hz, 1 H), 4.24 (dt, $J = 9.5, 6.3$ Hz, 1 H), 4.29 (dt, $J = 15.0, 2.1$ Hz, 1 H), 4.39 (m, 2 H), 4.56 (m, 3 H), 7.08 (t, $J = 7.9$ Hz, 1 H), 7.12 (s, 1 H), 7.16 (t, $J = 7.3$ Hz, 1 H), 7.20-7.35 (m, 11 H), 7.58 (d, $J = 7.8$ Hz, 1 H), 8.63 (br s, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 22.7, 29.7, 34.9, 58.6, 69.1, 69.5, 71.3, 72.2,

- 75 -

73.5, 74.4, 77.9, 78.4, 83.6, 104.9, 111.1, 112.7, 118.7, 119.1, 121.7, 122.5, 127.6, 127.8, 128.3, 128.4, 136.2, 138.0, 138.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 569.3029 [(M + H)⁺; calcd for C₁₅H₁₁O₅ N₂: 569.3015].

AC. Amide (+)-I-56.

To a solution of amine I-54 (8.4 mg, 0.012 mmol) in CH₂Cl₂ (1 ml) at 0°C was added Et₃N (1.8 ml, 0.013 mmol) and Ac₂O (1.2 ml, 0.013 mmol). After stirring for one minute, the mixture was poured into water. The aqueous layer was extracted with methylene chloride (3 x 20 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. ¹H NMR (500 MHz, CDCl₃, Crude) δ 1.47 (q, J = 11.9 Hz, 1 H), 1.95 (s, 3 H), 2.40 (m, 2 H), 2.53 (dt, J = 12.2, 4.8 Hz, 1 H), 2.99 (m, 2 H), 3.28 (m, 1 H), 3.39 (m, 2 H), 3.49 (m, 2 H), 3.65 (dd, J = 10.7, 5.0 Hz, 1 H), 3.78 (m, 2 H), 4.22 (m, 3 H), 4.34 (d, J = 7.6 Hz, 1 H), 4.40 (d, J = 11.4 Hz, 1 H), 4.56 (m, 3 H), 6.05 (br s, 1 H), 7.20-7.33 (m, 12 H), 7.39 (apparent t, J = 8.2 Hz, 2 H), 7.48 (m, 3 H), 7.85 (m, 2 H), 7.95 (d, J = 8.2 Hz, 1 H).

To a solution of the crude amide in MeOH (1 ml) was added 5M KOH (0.20 ml) and mixture was heated at reflux for 6 h. After cooling, the mixture was diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (2% methanol/methylene chloride) afforded I-56 (4.9 mg, 68% from I-54) as a pale yellow oil; $[\alpha]_D^{25}$ +18.4° (c 0.25, CHCl₃); IR (CHCl₃) 3480 (w), 3010 (m), 2940 (m), 2870 (m), 1675 (s), 1520 (w), 1456 (m), 1367 (w), 1250 (br, w), 1285 (br, s), 695 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.50 (q, J = 11.9 Hz, 1 H), 1.95 (s, 3 H), 2.41 (m, 2 H), 2.51 (dt, J = 12.2, 4.7 Hz, 1 H), 3.10 (t, J = 7.0 Hz, 2 H), 3.27 (m, 1 H), 3.40 (m, 2 H), 3.45-3.55 (m, 2 H), 3.66 (dd, J = 10.8, 4.9 Hz, 1 H), 3.76 (dd, J = 10.8, 1.8 Hz, 1 H), 3.85 (dt, J = 9.5, 7.3 Hz, 1 H), 4.19-4.29 (m, 3 H), 4.37 (d, J = 7.6 Hz, 1 H), 4.41 (d, J = 11.4 Hz, 1 H), 4.53-4.61 (m, 3 H), 7.09 (m, 2 H), 7.16 (m, 1 H),

- 76 -

7.20-7.36 (m, 11 H), 7.59 (d, $J = 8.2$ Hz, 1 H), 8.15 (br s, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 19.9, 23.9, 25.8, 34.8, 38.2, 58.3, 69.1, 69.6, 71.3, 72.2, 73.4, 74.1, 78.0, 78.1, 83.7, 104.8, 11.1, 112.7, 118.7, 119.2, 121.9, 122.2, 127.5, 127.7, 128.3, 128.4, 136.2, 138.0, 170.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 633.2923 [$(\text{M} + \text{Na})^+$; calcd for $\text{C}_{37}\text{H}_{42}\text{O}_6 \text{N}_2\text{Na}$: 633.2940].

AD. Alkane (+)-I-47.

To a solution of azide I-53 (0.020 g, 0.027 mmol) in EtOH (1 ml) was added 5% Pd/ CaCO_3 (6 mg, 33 wgt. %). The system was evacuated and back flushed with H_2 gas four times and then allowed to stir under an atmosphere of H_2 gas for 4 h. The mixture was filtered through celite, the celite was washed with Et_2O (20 ml), and the filtrate was concentrated in vacuo. Flash chromatography (20% methanol/methylene chloride) afforded the amine (12 mg, 62%) as a pale yellow oil; $[\alpha]_{\text{D}}^{25} +6.0^\circ$ (c 0.57, CHCl_3); IR (CHCl_3) 3059 (w), 3020 (w), 3017 (m), 2845 (m), 2878 (m), 1455 (m), 1372 (br, m), 1209 (w), 1179 (s), 1122 (s), 1095 (s), 720 (br, m), 600 (m), 570 (m) cm^{-1} ; ^1H NMR (500 MHz, CD_3OD) δ 1.35 (m, 2 H), 1.42-1.54 (m, 4 H), 2.45 (m, 1 H), 2.59 (t, $J = 7.4$ Hz, 1 H), 2.90 (t, $J = 5.9$ Hz, 2 H), 3.04 (m, 1 H), 3.32 (dt, $J = 9.4, 6.4$ Hz, 1 H), 3.38-3.46 (m, 3 H), 3.58 (dd, $J = 10.8, 4.9$ Hz, 1 H), 3.69 (dd, $J = 11.0, 1.5$ Hz, 1 H), 3.76 (dt, $J = 9.7, 6.5$ Hz, 1 H), 4.11 (dt, $J = 9.7, 5.9$ Hz, 1 H), 4.25 (d, $J = 7.5$ Hz, 1 H), 4.36 (d, $J = 11.6$ Hz, 1 H), 4.44 (d, $J = 12.0$ Hz, 1 H), 4.48 (d, $J = 12.0$ Hz, 1 H), 4.52 (d, $J = 11.5$ Hz, 1 H), 7.11-7.26 (m, 12 H), 7.37 (m, 3 H), 7.47 (m, 3 H), 7.80 (m, 2 H), 7.88 (d, $J = 8.3$ Hz, 1 H); ^{13}C NMR (62.9 MHz, CD_3OD) δ 24.3, 26.4, 30.8, 32.0, 35.9, 41.9, 69.3, 70.4, 71.7, 72.3, 73.4, 74.4, 77.0, 79.1, 106.2, 114.7, 120.8, 122.0, 124.4, 125.3, 125.7, 127.9, 128.7, 128.8, 129.0, 129.4, 130.4, 132.6, 135.1, 136.6, 139.4, 139.6; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 713.3251 [$(\text{M} + \text{H})^+$; calcd for $\text{C}_{41}\text{H}_{49}\text{SO}_7 \text{N}_2$: 713.3260].

To a solution of the amine (0.011 g, 0.016 mmol) in MeOH (1.5 ml) was added 5M KOH (0.30 ml) and mixture was

- 77 -

heated at reflux for 6 h. After cooling, the mixture was diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography (20% methanol/methylene chloride) afforded I-47 (5.2 mg, 58%) as a pale yellow oil; $[\alpha]_D^{25} +3.8^\circ$ (c 0.16, CHCl_3); IR (CHCl_3) 3492 (m), 3018 (m), 2960 (m), 2872 (m), 1455 (m), 1370 (w), 1208 (s), 1090 (br, s), 720 (br, s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.35 (m, 2 H), 1.48 (m, 2 H), 1.80 (br s, 2 H), 2.54 (m, 1 H), 2.73 (t, $J = 4.0$ Hz, 2 H), 3.12 (t, $J = 6.7$ Hz, 2 H), 3.21 (m, 1 H), 3.50 (m, 3 H), 3.67 (m, 2 H), 3.81 (d, $J = 10.1$ Hz, 1 H), 3.85 (dt, 9.5, 7.3 Hz, 1 H), 4.28 (dt, $J = 9.3, 6.2$ Hz, 1 H), 4.38 (d, $J = 5.6$ Hz, 1 H), 4.42 (d, $J = 11.3$ Hz, 1 H), 4.61 (m, 3 H), 7.11 (t, $J = 7.1$ Hz, 1 H), 7.15 (br s, 1 H), 7.19 (t, $J = 7.1$ Hz, 1 H), 7.25-7.37 (m, 11 H), 7.62 (d, $J = 7.8$ Hz, 1 H), 8.87 (br s, 1 H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 23.3, 25.6, 30.0, 32.8, 34.9, 41.7, 69.3, 69.5, 70.8, 71.2, 72.4, 73.4, 75.8, 78.0, 105.0, 111.0, 112.9, 118.7, 119.0, 121.6, 122.4, 127.5, 127.6, 127.7, 128.1, 128.3, 128.4, 136.2, 138.1, 138.4; high resolution mass spectrum (Cl, NH_3) m/z 573.3301 [(M + H) $^+$]; calcd for $\text{C}_{35}\text{H}_{45}\text{O}_5$ N_2 : 573.3328].

AE. Alkene (I-49).

To a solution of amine I-54 (0.018 g, 0.026 mmol) and quinoline (6 ml) in benzene (1.5 ml) was added Lindlar's catalyst (6 mg, 30 wgt. %). The system was evacuated and back flushed with H_2 gas four times and then allowed to stir under an atmosphere of H_2 gas for 4 h. The mixture was filtered through celite, the celite was washed with Et_2O (20 ml), and the filtrate was concentrated in vacuo. The residue was used without purification in the next reaction.

To a solution of the crude amine in MeOH (1.5 ml) was added 5M KOH (0.30 ml) and mixture was heated at reflux for 6 h. After cooling, the mixture was diluted with water and extracted with methylene chloride (3 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography

- 78 -

(methylene chloride/toluene/methanol; 9:8:3) afforded I-49 (1.5 mg, 10% from I-54) as a pale yellow oil; ¹H NMR (500 MHz, CDCl₃) δ 1.49 (m, 1 H), 2.06 (m, 2 H), 2.49 (m, 2 H), 2.57 (m, 1 H), 3.09 (t, J = 5.8 Hz, 2 H), 3.28 (m, 1 H), 3.51 (m, 2 H), 3.67 (m, 1 H), 3.75 (d, J = 10.8 Hz, 1 H), 3.87 (dt, 9.3, 7.4 Hz, 1 H), 3.94-4.08 (m, 2 H), 4.25 (dt, J = 9.3, 7.0 Hz, 1 H), 4.40 (m, 2 H), 4.56 (m, 3 H), 5.35 (m, 1 H), 5.60 (m, 1 H), 7.08 (t, J = 7.0 Hz, 1 H), 7.08 (s, 1 H), 7.14 (t, J = 7.1 Hz, 1 H), 7.21-7.35 (m, 11 H), 7.59 (d, 7.7 Hz, 1 H), 8.90 (br s, 1 H); high resolution mass spectrum (Cl, NH₃) m/z 571.3182 [(M + H)⁺; calcd for C₃₅H₄₃O₅, N₂: 571.3171].

AF. Benzoylamide (+)-I-59.

Triflate I-62 was generated in the following way: A stirred solution of alcohol I-61 (0.20 g, 0.27 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.089 g, 0.44 mmol) in dry dichloromethane (3 ml) at -11°C was treated with triflic anhydride (0.060 ml, 0.35 mmol). After 10 min, the mixture was diluted with water (100 ml), saturated sodium bicarbonate (2 ml) and extracted with dichloromethane (2 x 200 ml). The combined extracts were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo, affording a yellow oil which was used without purification in the next reaction.

To a stirred solution of N-benzoyl-5-amino-1-pentanol (0.28 g, 1.36 mmol) in THF (6 ml) was added sodium hydride (60% dispersion in oil, 0.11 g, 2.80 mmol). The mixture was allowed to stir for 1.5 h, then cooled to 0°C before triflate I-62 was added via cannula (4 ml THF). After stirring an additional 18 h, the mixture was added to water (100 ml) and extracted with ether (3 x 20 ml). The combined extracts were washed with water, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography (40% ethyl acetate/hexanes) afforded the amide (90 mg, 36%) as a pale yellow oil which was used immediately in the next reaction.

To a solution of the amine (0.055 g, 0.060 mmol) in MeOH (3 ml) was added 5M KOH (0.30 ml) and the mixture was heated at reflux for 3 h. After cooling, the mixture was

- 79 -

diluted with water and extracted with methylene chloride (4 x 10 ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (40% ethyl acetate/hexanes) afforded I-59

5 (42 mg, 90%) as a clear yellow oil; $[\alpha]_D^{25} +12.3^\circ$ (c 0.31, CHCl_3); IR (CHCl_3) 3480 (m), 3097 (w), 3069 (w), 3035 (w), 3010 (m), 2960 (m), 2875 (m), 1660 (br, m), 1582 (w), 1520 (br, m), 1489 (m), 1455 (m), 1360 (br, m), 1305 (br, w), 1285 (br, w), 1070 (br, s), 695 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ

10 1.43 (m, 2 H), 1.59 (m, 4 H), 3.11 (t, $J = 7.0$ Hz, 2 H), 3.36 (m, 2 H), 3.44 (m, 4 H), 3.49-3.56 (m, 2 H), 3.60 (dd, $J = 10.8, 5.5$ Hz, 1 H), 3.64 (t, $J = 9.0$ Hz, 1 H), 3.69 (dd, $J = 10.8, 1.6$ Hz, 1 H), 3.85 (dt, $J = 9.3, 7.4$ Hz, 1 H), 4.22 (dt, $J = 9.4, 6.8$ Hz, 1 H), 4.45 (d, $J = 7.8$ Hz, 1 H), 4.60 (d, $J = 11.0$ Hz, 1 H), 4.65 (d, $J = 11.0$ Hz, 1 H), 4.76 (d, $J = 10.9$ Hz, 1 H), 4.85 (m, 2 H), 4.91 (d, $J = 10.9$ Hz, 1 H), 6.09 (br s, 1 H), 7.03 (m, 1 H), 7.09 (m, 1 H), 7.16 (m, 1 H), 7.21 (m, 2 H), 7.25-7.33 (m, 14 H), 7.40 (m, 2 H), 7.47 (m, 1 H), 7.59 (d, $J = 7.8$ Hz, 1 H), 7.73 (m, 2 H), 8.22 (br s, 1 H); ^{13}C NMR

20 (125.8 MHz, CDCl_3) δ 23.7, 25.7, 29.3, 29.4, 40.0, 69.7, 70.1, 71.5, 74.7, 74.8, 74.9, 75.7, 78.1, 82.3, 84.7, 103.7, 11.2, 112.5, 118.6, 119.2, 122.2, 126.8, 127.5, 127.5, 127.6, 127.8, 127.9, 128.0, 128.3, 128.3, 128.4, 128.5, 131.3, 134.7, 136.2, 138.2, 138.5, 138.6, 167.6; high resolution mass spectrum

25 (FAB, m-nitrobenzyl alcohol) m/z 782.3900 (M^+ ; calcd for $\text{C}_{49}\text{H}_{54}\text{O}_7$, N_2 : 782.3931).

AG. Trifluoroacetamide (+)-I-58.

To a stirred solution of N-trifluoroacetyl-5-amino-1-pentanol (0.27 g, 1.36 mmol) in THF (4 ml) was added sodium

30 hydride (60% dispersion in oil, 0.12 g, 3.00 mmol). The mixture was allowed to stir for 1.5 h, then cooled to 0°C before triflate I-62 was added via cannula (8 ml dichloromethane). After stirring an additional 18 h, the mixture was added to water (100 ml) and extracted with

35 dichloromethane (2 x 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography (40% ethyl

- 80 -

acetate/hexanes) afforded the amide (178 mg, 84%) as a pale yellow oil which was used immediately in the next reaction.

A solution of the amide (0.010 g, 0.011 mmol), 1,5-dimethoxynaphthalene (0.0062 g, 0.033 mmol) and NaCNBH₃ (0.0021 g, 0.011 mmol) in EtOH (4.8 ml) and water (0.16 ml) was purged with argon then irradiated with a Hanovia apparatus through pyrex for 4 h. The solvent was removed in vacuo and the remaining oil was diluted with water and extracted with dichloromethane (3 x 10 ml). The combined extracts were washed with water, dried over magnesium sulfate, and concentrated in vacuo. Preparative TLC (0.5 mm, 3% MeOH/CH₂Cl₂, 2x) afforded the amide (5 mg, 59%) as a pale yellow oil; $[\alpha]_D^{25} +17.6^\circ$ (c 0.46, CH₂Cl₂); IR (CHCl₃) 3490 (m), 3100 (w), 3075 (w), 3034 (w), 3014 (w), 2945 (m), 2880 (m), 1692 (s), 1610 (w), 1460 (m), 1362 (w), 1230 (w), 1200 (m), 1152 (s), 1090 (s), 1070 (s), 1040 (m), 910 (w), 697 (w) cm⁻¹; ¹H NMR (500 MHz, d₆-DMSO, 380 K) δ 1.26 (m, 2 H), 1.42 (m, 2 H), 1.56 (m, 2 H), 3.00 (t superimposed on a br s, $J = 7.2$ Hz, 2 H), 3.00 (br s, 2 H), 3.35-3.42 (m, 3 H), 3.45 (m, 2 H), 3.68 (t, $J = 8.8$ Hz, 2 H), 3.81 (dt superimposed on a br s, $J = 9.6, 7.2$ Hz, 1 H), 3.81 (br s, 1 H), 4.10 (dt, $J = 9.7, 6.9$ Hz, 1 H), 4.56 (br d, $J = 7.2$ Hz, 1 H), 4.61 (d, $J = 11.6$ Hz, 1 H), 4.63 (d, $J = 11.3$ Hz, 1 H), 4.71 (d, $J = 11.5$ Hz, 1 H), 4.77 (d, $J = 11.5$ Hz, 1 H), 4.79 (d, $J = 11.4$ Hz, 1 H), 4.83 (d, $J = 11.5$ Hz, 1 H), 6.96 (m, 1 H), 7.05 (m, 1 H), 7.08 (br s, 1 H), 7.20-7.34 (m, 16 H), 7.49 (d, $J = 7.7$ Hz, 1 H); ¹³C NMR (125.8 MHz, CDCl₃) δ 22.7, 22.9, 25.8, 26.2, 28.2, 32.1, 32.2, 47.9, 48.4, 48.5, 48.7, 62.5, 62.6, 70.0, 70.2, 72.5, 74.5, 74.7, 74.7, 74.8, 75.0, 75.7, 75.8, 78.9, 79.5, 82.1, 82.2, 84.3, 84.5, 103.5, 103.5, 111.1, 111.2, 115.2, 115.4, 117.5, 117.7, 118.6, 119.3, 119.3, 122.0, 122.0, 122.0, 122.1, 127.6, 127.6, 127.7, 127.7, 127.9, 127.9, 127.9, 128.0, 128.1, 128.1, 128.2, 128.3, 128.3, 128.4, 128.4, 128.4, 128.5, 136.2, 136.2, 137.5, 137.9, 138.2, 138.3, 138.3, 138.4, 156.4, 156.7, 157.0, 157.3; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 775.3543 [(M + H)⁺; calcd for C₄₄H₅₀O₇ N₂F₃: 782.3931).

- 81 -

EXAMPLE 10.**Preparation of Ester Compounds.**

In order to distinguish these compounds from those previously described, each compound number is preceded by "II". The chemical structures and the synthesis schemes for the compounds of Example 10 are presented in Figure 2.

A. 1,2,4,6-Tetra-O-acetyl- β -D-glucopyranoside (II-4).

A solution of 3-deoxy-diacetone-D-glucose II-3 in 60% aqueous acetic acid (200 ml) was heated at 90°C for 1 h, evaporated and azeotroped with dry benzene (4 x 20 ml). The residue was taken up in dry pyridine (250 ml), acetic anhydride (107 ml, 1.13 mol), DMAP (2 mol%, 275 mg) was added, and the solution was stirred at room temperature for 30 minutes. The mixture was evaporated, diluted with water (40 ml) and extracted with methylene chloride (3 x 40 ml). The combined extracts were washed with brine (40 ml), dried over sodium sulphate and evaporated. The residue was recrystallized from ether to afford the pure β -anomer as a fine white powder (11.3 g). The supernatant was evaporated and purified by flash chromatography eluting with 45% ethyl acetate in hexane to give a mixture of α - and β -anomers II-4 as a colorless gum (23.0g, total yield 91.7%). β -Anomer II-4 : m.p. 127-128° (ether) (lit. 129-130°)¹³; $[\alpha]_D^{25}$ -17.14° (c 1.05, CH₃OH); IR (CHCl₃) 3010 (m), 2940 (w), 2870 (w), 1745 (s), 1510 (w), 1365 (m), 1230 (s), 1210 (s), 1030 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 5.67 (d, J = 7.9 Hz, 1 H), 4.89-4.81 (m, 2 H), 4.21 (dd, J = 5.1, 12.3 Hz, 1 H), 4.12 (dd, J = 2.5, 12.2 Hz, 1 H), 3.81-3.79 (m, 1 H), 2.60 (ddd, J = 5.0, 5.0, 12.3 Hz, 1 H), 2.10 (s, 3 H), 2.06 (s, 3 H), 2.03 (s, 3 H), 2.02 (s, 3 H), 1.64 (dd, J = 11.0, 23.2 Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 170.69, 169.43, 169.31, 169.19, 93.06, 75.68, 67.33, 65.00, 62.07, 32.69, 20.92, 20.77; high resolution mass spectrum (CI) m/z 367.0773 [(M+Cl)⁺]; calcd for C₁₄H₂₀O₉Cl: 367.0796]. Anal. calcd for C₁₄H₂₀O₉: C, 50.60; H, 6.07; found: C, 50.65; H, 6.16.

- 82 -

B. 2-(1H-Indol-3-yl)ethyl 2,4,6-Tri-O-acetyl-3-deoxy- β -D-glucopyranoside (II-5).

Hydrogen bromide (30% in acetic acid) was added dropwise to a solution of the tetraacetate II-4 (9.97 g, 30.0mmol) in methylene chloride at 0°C. Stirring was continued at room temperature for 3 h, the mixture was poured into saturated aqueous sodium bicarbonate (500 ml) and extracted with ether (3 x 100 ml). The combined extracts were washed with saturated aqueous sodium bicarbonate (200 ml) and brine (200 ml), dried over sodium sulphate and evaporated. The pale yellow oil was azeotroped with benzene (4 x 20 ml) and dried under vacuum. A solution of the crude bromide in benzene (200 ml) was introduced into a flask containing activated powdered 4 Angstrom molecular sieves (10 g) and tryptophol (4.84 g, 30.0 mmol). Hexane (50 ml) and silver oxide (21 g, 90 mmol) were added, and the mixture was stirred vigorously in the dark for 18h. The solution was filtered through celite, evaporated, and purified by flash chromatography eluting with 10% ether in methylene chloride to afford the triacetate II-5 as a pale pinkish oil. (8.37 g, 64.4%): $[\alpha]_D^{25} + 22.04^\circ$ (c 1.08, CHCl_3); IR (CHCl_3) cm^{-1} 3020 (w), 2965 (w), 1745 (s), 1370 (m), 1230 (s), 1220 (s), 1205 (s), 1050 (s), 1035 (m), 740 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.06 (br s, 1 H), 7.57 (d, $J = 16.9$ Hz, 1 H), 7.32 (d, $J = 17.8$ Hz, 1 H), 7.16 (ddd, $J = 1.0, 8.8$ Hz, 1 H), 7.09 (ddm, $J = 8, 8$ Hz, 1 H), 7.02 (d, $J = 2.2, 1$ H), 4.84-4.77 (m, 1 H), 4.49 (d, $J = 7.5$ Hz, 1 H), 3.84-3.77 (m, 3 H), 3.77 (dd, $J = 7.3, 16.9$ Hz, 1 H), 3.69-3.65 (m, 1 H), 3.04 (t, $J = 7.1$ Hz, 2 H), 2.52 (ddd, $J = 5.1, 5.1, 12.3$ Hz, 1 H), 2.04 (s, 1 H), 2.02 (s, 1 H), 1.89 (s, 1 H), 1.57 (dd, $J = 9.0, 22$ Hz, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 170.83, 169.52, 136.06, 127.43, 122.26, 121.82, 119.19, 118.64, 112.41, 111.05, 102.08, 74.82, 69.80, 68.39, 65.83, 62.66, 32.71, 25.53, 20.83, 20.75; high resolution mass spectrum (CI) m/z 434.1782 [(M+H) $^+$]; calcd for $\text{C}_{22}\text{H}_{28}\text{O}_8$: 434.1815].

- 83 -

C. 2-(1H-Indol-3-yl)ethyl
3-Deoxy- β -D-glucopyranoside (II-6).

Sodium methoxide (9.42 mmol) was added in aliquots to a stirred solution of the triacetate II-5 (1.17 g, 2.69 mmol) in methanol (50 ml) at 0°C, and the solution stirred at room temperature for 15 h. Amberlyst® 15 ion-exchange resin was added to pH 7, and the mixture was filtered, evaporated and purified by flash chromatography eluting with 15% methanol in methylene chloride to afford the title compound II-6 as a colorless oil (752 mg, 90.9%): $[\alpha]_D^{25} +76.19^\circ$ (c1.05, CH₃OH); IR (CHCl₃) 3600-3200 (br), 2900 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.55 (d, J = 8.8 Hz, 1 H), 7.30 (d, J = 9 Hz, 1 H), 7.10-6.96 (m, 3 H), 4.27 (d, J = 7.6 Hz, 1 H), 4.18-4.13 (m, 1 H), 3.85-3.80 (m, 2 H), 3.64 (dd, J = 5.9, 11.8 Hz, 1 H), 3.53-3.48 (m, 1 H), 3.42-3.37 (m, 1 H), 3.34-3.29 (m, 1 H), 3.25-3.21 (m, 1 H), 3.09-3.03 (m, 2 H), 2.28 (ddd, J = 4.9, 4.9, 21.1 Hz, 1 H), 1.47 (dd, J = 11.5, 23.4 Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 137.98, 128.87, 123.63, 122.21, 119.5, 119.28, 112.55, 112.16, 106.49, 81.68, 71.11, 69.38, 66.14, 62.79, 40.56, 26.78; high resolution mass spectrum (CI) m/z [(M+H)⁺]; calcd for C₁₆H₂₁O₅N:].

D. 2-(1H-Indol-3-yl)ethyl 6-(p-Toluenesulphonyl)-
2,4-di-O-(tert-butyldimethyl)silyl-3-deoxy- β -D-
glucopyranoside (II-11).

p-Toluenesulphonyl chloride (0.559 mmol, 106 mg) and DMAP (10 mg) were added to a stirred solution of triol II-6 (56 mg, 0.508 mmol) and triethylamine (4.06 mmol, 0.76 ml) in methylene chloride (10 ml) at 0°C and the solution was stirred at room temperature for 30 min. More p-toluenesulphonyl chloride (5 mg) was added and the solution was stirred for a further 1 h, poured into saturated aqueous sodium bicarbonate (40 ml), extracted with methylene chloride (2 x 20 ml) and the combined extracts were washed with brine (20 ml), dried over sodium sulphate and evaporated. The resulting yellow oil was dissolved in methylene chloride (12 ml) and 2,6-lutidine (0.36 ml, 3.05 mmol) was added, followed by dropwise addition of tributyltrimethylsilyl triflate (2.03 mmol, 0.47 ml) at 0°C.

- 84 -

The solution was stirred at room temperature for 16 h, diluted with saturated aqueous sodium bicarbonate (25 ml), extracted with methylene chloride (3 x 20 ml) and the combined organic extracts were washed with brine (50 ml), dried over sodium sulphate and evaporated. The residue was purified by flash chromatography eluting with 30% ethyl acetate in hexane to give the title compound II-11 as a colorless oil (201 mg, 57.4%): $[\alpha]_D^{25} +2.25^\circ$ (c 0.71, CHCl_3); IR (CHCl_3) 2960 (s), 2950 (s), 2900 (s), 2860 (s), 1800 (w), 1605 (w), 1460 (s), 1365 (s), 1260 (s), 1100 (s), 980 (s) 920-890 (br), 840 (s), 695 (s), 550 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.01 (br s, 1 H), 7.72 (d, $J = 8.3$ Hz, 2 H), 7.56 (d, $J = 7.8$ Hz, 1 H), 7.34 (d, $J = 8.1$ Hz, 1 H), 7.19 - 7.09 (m, 5 H), 4.21 (dd, $J = 2.0$, 10.3 Hz, 1 H), 4.18 (d, $J = 7.6$ Hz, 1 H), 4.07 - 3.99 (m, 2 H), 3.79 - 3.74 (m, 1 H), 3.51 - 3.34 (m, 3 H), 3.06 - 3.03 (m, 2 H), 2.29 (s, 3 H), 2.14 - 2.10 (m, 1 H), 1.52 (app. q, $J = 11.4$ Hz, 1 H), 0.86 (s, 9 H), 0.81 (s, 9 H), 0.03 (s, 6 H), 0.00 (s, 6 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 144.61, 136.10, 132.75, 129.67, 127.87, 127.52, 122.14, 121.79, 119.16, 118.56, 112.40, 111.08, 105.09, 77.26, 69.88, 69.29, 69.00, 65.7, 41.50, 25.71, 25.58, 21.42, 18.11, 17.72.

E. 2-(1H-Indol-3-yl)ethyl 6-Iodo-2,4-di-O-(tert-butyldimethyl)silyl-3-deoxy- β -D-glucopyranoside (II-12).

A solution of tosylate II-11 (147 mg, 0.213 mmol) and sodium iodide (4.26 mmol, 639 mg) in dry acetone (8.0 ml) was heated to reflux for 16 h, diluted with saturated aqueous sodium thiosulphate (15 ml) and extracted with methylene chloride (3 x 15 ml). The combined extracts were washed with brine (10 ml), dried over sodium sulphate and evaporated. The residue was purified by flash chromatography eluting with 10% ethyl acetate in hexane to give the title compound II-12 (r_f 0.40) (91.3 mg, 66.3%) and starting material II-11 (r_f 0.20) (37.1 mg, 25.2%). 2-(1H-Indol-3-yl)ethyl 6-(p-Toluenesulphonyl)-2,4-di-O-(tert-butyldimethyl)silyl-3-deoxy- β -D-glucopyranoside II-12: $[\alpha]_D^{25} -4.63^\circ$ (c 0.67, CHCl_3); IR (CHCl_3) 3490 (w), 3010 (w), 2960 (m), 2930 (m),

- 85 -

2895 (w), 2860 (m), 1350 (w), 1090 (s), 835 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.93 (br s, 1 H), 7.61 (d, J = 6.8 Hz, 1 H), 7.33 (d, J = 8.1 Hz, 1 H), 7.18 - 7.15 (m, 1 H), 7.11 - 7.08 (m, 2 H), 4.28 (d, J = 7.4 Hz, 1 H), 4.19 (dt, J = 6.4, 9.3 Hz), 3.84 (dt, J = 6.4, 9.3 Hz, 1 H), 3.53 - 3.39 (m, 3 H), 3.18 - 3.07 (m, 3 H), 2.15 - 2.10 (m, 1 H), 1.57 (app. q, J = 11.3 Hz, 1 H), 0.87 (s, 9 H), 0.86 (s, 9 H), 0.07 (s, 6 H), 0.06 (s, 6 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 136.11, 127.56, 122.06, 121.87, 119.23, 118.82, 112.58, 111.03, 105.35, 78.82, 70.00, 69.92, 69.50, 41.50, 25.82, 25.76, 25.70, 18.18, 17.83, 6.78, -4.08, -4.43, -4.63, -4.90.

F. 2-(1H-Indol-3-yl)ethyl 6-(Trimethyl)acetyl-2,4-di-O-(tert-butyldimethyl)silyl-3-deoxy- β -D-glucopyranoside (II-13).

15 Pivaloyl chloride (18.6 mmol, 2.31 ml) was added dropwise at 0°C to a solution of the triol II-6 (5.18 g, 16.9 mmol) and DMAP (20 mg) in methylene chloride (150 ml) and triethylamine (135 mmol, 25.6 ml). The solution was stirred for 20 minutes at room temperature, an extra 0.32 ml (0.15 mmol) of pivaloyl chloride was added, and stirring was continued for 15 minutes. The solution was poured into ice-cold 1N HCl (200 ml), extracted with methylene chloride (3 x 50 ml) and the extracts were washed with saturated aqueous sodium bicarbonate (150 ml) and back-extracted with 20 methylene chloride (50 ml). The combined organic extracts were washed with brine (100 ml), dried (sodium sulphate) and evaporated. The resulting yellow oil was dissolved in methylene chloride and 2,6-lutidine (12.0 ml, 135 mmol) was added, followed by tributyltrimethylsilyl triflate (50.7 mmol, 30 11.6 ml) added dropwise at 0°C . The solution was stirred at room temperature for 15 h, diluted with saturated aqueous sodium bicarbonate (100 ml), extracted with methylene chloride (3 x 50 ml) and the combined organic extracts were washed with brine (100 ml), dried over sodium sulphate and evaporated. 35 The residue was purified by flash chromatography eluting with 15% ethyl acetate in hexane to give the title compound II-13 as a colorless oil (8.23 g, 78.7%): $[\alpha]_{\text{D}}^{25} -1.31^\circ$ (c 3.29,

- 86 -

CHCl₃); IR (CHCl₃) 3480 (m), 3020 (m), 2960 (s), 2920 (s), 2890 (m), 2860 (s), 1730 (s), 1470 (m), 1420 (m), 1390 (m), 1250 (s), 1230 (s), 1155 (s), 1080 (s), 1045 (s), 920 (m), 835 (s), 780-725 (s), 660 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.97 (br s, 1 H), 7.56 (d, J = 7.8, 1 H), 7.33 (d, J = 8.1 Hz, 1 H), 7.16 (dd, J = 7, 9 Hz, 1 H), 7.09 (dd, J = 7, 8 Hz, 1 H), 7.04 (dd, J = 1.1, 1.1 Hz, 1 H), 4.38 (d, J = 11.6, 1 H), 4.23 (d, J = 7.4 Hz, 1 H), 4.13-4.03 (m, 2 H), 3.79 (m, 1 H), 3.57-3.41 (m, 3 H), 3.09 (dd, J = 7.2, 7.3 Hz, 1 H), 2.14 (dt, J = 4.9, 12.4 Hz, 1 H), 1.56 (dd, J = 11.4, 23.8 Hz, 1 H), 1.22 (s, 9 H), 0.89 (s, 9 H), 0.86 (s, 9 H), 0.08 (s, 3 H), 0.08 (s, 3 H), 0.06 (s, 3 H), 0.05 (s, 3 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 178.33, 136.12, 127.52, 121.98, 118.64, 112.49, 111.02, 105.23, 77.73, 69.83, 69.22, 66.58, 63.65, 41.78, 38.76, 27.16, 27.01, 25.74, 25.65, 18.16, 17.83, -3.60, -4.16, -4.45, -4.95; high resolution mass spectrum (+ve FAB) m/z 619.3705 (M⁺; calcd for C₃₃H₃₇O₆NSi₂: 619.3724).

G. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-(tert-butyl-dimethyl)silyl-3-deoxy-β-D-glucopyranoside (II-14).

Sodium methoxide (6.0 mmol, 1.32 ml) was added in aliquots with stirring to pivaloate II-13 (740 mg, 1.20 mmol) in methanol (50 ml) and stirring was continued for 15 h. The solution was neutralised with Amberlyst® 15 ion-exchange resin, filtered and evaporated. The residue was purified by flash chromatography eluting with 30% ethyl acetate in hexane to afford the title compound II-14 as a colorless oil (468 mg, 73.2%): [α]_D²⁵ +15.67° (c 5.68, CHCl₃); IR (CHCl₃) 3480 (m), 3000 (s), 2960 (s), 2925 (s), 2880 (s), 2845 (s), 1710 (s), 1415 (m), 1360 (s), 1250 (s), 1220 (s), 1085 (s), 1030 (s), 1000 (m), 905 (m), 875 (s), 830 (s), 520 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.97 (br s, 1 H), 7.57 (d, J = 7.8 Hz, 1 H), 7.33 (d, J = 7.8 Hz, 1 H), 7.17 (ddd, J = 1.1, 7.7, 7.5), 7.10 (ddd, J = 0.9, 8, 8 Hz), 7.03 (s, 1 H), 4.28 (d, J = 7.4 Hz, 1 H), 4.10 (dd, J = 8.5, 16.3 Hz, 1 H), 3.83-3.77 (m, 2 H), 3.63-3.56 (m, 2 H), 3.50-3.45 (m, 1 H), 3.28-3.24 (m, 1 H), 3.09 (t, J = 7 Hz, 2 H), 2.26-2.22 (m, 1 H), 1.58 (dd, J =

- 87 -

11.3, 22.1 Hz, 1 H), 0.88 (s, 9 H), 0.84 (s, 9 H), 0.09 (s, 6 H), 0.08 (s, 6 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 136.14, 127.45, 121.96, 119.27, 118.64, 112.35, 111.09, 105.35, 79.65, 70.16, 69.39, 66.09, 62.42, 41.62, 25.75, 25.67, 18.17, 17.84, -4.28, -4.46, -4.87, -5.01; high resolution mass spectrum (CI) m/z 535.3172 [(M+H) $^+$]; calcd for $\text{C}_{28}\text{H}_{49}\text{O}_5\text{NSi}_2$: 535.3149].

H. 2-(1*H*-Indol-3-yl)ethyl 2,4-Di-O-(*tert*-butyl-dimethyl)silyl-3-deoxy-O-(6-azidohexyl)- β -D-glucopyranoside (II-15a).

10 Triflic anhydride (2.15 mmol, 0.36 ml) was added dropwise at -78°C to a solution of the alcohol II-14 (764 mg, 1.43 mmol) and 2,2-di-*tert*-butyl-4-methylpyridine (2.57 mmol, 528 mg) in methylene chloride (45 ml). The solution was stirred for 20 minutes, warmed to room temperature for 20
15 minutes, poured into saturated aqueous sodium bicarbonate (80 ml) and extracted with methylene chloride (2 x 40 ml). The extracts were combined, washed with brine (40 ml), dried over sodium sulphate, evaporated and dried under vacuum. Sodium hexamethyldisilylazide (0.6 M in toluene, 1.86 mmol, 3.10 ml)
20 was added dropwise to a solution of 6-azidohexanol (494 mg, 3.45 mmol) in methylene chloride (40 ml) at 0°C . The colorless triflate (purified by thin layer chromatography) was dissolved in methylene chloride and then added to the above solution at 0°C via cannula. Stirring was continued at room
25 temperature for 38 h, the solution diluted with saturated aqueous sodium bicarbonate (50 ml) and extracted with methylene chloride (3 x 25 ml). The combined organic extracts were washed with brine (40 ml), dried over sodium sulphate and evaporated. Purification by flash chromatography (eluting
30 with methylene chloride) furnished the title compound II-15a as a colorless viscous oil (257 mg, 27.3%): $[\alpha]_{\text{D}}^{25} +10.05^\circ$ (c 2.13, CHCl_3); IR (CHCl_3) 3480 (w), 3000 (w), 2950 (m), 2930 (m), 2855 (m), 2090 (m), 1360 (m), 1250 (w), 1080 (s), 830 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.00 (br s, 1 H), 7.58 (d, $J = 7.7$ Hz, 1 H), 7.33 (d, $J = 8.2$ Hz, 1 H), 7.16 (dd, $J = 7.1, 7.1$, 1 H), 7.09 (dd, $J = 7.0, 7.0$ Hz, 1 H), 7.02 (s, 1 H), 4.23 (d, $J = 6.4$ Hz, 1 H), 4.15-4.09 (m, 1 H), 3.80-3.75

- 88 -

(m, 1 H), 3.66 (dd, $J = 1.9, 10.8$ Hz, 1 H), 3.58-3.41 (m, 4 H), 3.35-3.31 (m, 1 H), 3.14 (t, $J = 7.0$ Hz, 2 H), 3.09 (t, $J = 7.5$ Hz, 2 H), 2.17-2.12 (m, 1 H), 1.60-1.45 (m, 5 H), 1.35-1.24 (m, 4 H), 0.87 (s, 9 H), 0.86 (s, 9 H), 0.08 (s, 3 H), 0.07 (s, 3 H), 0.06 (s, 3 H), 0.04 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 136.11, 127.54, 121.96, 121.84, 119.17, 117.70, 112.52, 111.03, 105.35, 79.65, 51.31, 41.83, 29.50, 28.67, 26.51, 25.77, 25.69, 18.19, 17.86, -4.19, -4.43, -4.90, -4.96; high resolution mass spectrum (+ve FAB) m/z 661.4213 [(M+H⁺); calcd for $\text{C}_{34}\text{H}_{61}\text{N}_4\text{O}_5\text{Si}_2$: 661.4180].

I. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-(tert-butyl-dimethyl)silyl-3-deoxy-O-(5-azidopentyl)- β -D-glucopyranoside (II-15b).

The same procedure as detailed above, using 5-azido-1-pentanol (2.4 eq., 4.08 mmol, 461 mg) furnished the title compound II-15b as a colorless oil (284 mg, 25.9%): $[\alpha]_{\text{D}}^{25} +7.31^\circ$ (c 1.67, CHCl_3); IR (CHCl_3) 3460 (m), 3000 (m), 2940 (s), 2920 (s), 2850 (s), 2080 (s), 1450 (w), 1250 (m), 1110 (s), 1080 (s), 830 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.00 (br s, 1 H), 7.57 (d, $J = 7.7$ Hz, 1 H), 7.33 (d, $J = 8.2$ Hz, 1 H), 7.17 (dt, $J = 8.1, 1, 1$ H), 7.09 (t, $J = 7.0$ Hz, 1 H), 7.03 (d, $J = 2.2$, 1 H), 4.23 (d, $J = 7.4$ Hz, 1 H), 4.11 (app. dd, $J = 8.5, 16.6$ Hz, 1 H), 3.77 (app. dd, $J = 8.6, 16.8$ Hz, 1 H), 3.58 - 3.43 (m, 5 H), 3.34-3.31 (m, 1 H), 3.12 - 3.07 (m, 4 H), 2.17 - 2.12 (m, 2 H), 1.58-1.48 (m, 5 H), 1.38-1.33 (m, 2 H), 0.88 (s, 9 H), 0.86 (s, 9 H), 0.08 (s, 3 H), 0.07 (s, 3 H), 0.05 (s, 3 H), 0.04 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 136.12, 127.50, 121.99, 119.11, 118.64, 112.39, 111.055, 105.35, 79.62, 71.33, 70.13, 69.99, 69.30, 66.10, 51.21, 41.80, 29.15, 28.58, 25.74, 25.67, 23.32, 18.16, 17.83, -4.20, -4.45, -4.92, -4.98; high resolution mass spectrum (+ve Cl) m/z 646.3887 (M⁺; calcd for $\text{C}_{33}\text{H}_{58}\text{N}_4\text{O}_5\text{Si}$: 646.3946).

J. 2-(1H-Indol-3-yl)ethyl 3-Deoxy-O-(6-azidohexyl)- β -D-glucopyranoside (II-16a).

Tetrabutylammoniumfluoride (1.0 M in THF (1.74 mmol, 1.74 ml) was added to a solution of the azide II-15a (230 mg,

- 89 -

0.348 mmol) in THF (10 ml) and stirred for 1 h. The solution was evaporated and the residue was purified by flash chromatography eluting with 10% methanol in methylene chloride to afford the title compound II-16a as a colorless oil (150 mg, 100%): $[\alpha]_D^{25} +38.24^\circ$ (c 1.53, CHCl_3); IR (CHCl_3) 3600 (w), 3470 (m), 3000 (w), 2930 (m), 2860 (m), 2090 (s), 1200 (m), 1080 (s), 1060 (s), 710 (s), 655 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.08 (br s, 1 H), 7.60 (d, $J = 7.8$ Hz, 1 H), 7.34 (d, $J = 8.1$ Hz, 1 H), 7.17 (dd, $J = 7, 7$ Hz, 1 H), 7.10 (dd, $J = 7, 7$ Hz, 1 H), 7.03 (s, 1 H), 4.22-4.16 (m, 2 H), 3.85-3.38 (m, 10 H), 3.25-2.99 (m, 4 H), 2.35-2.27 (m, 1 H), 2.14 (br s, 1 H), 1.72-1.24 (m, 9 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 136.2, 122.12, 121.90, 119.43, 118.70, 111.25, 105.14, 76.06, 72.78, 71.85, 70.42, 69.17, 68.29, 51.32, 37.28, 29.33, 28.72, 26.46, 25.74, 25.61; high resolution mass spectrum (+ve FAB) m/z 432.2411 (M^+ ; calcd for $\text{C}_{22}\text{H}_{32}\text{N}_4\text{O}_5$: 432.2373).

K. 2-(1H-Indol-3-yl)ethyl 3-Deoxy-O-(5-azidopentyl)- β -D-glucopyranoside (II-16b).

The same procedure as detailed above afforded the title compound II-16b as a colorless oil (173 mg, 100%): $[\alpha]_D^{25} +31.01^\circ$ (c 0.79, CHCl_3); IR (CHCl_3) 3480 (m), 3005 (w), 2950 (m), 2880 (m), 2100 (s), 1455 (w), 1280 (w), 1090 (w), 1070 (s), 1060 (s), 1020 (w), 1010 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.02 (br s, 1 H), 7.61 (d, $J = 7$ Hz, 1 H), 7.33 (dd, $J = 0.7, 8.0$ Hz, 1 H), 7.18 (app. t, $J = 8$ Hz, 1 H), 7.11 (app. t, $J = 8$ Hz, 1 H), 7.03 (d, $J = 2.3$, 1 H), 4.22 - 4.18 (m, 2 H), 3.75-3.65 (m, 3 H), 3.61 (dd, $J = 7.2, 9.6$ Hz, 1 H), 3.52-3.38 (m, 4 H), 3.23 (t, $J = 6.9$ Hz, 2 H); 3.13 - 3.00 (m, 4 H), 2.35-2.31 (m, 1 H), 2.11 (br s, 1 H), 1.75-1.68 (m, 1 H), 1.61 - 1.35 (m, 7 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 1136.10, 127.45, 122.03, 121.84, 119.16, 118.52, 112.43, 111.23, 104.95, 76.53, 72.15, 71.54, 70.16, 68.27, 68.15, 52.83, 51.15, 37.37, 28.89, 28.45, 25.60, 25.19, 23.15, 19.74, 13.39; high resolution mass spectrum (+ve CI) m/z 436.2537 [$(\text{M}+\text{NH}_4)^+$]; calcd for $\text{C}_{21}\text{H}_{34}\text{N}_5\text{O}_5$: 436.2560].

- 90 -

L. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-(2,2-dimethyl-3-phenylpropanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-glucopyranoside (II-17a) and 2-(1H-Indol-3-yl)ethyl 2-O-(2,2-Dimethyl-3-phenylpropanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-glucopyranoside (II-18a).

2,2-Dimethyl-4-aminopyridine (40 mol%, 11 mg) was added to a vigorously stirred solution of diol II-16a (91.4 mg, 0.212 mmol), 2,2-dimethyl-3-phenylpropanoic acid (242 mg, 1.27 mmol) and dicyclohexylcarbodiimide (703 mg, 3.39 mmol) in chloroform (5 ml) and the mixture was refluxed for 40 h. The cooled solution was evaporated, taken up in ether, filtered and evaporated again. The residue was purified by flash chromatography eluting with 25% ethyl acetate in hexane to afford an impure component (RF 0.28). The gradient was increased to 40% ethyl acetate in hexane affording an impure component (RF 0.23). The higher-running compound was recolumned in 50% methylene chloride in hexane increasing to 10% ether in methylene chloride to afford the pure bis-ester II-17a as a colorless oil (97.2 mg, 61.1%). The lower-running compound was recolumned in 10% ether in methylene chloride to give the pure mono-ester II-18a as a colorless oil (39.5 mg, 31.3%).

2-(1H-Indol-3-yl)ethyl 2,4-Di-O-(2,2-dimethyl-3-phenylpropanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-glucopyranoside (II-17a) (bis-ester): $[\alpha]_D^{25} +36.18^\circ$ (c 0.34, CHCl₃); IR (CHCl₃) 3480 (w), 3020 (w), 2935 (m), 2860 (m), 1730 (s), 1725 (s), 1455 (w), 1120 (s), 1005 (w), 690 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.84 (br s, 1 H), 7.53 (d, J = 7.8 Hz, 1 H), 7.32 (d, J = 8.0 Hz, 1 H), 7.26-7.03 (m, 12 H), 6.99 (s, 1 H), 4.83-4.75 (m, 2 H), 4.54 (d, J = 7.8 Hz, 1 H), 4.09 (ddm, J = 8, 17 Hz, 1 H), 3.79 (ddm, J = 8, 16 Hz, 1 H), 3.67-3.64 (m, 1 H), 3.54 (d, J = 11.2 Hz, 1 H), 3.48-3.35 (m, 3 H), 3.18 (br s, 2 H), 3.05-3.02 (m, 2 H), 2.88 (d, J = 13.4 Hz, 1 H), 2.74 (d, J = 13.4 Hz, 1 H), 2.53-2.47 (m, 2 H), 1.80 (t, J = 8.4 Hz, 2 H), 1.52-1.14 (m, 9 H), 1.21 (s, 6 H), 1.14 (s, 3 H), 1.08 (s, 3 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 176.13, 175.99,

- 91 -

141.93, 137.67, 136.25, 130.23, 128.43, 128.22, 127.95, 127.46, 126.43, 125.93, 122.29, 121.70, 119.02, 118.57, 111.78, 111.21, 102.34, 77.20, 71.75, 70.00, 69.89, 68.73, 66.33, 45.94, 43.48, 42.43, 42.36, 33.88, 33.35, 31.42, 29.68, 29.59, 29.33, 26.52, 25.70, 25.36, 25.18, 24.92, 24.31; high resolution mass spectrum (+ve FAB) m/z 753.4263 [(M+H)⁺]; calcd for C₄₄H₅₇N₄O₇: 753.4227].

2-(1H-Indol-3-yl)ethyl 2-O-(2,2-Dimethyl-3-phenylpropanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-glucopyranoside (II-18a) (mono-ester): [α]_D²⁵ +31.15° (c 1.11, CHCl₃); IR (CHCl₃) 3680 (w), 3620 (w), 3480 (s), 3020 (s), 2975 (m), 2935 (m), 2875 (m), 2090 (m), 1725 (m), 1520 (m), 1470 (m), 1420 (m), 1220 (s), 1070 (m), 925 (m), 760 (s), 660 (s), 615 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.87 (br s, 1 H), 7.53 (d, J = 8.0 Hz, 1 H), 7.31 (d, J = 8.1 Hz, 1 H), 7.23-7.06 (m, 7 H), 6.98 (s, 1 H), 4.77-4.72 (m, 1 H), 4.50 (d, J = 7.7 Hz, 1 H), 4.07 (dd, J = 8.2, 16.0 Hz, 1 H), 3.78-3.70 (m, 2 H), 3.61 (app. t, J = 7.6 Hz, 1 H), 3.51-3.43 (m, 2 H), 3.23 (t, J = 6.9 Hz, 1 H), 3.01 (dt, J = 3, 7 Hz, 2 H), 2.84 (d, J = 13.3, 1 H), 2.76 (d, J = 13.3 Hz, 1 H), 2.42 (app. dt, J = 12.2, 5.0 Hz, 1 H), 1.58-1.47 (m, 5 H), 1.36-1.34 (m, 4 H), 1.12 (s, 3 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 176.15, 137.73, 136.07, 130.22, 127.96, 127.41, 126.43, 122.11, 121.88, 119.22, 118.60, 112.29, 111.06, 102.26, 76.02, 72.70, 69.71, 69.18, 68.71, 51.30, 45.87, 43.47, 35.80, 29.31, 28.69, 26.44, 25.68, 25.59, 25.13, 24.49; high resolution mass spectrum (+ve FAB) m/z 592.3228 [(M+H)⁺]; calcd for C₃₃H₄₄N₄O₆: 592.3261].

M. 2-(1H-Indol-3-yl)ethyl 2-O-(2,2-Dimethyl-3-phenylpropanoyl)-4-O-(2,2-dimethyl-3-phenylbutanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-glucopyranoside (II-19a).

2,2-Dimethyl-4-aminopyridine (2 mg) was added to a vigorously stirred solution of mono-ester II-18a (25.6 mg, 0.0430 mmol), 2,2-dimethyl-4-phenylbutanoic acid (49.5 mg, 0.258 mmol) and dicyclohexylcarbodiimide (88.7 mg, 0.430 mmol) in methylene chloride (2 ml) and the mixture refluxed for 18 h. The cooled solution was evaporated, taken up in ether,

- 92 -

filtered and evaporated. The residue was purified by flash chromatography eluting with 20% ethyl acetate in hexane to furnish the title compound II-19a as a colorless oil (29.0 mg, 87.7%): $[\alpha]_D^{25} +8.33^\circ$ (c 0.60, CHCl_3); IR (CHCl_3) 3480 (w), 3020 (w), 2940 (m), 2860 (m), 2100 (m), 1735 (s), 1455 (m), 1120 (s), 895 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.94 (br s, 1 H), 7.54 (d, $J = 7.9$ Hz, 1 H), 7.31 (d, $J = 8.1$ Hz, 1 H), 7.28-7.07 (m, 12 H), 6.99 (d, $J = 1.2$ Hz, 1 H), 4.86-4.78 (m, 2 H), 4.55 (d, $J = 7.8$ Hz, 1 H), 4.13 (ddm, $J = 8.5, 15.9$ Hz, 1 H), 3.80 (ddm, $J = 8.5, 16.4$ Hz, 1 H), 3.69-3.65 (m, 1 H), 3.56 (dd, $J = 2.1, 11.0$ Hz, 1 H), 3.51-3.36 (m, 3 H), 3.15 (t, $J = 7.0$ Hz, 2 H), 3.08-3.01 (m, 1 H), 2.87 (d, $J = 13.3$ Hz, 1 H), 2.74 (d, $J = 13.3$ Hz, 1 H), 2.56-2.47 (m, 3 H), 1.82 (t, $J = 8.8$ Hz, 2 H), 1.54-1.45 (m, 4 H), 1.36-1.25 (m, 6 H), 1.23 (s, 6 H), 1.13 (s, 3 H), 1.07 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 176.07, 175.90, 141.93, 137.69, 136.07, 130.23, 128.41, 128.20, 127.93, 127.43, 126.41, 125.92, 122.17, 121.87, 119.20, 118.61, 112.29, 111.06, 102.34, 77.17, 71.75, 70.03, 69.73, 68.77, 66.28, 51.30, 45.90, 43.44, 42.44, 42.33, 33.22, 31.41, 29.44, 28.68, 26.47, 25.63, 25.27, 25.16, 24.95, 24.28; high resolution mass spectrum (+ve FAB) m/z 767.4361 [(M+H) $^+$]; calcd for $\text{C}_{45}\text{H}_{59}\text{N}_4\text{O}_7$: 767.4384].

N. 2-(1H-Indol-3-yl)ethyl 2-O-(2,2-Dimethyl-3-phenylpropanoyl)-4-O-(2,2-dimethyl-3-phenylbutanoyl)-3-deoxy-O-(6-aminohexyl)- β -D-glucopyranoside (II-1a).

A solution of bis-ester II-19a (11.7 mg, 0.0152 mmol) and triphenylphosphine (9.97 mg, 0.0380 mmol) in THF (0.8 ml) and water (12 ml) was heated at 55°C for 15 h. The cooled solution was evaporated and purified by flash chromatography eluting with methanol/methylene chloride/acetic acid (10:90:1) increasing the gradient to (30:70:1). Fractions containing the title compound were treated with solid sodium bicarbonate, filtered, evaporated, redissolved in methylene chloride, filtered and evaporated, to afford the title compound II-1a as a colorless oil (10.6 mg, 93.8%): $[\alpha]_D^{25} +36.18^\circ$ (c 0.34, CHCl_3); IR (CHCl_3) 3480 (w), 3020 (w), 2935 (m), 2860 (m),

- 93 -

1730 (s), 1725 (s), 1455 (w), 1120 (s), 1005 (w), 690 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.84 (br s, 1 H), 7.53 (d, $J = 7.8$ Hz, 1 H), 7.32 (d, $J = 8.0$ Hz, 1 H), 7.26-7.03 (m, 12 H), 6.99 (s, 1 H), 4.83-4.75 (m, 2 H), 4.54 (d, $J = 7.8$ Hz, 1 H), 5 4.09 (ddm, $J = 8, 17$ Hz, 1 H), 3.79 (ddm, $J = 8, 16$ Hz, 1 H), 3.67-3.64 (m, 1 H), 3.54 (d, $J = 11.2$ Hz, 1 H), 3.48-3.35 (m, 3 H), 3.18 (br s, 2 H), 3.05-3.02 (m, 2 H), 2.88 (d, $J = 13.4$ Hz, 1 H), 2.74 (d, $J = 13.4$ Hz, 1 H), 2.53-2.47 (m, 2 H), 1.80 (t, $J = 8.4$ Hz, 2 H), 1.52-1.14 (m, 9 H), 1.21 (s, 6 H), 1.14 10 (s, 3 H), 1.08 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 176.13, 175.99, 141.93, 127.67, 136.25, 130.23, 128.43, 128.22, 127.95, 127.46, 126.43, 125.93, 122.29, 121.70, 119.02, 118.57, 111.78, 111.21, 102.34, 77.20, 71.75, 70.00, 69.89, 68.73, 66.33, 45.94, 43.48, 42.43, 42.36, 33.88, 33.35, 31.42, 15 29.68, 29.59, 29.33, 26.52, 25.70, 25.36, 25.18, 24.92, 24.31; high resolution mass spectrum (+ve FAB) m/z 741.4430 [(M+H) $^+$]; calcd for $\text{C}_{45}\text{H}_{61}\text{N}_2\text{O}_7$: 741.4478].

O. 2-(1*H*-Indol-3-yl)ethyl 2,4-Di-O-(2,2-dimethyl-3-phenylpropanoyl)-3-deoxy-O-(6-aminohexyl)- β -D-glucopyranoside (II-1c).

The same procedure as detailed above afforded the title compound II-1c as a colorless oil (27.5 mg, 81.6%): $[\alpha]_D^{25} +2.86^\circ$ (c 0.28, CHCl_3); IR (CHCl_3) 3680 (w), 3480 (w), 3025 (w), 3005 (w), 2965 (w), 2930 (m), 2860 (w), 1730 (s), 1600 25 (w), 1450 (w), 1115 (s), 895 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.55 (br s, 1 H), 7.51 (d, $J = 7.9$ Hz, 1 H), 7.32 (d, $J = 9.4$ Hz, 1 H), 7.28-7.03 (m, 12 H), 6.99 (s, 1 H), 4.81-4.76 (m, 1 H), 4.74-4.69 (m, 1 H), 4.51 (d, $J = 7.8$ Hz, 1 H), 4.09 (ddm, $J = 8, 14$ Hz, 1 H), 3.78 (ddm, $J = 8, 17$ Hz, 1 H), 30 3.62-3.58 (m, 1 H), 3.38-3.33 (m, 3 H), 3.06-3.00 (m, 2 H), 2.88 (d, $J = 13.3$ Hz, 1 H), 2.81 (s, 2 H), 2.75 (d, $J = 13.3$ Hz, 1 H), 2.47-2.42 (m, 1 H), 2.3 (br s, 2 H), 1.49-1.16 (m, 9 H), 1.14 (s, 6 H), 1.14 (s, 3 H), 1.08 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 176.04, 175.92, 137.69, 137.43, 136.14, 35 130.22, 127.98, 127.43, 126.53, 126.41, 122.23, 121.73, 119.07, 118.55, 112.01, 111.14, 102.24, 77.12, 71.74, 69.91, 69.73, 68.74, 66.56, 45.98, 45.88, 43.45, 33.18, 29.56, 26.55,

- 94 -

25.85, 25.60, 25.28, 24.83, 24.31; high resolution mass spectrum (+ve FAB) m/z 727.4341 ($M+H^+$; calcd for $C_{44}H_{59}N_2O_7$: 727.4322).

5 P. 2-(1*H*-Indol-3-yl)ethyl 2-O-(2,2-dimethyl-3-phenyl-propanoyl)-3-deoxy-O-(5-azidopentyl)- β -D-gluco-pyranoside (II-14b).

2,2-Dimethyl-4-aminopyridine (40 mol%, 18 mg) was added to a vigorously stirred solution of diol II-16b (146 mg, 0.349 mmol), 2,2-dimethyl-3-phenylpropanoic acid (333 mg, 1.75 mmol) and dicyclohexylcarbodiimide (1.09 g, 5.24 mmol) in chloroform (10 ml) and the mixture refluxed for 18 h. The cooled solution was evaporated, taken up in ether, filtered and evaporated. The residue was purified by flash chromatography eluting with 40% ethyl acetate in hexane to afford the somewhat impure mono-ester (RF 0.20). The eluant was changed to 10% methanol/dichloromethane to afford recovered starting material (61.2 mg, 41.9%). The mono-ester was further purified by flash chromatography eluting with 10% ether/dichloromethane to give the title compound II-18b as a colorless oil (65.3 mg, 32.1%): $[\alpha]_D^{25} +36.61^\circ$ (c 1.21, $CHCl_3$); IR ($CHCl_3$) 3500 (m), 3010 (w), 2940 (m), 2880 (m), 2100 (s), 1460 (m), 1120 (s), 1070 (s) cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 7.92 (br s, 1 H), 7.54 (d, $J = 7.9$ Hz, 1 H), 7.32 (d, $J = 8.1$ Hz, 1 H), 7.26-7.07 (m, 7 H), 7.08 (d, $J = 7$, 1 H), 4.79-4.74 (m, 1 H), 4.52 (d, $J = 7.6$ Hz, 1 H), 4.10 (m, 1 H), 3.80-3.71 (m, 3 H), 3.62 (dd, $J = 7.3, 9.6$ Hz, 1 H), 3.54-3.45 (m, 3 H), 3.24 (t, $J = 6.9$ Hz, 2 H), 3.12 (br s, 1 H), 3.06 - 3.00 (m, 2 H), 2.86 (d, $J = 13.4$ Hz, 1 H), 2.77 (d, $J = 13.4$ Hz, 1 H), 2.46 - 2.41 (m, 1 H), 1.63 - 1.38 (m, 7 H), 1.14 (s, 3 H), 1.09 (s, 3 H); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 176.16, 137.67, 136.05, 130.17, 127.93, 127.37, 126.40, 122.12, 121.80, 119.13, 118.53, 112.12, 111.06, 102.20, 76.27, 72.43, 71.64, 69.70, 69.18, 68.30, 51.18, 45.83, 43.44, 35.82, 28.95, 28.51, 25.65, 25.10, 24.45, 23.20; high resolution mass spectrum (+ve Cl) m/z 578.3107 (M^+ ; calcd for $C_{32}H_{42}N_4O_6$: 578.3104).

- 95 -

Q. 2-(1*H*-Indol-3-yl)ethyl 2-O-(2,2-Dimethyl-3-phenyl-propanoyl)-4-O-(2,2-dimethyl-3-phenylbutanoyl)-3-deoxy-0-(5-azidopentyl)- β -D-glucopyranoside (II-19b).

5 2,2-Dimethyl-4-aminopyridine (2 mg) was added to a vigorously stirred solution of mono-ester II-18b (34.8 mg, 0.0598 mmol), 2,2-dimethyl-4-phenylbutanoic acid (68.9 mg, 0.359 mmol) and dicyclohexylcarbodiimide (123 mg, 0.598 mmol) in methylene chloride (2.5 ml) and the mixture was refluxed
10 for 20 h. The cooled solution was evaporated, taken up in ether, filtered and evaporated. The residue was purified by flash chromatography eluting with 20% ethyl acetate in hexane to furnish the title compound II-19b as a colorless oil (39.4 mg, 87.2%): $[\alpha]_D^{25} +6.03^\circ$ (c 0.58, CHCl₃); IR (CHCl₃) 33490
15 (w), 2940 (m), 2930 (m), 2100 (m), 1735 (s), 1730 (s), 1140 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.89 (br s, 1 H), 7.53 (d, J = 7.9 Hz, 1 H), 7.31 (dd, J = 0.5, 8.1 Hz, 1 H), 7.27-7.06 (m, 12 H), 7.00 (d, J = 2.0 Hz, 1 H), 4.84-4.78 (m, 2 H), 4.53 (d, J = 7.8 Hz, 1 H), 4.12 (dt, J = 6.7, 8.6 Hz, 1 H),
20 3.80 - 3.76 (m, 1 H), 3.54 (dd, J = 2.0, 11.0 Hz), 3.49 - 3.35 (m, 3 H), 3.12 (t, J = 6.9 Hz, 2 H), 3.07-3.00 (m, 1 H), 2.85 (d, J = 13.3 Hz, 1 H), 2.73 (d, J = 13.3 Hz, 1 H), 2.55 - 2.48 (m, 2 H), 1.81 (t, J = 8.8 Hz, 2 H), 1.54-1.44 (m, 7 H), 1.36 -1.29 (m, 2 H), 1.22 (s, 6 H), 1.12 (s, 3 H), 1.06 (s, 3 H);
25 ¹³C NMR (62.9 MHz, CDCl₃) δ 176.06, 175.89, 141.90, 137.66, 136.06, 130.21, 128.40, 128.17, 127.90, 127.41, 126.39, 125.90, 122.16, 121.85, 119.17, 118.57, 112.25, 111.06, 102.32, 77.12, 71.56, 70.03, 69.71, 68.74, 66.24, 51.21, 45.89, 43.42, 42.39, 42.33, 33.29, 31.37, 29.07, 28.54, 25.63,
30 25.25, 25.12, 24.92, 24.27, 23.27; high resolution mass spectrum (+ve CI) m/z 753.4261 [(M+H)⁺]; calcd for C₄₄H₅₇N₄O₇: 753.4228].

- 96 -

R. 2 - (1 H - I n d o l - 3 - y l) e t h y l
2 - O - (2 , 2 - D i m e t h y l - 3 - p h e n y l -
propanoyl)-4-O-(2,2-dimethyl-3-phenylbutanoyl)-3-
deoxy-O-(5-aminopentyl)- β -D-glucopyranoside (II-
1b).

A solution of bis-ester II-19b (26.7 mg, 0.0353 mmol) and triphenylphosphine (23.2 mg, 0.0833 mmol) in THF (1.5 ml) and water (20 ml) was heated at 55°C for 15 h. The cooled solution was evaporated and purified by flash chromatography eluting with methanol/methylene chloride/acetic acid (10:90:1) increasing the gradient to (30:70:1). Fractions containing the title compound were treated with solid sodium bicarbonate, filtered, evaporated, redissolved in methylene chloride, refiltered and evaporated to afford the title compound II-1b as a colorless oil (18.7 mg, 72.5%): $[\alpha]_D^{25} +25.00^\circ$ (c 0.32, CHCl_3); IR (CHCl_3) 3480 (w), 3010 (s), 2920 (m), 2860 (w), 2390 (m), 1730 (m), 1520 (m), 1470 (m), 1420 (m), 1210 (s), 1120 (m), 920 (m), 840 (m), 750 (s), 660 (s), 615 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 9.13 (br s, 1 H), 7.52 (d, $J = 7.9$ Hz, 1 H), 7.32-7.11 (m, 12 H), 7.05 (t, $J = 5.7$ Hz, 1 H), 6.98 (s, 1 H), 4.83-4.75 (m, 2 H), 4.68 (br s, 2 H), 4.53 (d, $J = 7.8$ Hz, 1 H), 4.10 - 4.05 (m, 1 H), 3.81 - 3.76 (m, 1 H), 3.66 - 3.63 (m, 1 H), 3.51 (dd, $J = 1.8, 11.1$ Hz, 1 H), 3.45 - 3.31 (m, 3 H), 3.06 - 3.00 (m, 2 H), 2.89 (d, $J = 13.3$ Hz, 1 H), 2.74 (d, $J = 13.3$ Hz, 1 H), 2.55 - 2.45 (m, 4 H), 1.80 (t, $J = 8.8$ Hz, 2 H), 1.52-1.34 (m, 5 H), 1.30 - 1.21 (m, 3 H), 1.21 (s, 6 H), 1.15 (s, 3 H), 1.09 (s, 3 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 176.17, 175.99, 141.90, 137.67, 136.24, 130.23, 128.43, 128.21, 127.96, 126.45, 125.95, 122.31, 121.70, 119.01, 118.55, 11.82, 111.26, 102.28, 71.48, 69.83, 68.72, 66.27, 45.95, 43.51, 42.36, 33.32, 31.42, 29.70, 29.22, 25.61, 25.39, 25.17, 24.95, 24.32, 23.28; high resolution mass spectrum (-ve CI) m/z 761.3902 [$(\text{M}+\text{Cl})^-$]; calcd for $\text{C}_{44}\text{H}_{58}\text{N}_2\text{O}_7\text{Cl}$: 761.3932].

- 97 -

S. 2-(1*H*-Indol-3-yl)ethyl 2-O-(3-phenylpropanoyl)-3-deoxy-O-(6-azidohexyl)- β -D-glucopyranoside (II-23).

A solution of diol II-16a (143 mg, 0.331 mmol) in methylene chloride (10 ml) was added dropwise to a stirred solution of hydrocinnamic acid (0.331 mmol, 49.7 mg), dicyclohexylcarbodiimide (0.331 mmol, 68.2 mg) and 2,2'-dimethyl-4-aminopyridine (1 mg) in methylene chloride at 0°C. The solution was warmed to room temperature and stirred for 16 h, evaporated, taken up in ether, filtered and evaporated. The residue was purified by flash chromatography eluting with 45% ethyl acetate/hexane to give a higher component (C-4 monoester II-24) (Rf 0.25), mixed fractions and a lower component (C-2 monoester II-23) (Rf 0.23). The mixed fractions were combined and the process was repeated twice. This produced a pure sample of the lower, C-2 monoester II-23 as a colorless oil (35.0 mg, 18.7%): $[\alpha]_D^{25} +25.00^\circ$ (c 0.32, CHCl₃); IR (CHCl₃) 3480 (w), 3010 (s), 2920 (m), 2860 (w), 2390 (m), 1730 (m), 1520 (m), 1470 (m), 1420 (m), 1210 (s), 1120 (m), 920 (m), 840 (m), 750 (s), 660 (s), 615 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 9.13 (br s, 1 H), 7.52 (d, *J* = 7.9 Hz, 1 H), 7.32 - 7.11 (m, 12 H), 7.05 (t, *J* = 5.7 Hz, 1 H), 6.98 (s, 1 H), 4.83 - 4.75 (m, 2 H), 4.68 (br s, 2 H), 4.53 (d, *J* = 7.8 Hz, 1 H), 4.10 - 4.05 (m, 1 H), 3.81 - 3.76 (m, 1 H), 3.66 - 3.63 (m, 1 H), 3.51 (dd, *J* = 1.8, 11.1 Hz, 1 H), 3.45 - 3.31 (m, 3 H), 3.06 - 3.00 (m, 2 H), 2.89 (d, *J* = 13.3 Hz, 1 H), 2.74 (d, *J* = 13.3 Hz, 1 H), 2.55 - 2.45 (m, 4 H), 1.80 (t, *J* = 8.8 Hz, 2 H), 1.52-1.34 (m, 5 H), 1.30 - 1.21 (m, 3 H), 1.21 (s, 6 H), 1.15 (s, 3 H), 1.09 (s, 3 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 176.17, 175.99, 141.90, 137.67, 136.24, 130.23, 128.43, 128.21, 127.96, 126.45, 125.95, 122.31, 121.70, 119.01, 118.55, 11.82, 111.26, 102.28, 71.48, 69.83, 68.72, 66.27, 45.95, 43.51, 42.36, 33.32, 31.42, 29.70, 29.22, 25.61, 25.39, 25.17, 24.95, 24.32, 23.28; high resolution mass spectrum (-ve CI) *m/z* 761.3902 [(M+Cl)⁻]; calcd for C₄₄H₅₈N₂O₇Cl: 761.3932].

- 98 -

T. 2 - (1 H - I n d o l - 3 - y l) e t h y l
2-O-(3-phenylpropanoyl)-4-O-
(4-phenylbutanoyl)-3-deoxy-O-(6-azidoheptyl)- β -D-
glucopyranoside (II-25).

5 2,2-Dimethyl-4-aminopyridine (1 mg) was added to a
vigorously stirred solution of mono-ester II-23 (13.8 mg,
0.0245 mmol), 4-phenylbutyric acid (8.0 mg, 0.0490 mmol) and
dicyclohexylcarbodiimide (20.2 mg, 0.0980 mmol) in methylene
chloride (1.5 ml) and the mixture was stirred at room
10 temperature for 20 h, evaporated, taken up in ether,
refiltered and evaporated. The residue was purified by flash
chromatography eluting with 25% ethyl acetate/hexane to
furnish the title compound II-25 as a colorless oil (17.0 mg,
97.9%): $[\alpha]_D^{25} +9.15^\circ$ (c 0.59, CHCl₃); IR (CHCl₃) 3490 (m),
15 3020 (w), 2950 (m), 2870 (m), 2100 (s), 1745 (s), 1460 (m),
1160 (m), 1135 (m), 1080 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ
7.86 (br s, 1 H), 7.56 (d, J = 7.3 Hz, 1 H), 7.33 - 7.07 (m,
13 H), 6.99 (d, J = 2.3 Hz, 1 H), 4.82 - 4.77 (m, 2 H), 4.46
(d, J = 7.5 Hz, 1 H), 4.14 - 4.10 (m, 1 H), 3.76 - 3.72 (m,
20 1 H), 3.62 - 3.58 (m, 1 H), 3.54 - 3.50 (m, 1 H), 3.47 - 3.35
(m, 3 H), 3.17 (t, J = 7.0 Hz, 2 H), 3.03 - 3.00 (m, 2 H),
2.92 - 2.83 (m, 3 H), 2.63 - 2.57 (m, 3 H), 2.49 - 2.41 (m,
3 H), 2.27 (t, J = 7.4 Hz, 2 H), 2.27 - 2.15 (obs m, 1 H),
1.95 - 1.87 (m, 3 H), 1.57 - 1.24 (m, 11 H); ¹³C NMR (125 MHz,
25 CDCl₃) δ 172.00, 171.44, 141.14, 140.44, 136.07, 128.47,
128.43, 128.29, 127.53, 126.26, 126.07, 122.17, 121.91,
119.26, 118.75, 112.71, 111.04, 101.94, 76.82, 71.69, 70.17,
69.61, 68.69, 66.44, 51.35, 35.63, 35.05, 33.59, 32.87, 30.74,
29.47, 28.73, 26.50, 26.45, 25.63, 25.58; high resolution mass
30 spectrum (+ve FAB) m/z 710.3717 (M⁺; calcd for C₄₁H₅₀N₄O₇:
710.3680).

U. 2 - (1 H - I n d o l - 3 - y l) e t h y l
2-O-(3-phenylpropanoyl)-4-O-
(4-phenylbutanoyl)-3-deoxy-O-(6-aminoheptyl)- β -D-
35 glucopyranoside (II-20).

A solution of bis-ester 25 (17.0 mg, 0.0239 mmol) and
triphenylphosphine (15.6 mg, 0.0599 mmol) in THF (2.0 ml) and

- 99 -

water (20 ml) was heated at 55°C for 16 h. The cooled solution was evaporated and purified by flash chromatography eluting with methanol/methylene chloride/acetic acid (10:90:1) increasing the gradient to (30:70:1). Fractions containing the title compound were treated with solid sodium bicarbonate, filtered, evaporated, redissolved in methylene chloride, filtered and evaporated, to afford the title compound II-20 as a colorless oil (16.0 mg, 97.8%) [α]_D²⁵ +6.25° (c 0.24, CHCl₃); IR (CHCl₃) 3480 (w), 3020 (w), 2930 (s), 2860 (m), 1740 (s), 1450 (w), 1155 (m), 1140 (m), 690 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.87 (br s, 1 H), 7.55 (d, *J* = 8.0 Hz, 1 H), 7.30 - 7.04 (m, 13 H), 6.98 (s, 1 H), 4.82 - 4.73 (m, 2 H), 5.0 - 4.7 (br s, 2 H), 4.47 (d, *J* = 7.4 Hz, 1 H), 4.10 - 4.05 (m, 1 H), 3.78 - 3.74 (m, 1 H), 3.62 - 3.58 (m, 1 H), 3.49 (dd, *J* = 2.6, 11.1 Hz, 1 H), 3.44 - 3.33 (m, 3 H), 3.02 (t, *J* = 7.3 Hz, 2 H), 2.87 (t, *J* = 7.8 Hz, 2 H), 2.63 - 2.56 (m, 3 H), 2.50 (t, *J* = 8 Hz, 2 H), 1.96 - 1.88 (m, 4 H), 1.53 - 1.22 (m, 11 H); ¹³C NMR (125 MHz, CDCl₃) δ 172.03, 171.50, 141.11, 140.38, 136.20, 128.46, 128.43, 128.28, 127.53, 126.26, 126.07, 122.26, 121.72, 119.05, 118.68, 112.19, 111.13, 101.88, 71.71, 70.09, 69.69, 68.69, 66.41, 40.46, 35.69, 35.03, 33.56, 32.86, 30.77, 29.69, 29.59, 26.51, 26.43, 25.76, 25.61.

V. 2-(1*H*-Indol-3-yl)ethyl 2,4-Di-O-(3-phenylpropanoyl)-3-deoxy-O-(6-azidohexyl)- β -D-glucopyranoside (II-22).

2,2'-dimethyl-4-aminopyridine (1 mg) was added to a stirred solution of diol II-16a (18.0 mg, 0.0417 mmol), hydrocinnamic acid (0.104 mmol, 15.7 mg) and dicyclohexylcarbodiimide (0.209 mmol, 42.9 mg) in methylene chloride (2.0 ml). The solution was stirred for 1 h, evaporated, taken up in ether, filtered and evaporated. The residue was purified by flash chromatography eluting with 30% ethyl acetate/hexane to give the title compound II-22 as a colorless oil (27.8 mg, 95.9%): [α]_D²⁵ +5.96° (c 0.94, CHCl₃); IR (CHCl₃) 3480 (m), 3010 (w), 2950 (m), 2860 (m), 2100 (s), 1745 (m), 1300 (w), 1290 (m), 1260 (w), 1160 (m), 1140 (m),

- 100 -

1080 (m), 690 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.86 (br s, 1 H), 7.57 (d, $J = 6.9$ Hz, 1 H), 7.34 - 7.02 (m, 13 H), 6.99 (d, $J = 2.4$ Hz, 1 H), 4.81 - 4.75 (m, 2 H), 4.44 (d, $J = 7.4$ Hz, 1 H), 4.14 - 4.09 (m, 1 H), 3.80 - 3.70 (m, 1 H), 3.59 - 3.55 (m, 1 H), 3.50 - 3.31 (m, 4 H), 3.18 (t, $J = 6.9$ Hz, 2 H), 3.01 (t, $J = 6.3$ Hz, 2 H), 2.91 (t, $J = 7.5$ Hz, 2 H), 2.85 (t, $J = 7.9$ Hz, 2 H), 2.63 - 2.37 (m, 5 H), 1.57 - 1.24 (m, 9 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 171.42, 140.09, 140.08, 128.48, 128.25, 128.20, 126.33, 126.23, 122.17, 121.84, 119.18, 118.70, 112.55, 111.02, 101.88, 76.67, 71.58, 70.02, 69.59, 68.63, 66.52, 51.29, 35.72, 35.59, 32.75, 30.79, 30.69, 29.42, 28.68, 26.45, 25.56; high resolution mass spectrum (-ve CI) m/z 731.3245 [$(\text{M}+\text{Cl})^+$; calcd for $\text{C}_{40}\text{H}_{48}\text{N}_4\text{O}_7\text{Cl}$: 731.3211].

W. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-(3-phenylpropanoyl)-3-deoxy-O-(6-aminohexyl)- β -D-glucopyranoside (II-21).

The same procedure as that detailed above for the preparation of compound II-20 yielded the title compound II-21 as a clear colorless oil (20.1 mg, 83.9%): $[\alpha]_{\text{D}}^{25} +23.10^\circ$ (c 0.58, CHCl_3); IR (CHCl_3) 3480 (w), 3020 (w), 2920 (m), 2850 (w), 1745 (s), 1455 (w), 1155 (m), 690 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 9.08 (br s, 1 H), 7.54 (d, $J = 7.9$ Hz, 1 H), 7.32 - 7.00 (m, 13 H), 6.96 (s, 1 H), 6.65 (br s, 1 H), 4.80 - 4.70 (m, 2 H), 4.46 (d, $J = 7.4$ Hz, 1 H), 4.08 - 4.03 (m, 1 H), 3.77 - 3.72 (m, 1 H), 3.59 - 3.55 (m, 1 H), 3.51 - 3.28 (m, 4 H), 3.02 (t, $J = 7.5$ Hz, 2 H), 2.97 - 2.86 (m, 4 H), 2.64 - 2.50 (m, 5 H), 2.42 - 2.37 (m, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 171.47, 140.06, 136.23, 128.51, 128.46, 128.26, 128.21, 127.51, 126.35, 126.25, 122.25, 121.66, 119.00, 118.66, 112.04, 111.14, 101.84, 76.90, 71.56, 69.98, 69.73, 68.67, 66.51, 35.75, 35.69, 32.77, 30.82, 30.77, 29.69, 29.57, 26.45, 25.68, 25.62.

X. 4-Azido-1-butyne (II-32).

p-Toluenesulphonyl chloride (127 mmol, 24.3 g) was added in aliquots to a solution of 3-butyne-1-ol (84.9 mmol, 5.86 g) in pyridine (20 ml) at 0°C and DMAP was added (10 mg). The mixture was allowed to stand for 15 h, then poured into

- 101 -

water (100 ml) and extracted with ether (100 ml). The ether extract was washed with 1 N HCl (100 ml), water (100 ml) and brine (50 ml), dried over sodium sulphate and evaporated to afford a yellow oil. To a stirred solution of this tosylate
5 in DMSO (100 ml) at 35°C was added sodium azide (170 mmol, 11.0 g). After stirring for 3 h, the mixture was poured into ether (50 ml), washed with water (3 x 100 ml), dried over sodium sulphate and evaporated at 0°C (water aspirator). Cautious distillation into a flask cooled to -78°C yielded the
10 pure azide II-32 as a colorless, volatile liquid (b.p. 30-32°C at 12 mmHg) (3.90 g, 48.3%): IR (CHCl₃) 3300 (s), 3000 (m), 2950 (m), 2880 (w), 2120 (s), 1450 (m), 1420 (m), 1350 (m), 1320 (m), 1290-1210 (br), 1050 (w), 950 (w), 910 (w), 630 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 3.40 (t, J = 6.9 Hz, 2 H),
15 2.48 - 2.44 (m, 2 H), 2.04 (t, J = 2.8 Hz, 1 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 80.29, 70.44, 49.62, 19.39.

Y. 5-Azido-2-pentyn-1-ol (II-33).

n-Butyl lithium (1.6 M in hexane, 18.8 mmol, 11.7 ml) was added dropwise at -78°C to a solution of 4-azido-1-butyne
20 II-32 (1.28 g, 17.3 mmol) in THF (35 ml). After stirring the resulting green solution for 1 h, paraformaldehyde was added in one portion, the solution was stirred for 5 min, and then warmed to room temperature for 2 h (an orange suspension gradually formed). The reaction mixture was poured into
25 saturated aqueous ammonium chloride (100 ml) and extracted with ether (3 x 50 ml). The combined extracts were washed with brine (50 ml), dried over sodium sulphate and evaporated to give a yellow oil. This was purified by flash chromatography eluting with pentane/ether 1:1 to afford the
30 title compound II-33 as a pale yellow oil (930 mg, 70.7%): IR (CHCl₃) 3600 (m), 3000 (m), 2940 (m), 2880 (m), 2100 (s), 1550 (w), 1380 (m), 1270 (m), 1220 (br), 1140 (m), 1000 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 4.24 - 4.22 (m, 2 H), 3.37 (t, J = 6.8 Hz, 2 H), 2.51 - 2.48 (m, 2 H), 1.84 (t, J = 6.0 Hz, 1 H);
35 ¹³C NMR (62.9 MHz, CDCl₃) δ 77.42, 70.47, 49.68, 19.45.

- 102 -

Z. 5-Azido-1-iodo-2-pentyne (II-31).

Iodine (2.94 g, 11.6 mmol) was added to a stirred solution of triphenylphosphine (12.2 mmol, 3.20 g) and imidazole (14.5 mmol, 987 mg) in THF (25 ml) at 0°C. To the
5 resulting brown solution was added 5-azido-2-pentyn-1-ol 33 (725 mg, 5.80 mmol) in THF (10 ml). The mixture was warmed to room temperature, stirred for 10 min and evaporated (water aspirator). Pentane was added and the solid was filtered off. Evaporation yielded the iodide II-31 (contaminated with a
10 small amount of triphenylphosphine) (905 mg, 66.4%).

AA. 2-[(N-Benzenesulphonyl)indol-3-yl]ethyl 4,6-Di-O-isopropylidene-3-deoxy- β -D-glucopyranoside (II-29).

Triol II-28 (25.0 mg, 0.0534 mmol) was stirred with
15 dl-camphorsulphonic acid (1 mg) in 2,2-dimethoxypropane (2.0 ml) for 15 h, triethylamine (0.05 ml) was added and the solution was evaporated. The residue was purified by flash chromatography (50% ethyl acetate/hexane) to yield the title compound II-29 as a colorless foam (26.9 mg, 99.1%): $[\alpha]_D^{25}$
20 +38.9° (c 0.99, CHCl₃); IR (CHCl₃) 3600 (w), 3010 (w), 2890 (w), 1730 (w), 1520 (w), 1450 (m), 1380 (m), 1220 (s), 1210 (s), 1180 (m), 1100 (m), 1055 (m), 930 (m), 780-720 (br), 660 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.96 (br s, 1 H), 7.84 (d, J = 7.8 Hz, 2 H), 7.52 - 7.39 (m, 4 H), 7.30 (app. t, J = 8.1
25 Hz, 1 H), 7.22 (app. t, J = 8.3 Hz, 2 H), 4.23 (d, J = 7.5 Hz, 1 H), 4.18 (dt, J = 6.6, 9.5 Hz, 1 H), 3.87 (dd, J = 5.3, 10.9 Hz, 1 H), 3.77 - 3.73 (m, 2 H), 3.64 - 3.59 (m, 1 H), 3.54 - 3.49 (m, 1 H), 3.23 - 3.19 (m, 1 H), 3.02 - 2.92 (m, 2 H), 2.27 - 2.11 (m, 1 H), 1.56 (app. q, J = 18.5 Hz, 1 H), 1.47
30 (s, 3 H), 1.39 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.25, 135.16, 133.73, 130.99, 129.23, 126.69, 124.87, 123.49, 123.20, 119.63, 119.34, 113.79, 105.43, 99.34, 71.68, 69.22, 69.03, 68.40, 62.44, 35.38, 29.11, 25.48, 19.01.

- 103 -

AB. 2 - [(N-Benzenesulphonyl)indol-3-yl]ethyl
2-O-(5-azido-2-pentynyl)-4,6-di-O-iso-
propylidene-3-deoxy- β -D-glucopyranoside (II-34).

Sodium hydride (60% dispersion in mineral oil, 0.276
5 mmol, 11.0 mg) was added to a solution of acetone II-29 (100
mg, 0.197 mmol) and 5-azido-1-iodo-2-pentyne II-31 (93 mg,
0.39 mmol) in dry acetonitrile (3.0 ml) at 0°C followed by the
addition of 15-crown-5 ether (0.001 ml). The solution was
warmed to room temperature and stirred for 36 h (a brown color
10 gradually appeared), then poured into saturated aqueous sodium
bicarbonate (10 ml) and extracted with methylene chloride (3
x 5 ml). The combined extracts were washed with brine (10
ml), dried over sodium sulphate and evaporated. The residue
was purified by flash chromatography eluting with 30% ethyl
15 acetate/hexane to afford the title compound II-34 as a
colorless oil (30.7 mg, 25.4%). The gradient was increased
to 50% ethyl acetate/hexane to yield the starting material II-
29 as a colorless oil (65 mg, 65%).

2 - [(N-Benzenesulphonyl)indol-3-yl]ethyl
20 2-O-(5-azido-2-pentynyl)-4,6-di-O-isopropylidene-3-deoxy- β -
D-glucopyranoside: $[\alpha]_D^{25} +11.59^\circ$ (c 0.63, CHCl₃); IR (CHCl₃)
3020 (m), 2950 (w), 2890 (w), 2890 (w), 2110 (s), 1450 (m),
1370 (m), 1260 (m), 1175 (s), 1090 (s), 1080 (s), 850 (w), 600
(m), 570 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.95 (d, *J* = 7.6
25 Hz, 1 H), 7.84 (d, *J* = 8 Hz, 2 H), 7.52 - 7.39 (m, 5 H), 7.29
(dt, *J* = 1.2, 7.4 Hz, 1 H), 7.23 - 7.20 (m, 1 H), 4.37 (d, *J*
= 7.5 Hz, 1 H), 4.24 - 4.20 (m, 2 H), 4.17 - 4.12 (m, 1 H),
3.86 (dd, *J* = 10.8, 5.3 Hz, 1 H), 3.81 - 3.72 (m, 2 H), 3.62
- 3.57 (m, 1 H), 3.49 - 3.42 (m, 1 H), 3.35 (t, *J* = 6.8 Hz,
30 2 H), 3.20 - 3.15 (m, 1 H), 2.96 (t, *J* = 6.2 Hz, 2 H), 2.50 -
2.46 (m, 2 H), 2.32 - 2.28 (m, 1 H), 1.52 (app. q, *J* = 11.7
Hz, 1 H), 1.47 (s, 3 H), 1.39 (s, 3 H); ¹³C NMR (62.9 MHz,
CDCl₃) δ 138.50, 135.09, 133.65, 130.95, 129.17, 126.70,
124.73, 123.11, 119.61, 119.35, 113.67, 105.00, 99.26, 82.59,
35 78.29, 74.96, 71.15, 68.59, 68.32, 62.46, 58.32, 49.74, 35.02,
29.11, 25.49, 19.84, 19.01.

EXAMPLE 11.**Preparation of Other Compounds.**

To distinguish the compounds described in this example from those described in other examples, a "III" precedes each compound number. The chemical structures and synthetic schemes of Example 11 are presented in Figure 1.

A. N-(Phenylsulfonyl)tryptophol (III-12).**(a). 1-O-tert-Butyldimethylsilyl-2-(3-indolyl)-ethanol.**

A solution of tryptophol (5.0 g, 31 mmol) in DMF (30 ml) was treated with imidazole (4.64 g, 68 mmol) and cooled to 0°C. tert-Butyldimethylsilyl chloride (5.14 g, 34.1 mmol) was added and the mixture was stirred at room temperature for 16 h. The mixture was then diluted with ethyl acetate (100 ml) and washed with water (2 x 100 ml) and the aqueous solutions were extracted with ethyl acetate (200 ml). The combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (30% ether/petroleum ether) yielded the title compound (8.43 g, 99% yield) as a colorless oil: IR (CCl₄) 3910 (s), 3060 (w), 2960 (s), 2930 (s), 2850 (s), 1450 (m), 1370 (w), 1260 (s), 1100 (s), 900 (m), 840 (s), 780 (s), 750 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.26 (br s, 1 H), 7.99 (d, J = 7.8 Hz, 1 H), 7.64-7.50 (m, 4 H), 4.28 (t, J = 7.3 Hz, 2 H), 3.38 (t, J = 7.3 Hz, 2 H), 1.29 (s, 9 H), -0.43 (s, 6 H); ¹³C NMR (125 MHz, CDCl₃) δ 136.08, 127.62, 122.08, 121.75, 119.12, 118.79, 112.84, 111.04, 63.89, 28.98, 25.98, 18.34, -5.29; high resolution mass spectrum (Cl, NH₃) m/z 276.1750 [(M+H)⁺; calcd for C₁₆H₂₅NOSi: 276.1783].

(b). 1-O-tert-Butyldimethylsilyl-2-[3-(1-N-phenyl-sulfonyl)indolyl]ethanol.

A suspension of sodium hydride (1.91 g, 60% oil dispersion) in dry DMF (64 ml) was cooled to 0°C and a solution of 1-O-tert-butyldimethylsilyl-2-(3-indolyl)ethanol (8.43 g, 30.6 mmol) in DMF (30 ml) was added. The mixture was stirred at room temperature for 30 min, recooled to 0°C, and treated dropwise with benzenesulfonyl chloride (5.30 ml, 39.7

- 105 -

mmol). The reaction was then stirred at room temperature for 16 h, quenched with saturated aqueous ammonium chloride (100 ml), and extracted with ether (3 x 200 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (30% ether/petroleum ether) afforded the title compound (7.37 g, 79% yield) as a colorless oil: ^1H NMR (500 MHz, CDCl_3) δ 7.77 (d, J = 8.4 Hz, 1 H), 7.62 (d, J = 7.5 Hz, 2 H), 7.26-6.98 (m, 7 H), 3.64 (t, J = 6.7 Hz, 2 H), 2.64 (t, J = 6.7 Hz, 2 H), 0.64 (s, 9 H), -0.24 (s, 6 H); ^{13}C NMR (125 MHz, CDCl_3) δ 135.10, 133.55, 131.21, 129.12, 126.65, 124.56, 123.42, 122.00, 120.31, 119.57, 113.59, 62.51, 28.51, 25.87, 18.22, -5.44; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 433.1920 $[(\text{M}+\text{NH}_4)^+]$; calcd for $\text{C}_{22}\text{H}_{29}\text{NSO}_3\text{Si}$: 433.1971].

(c). N-Phenylsulfonyltryptophol (III-12).

Tetrabutylammonium fluoride (21 ml, 1 M in THF) was added to a solution of 1-O-tert-butyldimethylsilyl-2-[3-(1-N-phenyl-sulfonyl)indolyl]ethanol (6.6 g, 22 mmol) in THF (100 ml) and the reaction was stirred at room temperature for 16 h. The mixture was then diluted with ethyl acetate (100 ml) and extracted with water (2 x 100 ml). The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (40% ethyl acetate/petroleum ether) furnished III-11 (4.00 g, 84% yield) as a pale yellow oil which crystallized upon standing: mp 63-64°C; IR (CCl_4) 3580 (m), 3400 (m), 3100 (w), 3080 (w), 2950 (m), 2890 (m), 1460 (s), 1360 (s), 1280 (m), 1160 (s), 1120 (s), 1100 (m), 1080 (w), 1060 (w), 1020 (w), 980 (w), 750 (s), 720 (s), 690 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.86 (d, J = 7.6 Hz, 1 H), 7.70 (d, J = 7.6 Hz, 2 H), 7.32-7.04 (m, 7 H), 3.68 (t, J = 6.2 Hz, 2 H), 2.72 (t, J = 6.2 Hz, 2 H), 2.36 (br s, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 137.79, 134.99, 133.55, 130.78, 129.00, 126.43, 124.63, 123.39, 123.05, 119.67, 119.38, 61.40, 28.07; high resolution mass spectrum (Cl , NH_3) m/z 301.0748 (M^+ ; calcd for $\text{C}_{16}\text{H}_{15}\text{NO}_3\text{S}$: 301.0772).

- 106 -

B. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3,4,6-Tetra-O-acetyl- β -D-glucopyranoside (III-13).

A solution of III-12 (537 mg, 1.78 mmol) in dry benzene
5 (3 ml) was added to a suspension of powdered, activated 4
Angstrom molecular sieves (0.89 g) and silver(I) oxide (412
mg, 17.8 mmol) in dry hexane (9 ml) at room temperature. A
solution of bromide III-11 (804 mg, 1.95 mmol) in dry benzene
(3 ml) was then added, the flask was covered with aluminum
10 foil and the mixture allowed to stir for 2 days at room
temperature. More silver(I) oxide (206 mg, 8.9 mmol) and
benzene (1 ml) were added and the reaction was stirred at room
temperature for an additional 2 days. After filtration
through Celite, concentration in vacuo and recrystallization
15 (ethyl acetate/petroleum ether) afforded pure II-13 (580 mg)
as a white solid. Concentration of the filtrate in vacuo and
flash chromatography (5% ether/dichloromethane) afforded III-
13 admixed with the α anomer and the corresponding ortho
ester. Further flash chromatography (70% ether/petroleum
20 ether) then gave an additional 134 mg of pure III-13 (64%
total yield): mp 145-146°C; $[\alpha]_D^{25}$ -16° (c 0.14,
acetonitrile); UV (1.05 x 10⁻⁴ M, acetonitrile) λ_{max} 253.6 (ϵ
1.19 x 10⁴), 214.0 (2.50 x 10⁴) nm; IR (thin film) 3028 (w),
2950 (w), 2880 (w), 1758 (s), 1450 (m), 1377 (s), 1225 (s),
25 1178 (s), 1122 (m), 1040 (s), 910 (w), 754 (s), 688 (w) cm⁻¹;
¹H NMR (500 MHz, CDCl₃) δ 7.96 (d, J = 8.3 Hz, 1 H), 7.87-7.21
(m, 9 H), 5.18 (dd, J = 9.5, 9.5 Hz, 1 H), 5.09 (dd, J = 9.6,
9.6 Hz, 1 H), 5.00 (dd, J = 9.5, 8.0 Hz, 1 H), 4.53 (d, J =
8.0 Hz, 1 H), 4.26 (dd, J = 12.3, 4.7 Hz, 1 H), 4.18-4.12 (m,
30 2 H), 3.76 (ddd, J = 9.3, 6.9, 6.9 Hz, 1 H), 3.69 (ddd, J =
9.8, 4.6, 2.4 Hz, 1 H), 2.94 (t, J = 6.6 Hz, 2 H), 2.07 (s,
3 H), 2.02 (s, 3 H), 2.00 (s, 3 H), 1.89 (s, 3 H); ¹³C NMR
(125 MHz, CDCl₃) δ 170.66, 170.24, 169.34, 138.24, 135.08,
133.70, 130.94, 129.22, 126.73, 124.75, 123.56, 123.21,
35 119.57, 119.42, 113.65, 106.61, 100.70, 72.87, 71.16, 68.75,
68.39, 61.91, 25.31, 20.72, 20.57, 20.43; high resolution mass
spectrum (Cl, NH₃) m/z 649.2021 [(M+NH₄)⁺; calcd for C₃₀H₃₃NO₁₂S:

- 107 -

649.2054]. Anal. Calcd for $C_{30}H_{33}NO_{12}S$: C, 57.04; H, 5.27. Found: C, 56.75; H, 5.30.

C. 2-(N-Phenylsulfonylindol-3-yl)ethyl
 β -D-glucopyranoside (III-14).

5 Sodium methoxide (221 mg, 4.09 mmol) was added to a suspension of III-13 (3.22 g, 5.12 mmol) in methanol (26 ml) at room temperature. After 20 min, the resultant solution was diluted with methanol (26 ml) and neutralized with Amberlyst®
15 ion exchange resin. The resin was quickly removed by filtration to avoid formation of the methyl glucoside. Concentration and flash chromatography (5:1:1 dichloromethane/methanol/acetone) afforded III-13 (2.09, 88% yield) as a white foam: $[\alpha]_D^{25}$ -23° (c 0.09, acetonitrile); UV (1.62×10^{-4} M, acetonitrile) λ_{max} 253.6 (ϵ 1.17×10^4),
214.0 (1.93×10^4) nm; IR (film) 3390 (s), 3065 (w), 3015 (w), 2920 (m), 2880 (m), 1450 (s), 1363 (s), 1282 (m), 1175 (s), 1123 (s), 1085 (s), 1021 (s), 748 (s), 725 (m), 686 (m), cm^{-1} ; 1H NMR (500 MHz, acetone- d_6) δ 8.00-7.97 (m, 3H), 7.71 (s, 1H), 7.64-7.53 (m, 4H), 7.35-7.31 (m, 1H), 7.26-7.23 (m, 1H),
20 4.40 (d, J = 7.7 Hz, 1H), 4.30 (d, J = 3.7 Hz, 1H), 4.25 (d, J = 3.7 Hz, 1H), 4.22 (d, J = 4.0 Hz, 1H), 4.16 (ddd, J = 9.7, 6.7, 6.7 Hz, 1H), 3.89-3.82 (m, 2H), 3.70 (ddd, J = 11.8, 5.9, 5.9 Hz, 1H), 3.58 (t, J = 6.4 Hz, 1H), 3.45 (ddd, J = 8.8, 8.8, 3.8 Hz, 1H), 3.39 (ddd, J = 8.5, 8.5, 4.0 Hz, 1H), 3.34
25 (ddd, J = 9.3, 5.5, 2.7 Hz, 1H), 3.25 (ddd, J = 8.6, 7.8, 3.8 Hz, 1H), 2.98 (t, J = 6.6 Hz, 2H); ^{13}C NMR (125 MHz, acetone- d_6) δ 139.40, 136.57, 134.87, 132.21, 130.31, 127.67, 125.41, 125.30, 124.07, 121.33, 120.56, 114.35, 104.07, 78.07, 77.53, 74.93, 71.73, 68.76, 63.00, 49.72, 25.92; high
30 resolution mass spectrum (CI, NH_3) m/z 481.1656 [(M+ NH_4) $^+$; calcd for $C_{22}H_{25}NO_8S$: 481.1634].

D. 2-(N-Phenylsulfonylindol-3-yl)ethyl
6-O-tert-Butyldiphenylsilyl- β -D-glucopyranoside
(III-15).

35 At room temperature a stirred solution of III-14 (7.11 g, 15.4 mmol) in dry DMF (51 ml) was treated with imidazole (2.93 g, 43.1 mmol) followed by tert-butyldiphenylsilyl

- 108 -

chloride (5.58 g, 21.6 mmol). The solution was heated at 50°C for 24 h. After concentration in vacuo, the mixture was diluted with ethyl acetate (250 ml) and washed with water (100 ml). The organic phase was then washed with brine (100 ml),
5 dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (5% methanol/dichloromethane) provided pure III-15 (9.15 g, 85% yield) as a white foam: $[\alpha]_D^{25}$ -26° (c 0.14, acetonitrile); UV (5×10^{-5} M, acetonitrile) λ_{\max} 280.0 (ϵ 7.1×10^3), 220.8 (5.17×10^4) nm;
10 IR (film) 3410 (s), 3070 (w), 3045 (w), 3010 (w), 2925 (m), 2885 (m), 2855 (m), 1474 (w), 1458 (w), 1430 (m), 1363 (w), 1220 (w), 1113 (s), 1047 (s), 1010 (s), 823 (m), 805 (w), 742 (s), 704 (s) cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 8.02 (d, J = 8.3 Hz, 1 H), 7.96-7.95 (m, 2 H), 7.78-7.74 (m, 4 H), 7.70 (s,
15 1 H), 7.57 (d, J = 7.8 Hz, 1 H), 7.54-7.50 (m, 1 H), 7.47-7.43 (m, 2 H), 7.39-7.30 (m, 7 H), 7.21-7.18 (m, 1 H), 4.49 (m, 2 H), 4.46 (d, J = 7.7 Hz, 1 H), 4.20 (ddd, J = 9.7, 6.7, 6.7 Hz, 1 H), 4.11 (dd, J = 11.2, 0.9 Hz, 1 H), 3.96 (dd, J = 11.0, 4.9 Hz, 1 H), 3.85 (ddd, J = 9.7, 6.9, 6.9 Hz, 1 H),
20 3.52 (m, 2 H), 3.38-3.34 (m, 1 H), 3.05 (t, J = 6.6 Hz, 2 H), 2.86 (s, 1 H), 2.75 (s, 1 H), 1.02 (s, 9 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 206.17, 138.97, 136.39, 136.30, 135.95, 134.83, 134.60, 134.47, 132.13, 130.45, 130.41, 130.26, 128.47, 127.59, 125.40, 125.01, 124.04, 121.24, 120.60, 114.31,
25 104.11, 78.17, 77.76, 74.94, 71.14, 68.93, 64.72, 27.12, 26.10, 19.82; high resolution mass spectrum (Cl, NH_3) m/z 684.2532 [(M-OH) $^+$; calcd for $\text{C}_{38}\text{H}_{43}\text{NO}_8\text{SSi}$: 684.2449]. Anal. Calcd for $\text{C}_{38}\text{H}_{43}\text{NO}_8\text{SSi}$: C, 65.03; H, 6.18. Found: C, 64.96; H, 6.28.

30 E. 2-(N-Phenylsulfonylindol-3-yl)ethyl
2,3,4-Tri-O-benzyl-6-O-tert-butyl-diphenylsilyl-
 β -D-glucopyranoside (III-16).

A solution of III-15 (1.62 g, 2.31 mmol) in THF (7 ml) was added to a stirred suspension of sodium hydride (323 mg, 60% oil dispersion, 8.08 mmol) in THF (5 ml) at 0°C. After
35 the mixture was stirred for 1 h at room temperature and recooled to 0°C, benzyl bromide (1.09 ml, 9.24 mmol) was added

- 109 -

dropwise followed by tetrabutylammonium iodide (85 mg, 0.23 mmol). The reaction was then allowed to stir for 3 days at room temperature. The resultant suspension was diluted with saturated aqueous ammonium chloride (3 ml) at 0°C and
5 extracted with ether (2 x 80 ml). The combined extracts were washed with saturated aqueous ammonium chloride (30 ml) and brine (30 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (20% ether/petroleum ether) afforded III-16 (1.66 g, 74% yield) as
10 a white foam: $[\alpha]_D^{25} -7.0^\circ$ (c 0.12, acetonitrile); UV (5.90 x 10⁻⁵ M, acetonitrile) λ_{\max} 253.6 (ϵ 2.90 x 10³), 213.6 (5.11 x 10⁴) nm; IR (film) 3065 (m), 3030 (m), 2930 (s), 2855 (s), 1608 (w), 1590 (w), 1496 (w), 1472 (w), 1464 (w), 1449 (s), 1429 (m), 1377 (s), 1338 (w), 1312 (w), 1280 (m), 1215 (m),
15 1176 (s), 1113 (s), 1088 (s), 1072 (s), 1029 (s), 952 (w), 920 (w), 825 (m), 805 (w), 746 (s), 700 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.99 (d, J = 8.3 Hz, 1 H), 7.82 (d, J = 7.5 Hz, 2 H), 7.73 (d, J = 6.7 Hz, 2 H), 7.68 (d, J = 6.7 Hz, 2 H), 7.50 (d, J = 7.8 Hz, 1 H), 7.44-7.17 (m, 27 H), 4.91 (d, J = 10.9 Hz, 1 H), 4.88 (d, J = 11.2 Hz, 1 H), 4.80 (d, J = 10.7 Hz, 1 H),
20 4.77 (d, J = 11.2 Hz, 1 H), 4.68 (d, J = 10.8 Hz, 1 H), 4.63 (d, J = 10.8 Hz, 1 H), 4.44 (d, J = 7.7 Hz, 1 H), 4.19 (dd, J = 14.6, 7.1 Hz, 1 H), 3.92 (d, J = 2.9 Hz, 2 H), 3.81 (dd, J = 15.4, 7.6 Hz, 1 H), 3.74 (dd, J = 8.8, 8.8 Hz, 1 H), 3.64
25 (dd, J = 9.1, 9.1 Hz, 1 H), 3.46 (dd, J = 8.1, 8.1 Hz, 1 H), 3.35 (apparent d, J = 7.6 Hz, 1 H), 3.05 (t, J = 7.0 Hz, 2 H), 1.04 (s, 9 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.58, 138.47, 138.32, 138.19, 135.83, 135.35, 135.23, 133.64, 133.58, 133.18, 130.96, 129.60, 129.13, 128.39, 128.30, 127.97,
30 127.90, 127.72, 127.66, 127.55, 127.51, 126.63, 124.77, 123.38, 123.16, 119.74, 119.57, 113.71, 103.62, 84.71, 82.55, 77.66, 75.81, 75.79, 75.10, 74.80, 68.53, 62.80, 26.78, 25.90, 19.29; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 972.4071 [(M+H)⁺; calcd for C₅₉H₆₁NO₈SSi:
35 972.3970].

- 110 -

F. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3,4-Tri-O-benzyl- β -D-glucopyranoside (III-17).

Tetrabutylammonium fluoride (1 M in THF, 2.4 ml, 2.4 mmol) was added to a stirred solution of III-16 (1.55 g, 1.60 mmol) in THF (8 ml) at room temperature. After 7 h the reaction mixture was diluted with ethyl acetate (70 ml), washed with water (30 ml) and brine (30 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (30% ethyl acetate/petroleum ether) afforded III-17 (1.10 g, 94% yield) as a clear oil: $[\alpha]_D^{25} -13^\circ$ (c 0.14, acetonitrile); UV (9.21×10^{-5} M, acetonitrile) λ_{\max} 254.0 (ϵ 2.81×10^3), 211.6 (3.19×10^4) nm; IR (film) 3480 (w), 3065 (w), 3035 (w), 2920 (m), 2878 (m), 1498 (w), 1450 (s), 1365 (s), 1280 (w), 1220 (m), 1176 (s), 1123 (s), 1090 (s), 1073 (s), 1030 (s), 750 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.84 (d, $J = 8.3$ Hz, 1 H), 7.82 (d, $J = 7.9$ Hz, 2 H), 7.53 (s, 1 H), 7.48-7.17 (m, 21 H), 4.92 (d, $J = 11.0$ Hz, 1 H), 4.86 (d, $J = 10.9$ Hz, 1 H), 4.81 (d, $J = 11.0$ Hz, 1 H), 4.74 (d, $J = 11.0$ Hz, 1 H), 4.65 (d, $J = 10.9$ Hz, 1 H), 4.62 (d, $J = 11.0$ Hz, 1 H), 4.48 (d, $J = 7.8$ Hz, 1 H), 4.20 (ddd, $J = 9.4, 7.0, 7.0$ Hz, 1 H), 3.91-3.86 (m, 2 H), 3.73 (dd, $J = 3.5, 11.9$ Hz, 1 H), 3.63 (ddd, $J = 9.0, 9.0, 18.0$ Hz, 2 H), 3.40 (apparent t, $J = 8.0$ Hz, 1 H), 3.35 (ddd, $J = 9.4, 4.2, 2.6$ Hz, 1 H), 3.04-2.93 (m, 2 H), 2.06 (s, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.48, 138.21, 138.13, 137.95, 135.09, 133.60, 130.92, 129.10, 128.40, 128.30, 128.25, 128.22, 127.98, 127.90, 127.82, 127.76, 127.55, 126.58, 124.72, 123.57, 123.12, 119.61, 119.31, 113.66, 103.59, 84.39, 82.25, 77.37, 75.56, 75.16, 74.99, 74.75, 68.60, 61.77, 25.57; high resolution mass spectrum (CI, NH_3) m/z 734.2743 [(M+H) $^+$; calcd for $\text{C}_{43}\text{H}_{43}\text{NO}_8\text{S}$: 734.2774]. Anal. Calcd for $\text{C}_{43}\text{H}_{43}\text{NO}_8\text{S}$: C, 70.37; H, 5.91. Found: C, 70.30; H, 6.08.

G. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3,4-Tri-O-benzyl-6-O-(5-azidopentyl)- β -D-
glucopyranoside (III-19a).

Sodium azide (1.83 g, 28.2 mmol) was added to a stirred solution of 5-bromo-1-pentanol (0.79 g, 4.7 mmol) in DMSO (15

- 111 -

ml). The resultant mixture was stirred at room temperature for 2.5 h, diluted with water, and extracted with diethyl ether. The combined organic solutions were washed with saturated aqueous sodium bicarbonate and brine, dried over
5 magnesium sulfate, filtered, and concentrated in vacuo. The azide was used without purification in the next step.

A stirred solution of crude 5-azido-1-pentanol (280 mg, equivalent to 2.17 mmol) and 2,6-di-tert-butyl-4-methylpyridine (441 mg, 2.17 mmol) in
10 dichloromethane (9 ml) was treated dropwise with triflic anhydride (0.36 ml, 2.17 mmol). After 10 min the mixture was poured into brine (40 ml) and extracted with dichloromethane (2 x 40 ml). The combined organic layers were dried over magnesium sulfate, filtered, and concentrated in vacuo. The
15 resultant triflate was used without purification in the next step. Sodium hydride (12.4 mg, 0.31 mmol, 60% dispersion in oil) was added to a solution of alcohol 17 (225 mg, 0.309 mmol) and crude azidotriflate (161 mg, equivalent to 0.62 mmol) in dichloromethane (4 ml) at room temperature. The
20 mixture was stirred for 24 h, diluted with dichloromethane (40 ml), and poured into saturated aqueous ammonium chloride (40 ml). The aqueous phase was extracted with dichloromethane and the combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash
25 chromatography (15% ethyl acetate/hexane) furnished III-19a (248 mg, 95% yield) as a colorless oil: $[\alpha]_D^{25} +1.3^\circ$ (c 0.48, CHCl₃); IR (CHCl₃) 3070 (w), 3015 (m), 2935 (s), 2875 (s), 2100 (s), 1450 (s), 1370 (s), 1280 (w), 1178 (m), 1122 (m), 1070 (s), 695 (m), 597 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃)
30 δ 7.93 (d, J = 8.4 Hz, 1 H), 7.78 (apparent d, J = 8.4 Hz, 2 H), 7.44-7.41 (m, 3 H), 7.39-7.10 (m, 19 H), 4.86 (d, J = 10.9 Hz, 1 H), 4.81 (d, J = 10.9 Hz, 1 H), 4.73 (d, J = 11.0 Hz, 1 H), 4.67 (d, J = 11.0 Hz, 1 H), 4.56 (d, J = 10.9 Hz, 1 H), 4.54 (d, J = 11.0 Hz, 1 H), 4.36 (dd, J = 7.8, 1.0 Hz, 1 H),
35 4.15 (dt, J = 9.5, 7.1 Hz, 1 H), 3.79 (dt, J = 9.5, 7.3 Hz, 1 H), 3.64-3.44 (m, 5 H), 3.36 (m, 3 H), 3.13 (t, J = 7.0 Hz, 2 H), 2.96 (t, J = 7.0 Hz, 2 H), 1.56-1.48 (m, 4 H), 1.39-1.31

- 112 -

(m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.56, 138.31, 138.28, 138.22, 135.18, 133.60, 130.96, 129.13, 128.42, 128.35, 128.28, 128.00, 127.85, 127.82, 127.77, 127.57, 127.51, 126.67, 124.74, 123.47, 123.11, 119.65, 119.44, 113.72, 103.74, 84.64, 82.25, 77.93, 75.66, 74.97, 74.90, 74.75, 71.40, 69.70, 68.76, 29.67, 29.18, 28.66, 25.71, 23.41; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 867.3532 (M^+ ; calcd for $\text{C}_{48}\text{H}_{52}\text{N}_4\text{O}_8\text{S}$: 867.3494).

H. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-

10 6-O-(5-aminoentyl)- β -D-glucopyranoside (III-4a).

A stirred solution of azide III-19a (31 mg, 0.037 mmol) in THF (2 ml) and water (0.032 ml) was treated with triphenylphosphine (25 mg, 0.095 mmol). The mixture was heated at reflux for 2.5 h, cooled, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) furnished the corresponding amine (26 mg, 86% yield) as a colorless oil: ($[\alpha]_D^{25}$) xx° (c 0.xx, CHCl_3); IR (CHCl_3) $xxx\text{ cm}^{-1}$; ^1H NMR (500 MHz, CDCl_3) δ 8.03 (d, J = 8.2 Hz, 1 H), 7.89 (dd, J = 8.5, 0.9 Hz, 2 H), 7.39-7.21 (m, 22 H), 4.96 (d, J = 10.9 Hz, 1 H), 4.91 (d, J = 10.9 Hz, 1 H), 4.84 (d, J = 10.9 Hz, 1 H), 4.78 (d, J = 11.3 Hz, 1 H), 4.67 (d, J = 10.8 Hz, 1 H), 4.65 (d, J = 11.0 Hz, 1 H), 4.47 (d, J = 7.8 Hz, 1 H), 4.26 (dt, J = 9.5, 6.9 Hz, 1 H), 3.90 (dt, J = 9.5, 7.1 Hz, 1 H), 3.75-3.62 (m, 4 H), 3.56 (dt, J = 9.4, 6.5 Hz, 1 H), 3.49-3.44 (m, 3 H), 3.06 (t, J = 6.9 Hz, 2 H), 2.68 (t, J = 6.9 Hz, 2 H), 1.91 (br s, 2 H), 1.66-1.58 (m, 2 H), 1.50-1.34 (m, 4 H); ^{13}C NMR (62.5 MHz, CDCl_3) δ 138.49, 138.23, 138.14, 135.25, 133.56, 133.20, 132.08, 131.56, 131.90, 130.09, 129.08, 128.52, 128.32, 128.23, 127.93, 127.79, 127.52, 126.59, 124.67, 123.39, 123.06, 119.60, 119.40, 113.62, 103.65, 84.56, 82.17, 77.85, 75.60, 74.91, 74.80, 74.68, 71.56, 69.56, 68.68, 41.88, 33.18, 29.37, 25.63, 23.36; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 819.3687 (M^+ ; calcd for $\text{C}_{48}\text{H}_{54}\text{N}_2\text{O}_8\text{S}$: 819.3679).

35 The above amine (26 mg, 0.032 mmol) was dissolved in ethanol (4 ml) and treated with 5 M aqueous sodium hydroxide (0.65 ml). The resultant mixture was heated at reflux for 3

- 113 -

h, cooled, diluted with brine, and poured into dichloromethane. The aqueous layer was extracted with dichloromethane (2 x 40 ml) and the combined organic solutions were dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) afforded III-4a (19.7 mg, 91% yield) as a colorless oil: $[\alpha]_D^{25} +13^\circ$ (c 0.03, CHCl_3); IR (CHCl_3) 3009 (s), 2930 (m), 2860 (m), 1450 (w), 1360 (w), 1200 (s), 1062 (s), 920 (w), 690 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.75 (br s, 1 H), 7.59 (d, $J = 7.9$ Hz, 1 H), 7.38-7.24 (m, 16 H), 7.17 (t, $J = 7.2$ Hz, 1 H), 7.10 (t, $J = 7.2$ Hz, 1 H), 7.07 (s, 1 H), 4.93 (d, $J = 10.9$ Hz, 1 H), 4.89 (d, $J = 11.0$ Hz, 1 H), 4.85 (d, $J = 11.0$ Hz, 1 H), 4.80 (d, $J = 10.9$ Hz, 1 H), 4.71 (d, $J = 11.0$ Hz, 1 H), 4.57 (d, $J = 11.0$ Hz, 1 H), 4.48 (d, $J = 7.8$ Hz, 1 H), 4.18 (dt, $J = 9.4, 7.1$ Hz, 1 H), 3.88 (dt, $J = 9.4, 7.1$ Hz, 1 H), 3.68-3.64 (m, 2 H), 3.55-3.35 (m, 6 H), 3.12 (t, $J = 7.1$ Hz, 2 H), 2.43 (br t, $J = 7.1$ Hz, 2 H), 1.59-1.54 (m, 2 H), 1.52-1.54 (m, 2 H), 1.37-1.28 (m, 4 H); ^{13}C NMR (62.5 MHz, CDCl_3) δ 138.48, 138.20, 138.05, 136.14, 130.90, 128.97, 128.45, 128.37, 128.07, 127.88, 127.61, 127.40, 122.47, 121.87, 119.17, 118.64, 112.15, 111.44, 103.70, 84.62, 82.29, 77.88, 77.21, 75.68, 74.97, 74.79, 74.56, 71.03, 70.46, 69.51, 66.80, 29.69, 28.89, 28.64, 25.77, 22.95; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) calcd for $\text{C}_{42}\text{H}_{50}\text{N}_2\text{O}_6$.

I. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-O-(4-azidobutyl)- β -D-glucopyranoside (III-19b).

Alcohol 17 (0.164 g, 0.223 mmol) and 2,6-di-tert-butyl-4-methyl-pyridine (0.06 g, 0.29 mmol) were dissolved in dichloromethane (5 ml) and triflic anhydride (0.041 ml, 0.246 mmol) was added dropwise. The mixture was stirred at room temperature for 10 min, diluted with dichloromethane (40 ml), and poured into brine (40 ml). The organic phase was dried over magnesium sulfate, filtered, and concentrated. The resultant white solid was redissolved in dichloromethane (3 ml) and treated sequentially with 4-azido-1-butanol (0.13 g, 1.21 mmol), prepared in a similar manner to 5-azido-1-pentanol above, and sodium hydride (0.045 g, 1.13 mmol, 60% dispersion

- 114 -

in oil). The mixture was then stirred for 24 h, diluted with dichloromethane (40 ml), and poured into saturated aqueous ammonium chloride (40 ml). The aqueous phase was extracted with dichloromethane (2 x 20 ml) and the combined organic solutions were washed with brine (40 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (15% ethyl acetate/hexane) yielded III-19b (85.2 mg, 56% yield) as a colorless oil: $[\alpha]_D^{25} +10.2^\circ$ (c 0.3, CH_2Cl_2); IR (CH_2Cl_2) 3485 (m), 3044 (w), 2910 (m), 2885 (m), 2090 (s), 1735 (m), 1610 (w), 1460 (m), 1420 (m), 1360 (m), 1250 (m), 1060 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.84 (br s, 1 H), 7.60 (d, $J = 7.8$ Hz, 1 H), 7.16-7.33 (m, 17 H), 7.11 (apparent t, $J = 7.2$ Hz, 1 H), 7.03 (br s, 1 H), 4.91 (d, $J = 10.9$ Hz, 1 H), 4.86 (d, $J = 11.0$ Hz, 1 H), 4.80 (d, $J = 11.0$ Hz, 1 H), 4.78 (d, $J = 10.9$ Hz, 1 H), 4.64 (d, $J = 11.0$ Hz, 1 H), 4.59 (d, $J = 7.8$ Hz, 1 H), 4.43 (d, $J = 7.8$ Hz, 1 H), 4.24 (dt, $J = 9.3, 6.8$ Hz, 1 H), 3.86 (dt, $J = 9.3, 7.4$ Hz, 1 H), 3.68-3.60 (m, 3 H), 3.57-3.51 (m, 2 H), 3.44 (t, $J = 5.9$ Hz, 2 H), 3.46-3.40 (m, 1 H), 3.24 (br t, $J = 6.5$ Hz, 2 H), 3.12 (t, $J = 6.9$ Hz, 2 H), 1.65-1.62 (m, 4 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.60, 138.56, 138.24, 136.17, 128.43, 128.28, 128.04, 127.90, 127.86, 127.78, 127.60, 127.53, 122.12, 121.96, 119.29, 118.73, 112.81, 111.10, 103.71, 84.70, 82.33, 77.99, 75.69, 74.97, 74.84, 74.69, 70.97, 70.05, 69.76, 51.29, 26.88, 25.84, 25.81.

J. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-O-(4-aminobutyl)- β -D-glucopyranoside (III-4b).

A solution of azide III-19b (0.037 g, 0.056 mmol) in THF (3 ml) was treated sequentially with water (0.025 ml, 1.39 mmol) and triphenylphosphine (0.29 g, 0.11 mmol). The mixture was then heated at 60°C for 6 h, cooled, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) yielded III-4b (26.6 mg, 72% yield) as a colorless oil: $[\alpha]_D^{25}$ (CH_2Cl_2); IR (CH_2Cl_2) 3700 (w), 3487 (m), 3028 (m), 3020 (m), 2918 (s), 2878 (s), 1608 (w), 1498 (w), 1277 (m), 1212 (m), 1072 (s), 1465 (s), 1371 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.29 (br s, 1 H), 7.60 (d, $J = 7.8$ Hz, 1 H), 7.34-7.60 (m,

- 115 -

18 H), 7.09 (br s, 1 H), 4.92 (d, $J = 10.9$ Hz, 1 H), 4.86 (d, $J = 10.9$ Hz, 1 H), 4.83 (d, $J = 11.0$ Hz, 1 H), 4.79 (d, $J = 10.9$ Hz, 1 H), 4.66 (d, $J = 11.0$ Hz, 1 H), 4.61 (d, $J = 10.9$ Hz, 1 H), 4.45 (d, $J = 7.8$ Hz, 1 H), 4.24 (dt, $J = 9.3, 6.9$ Hz, 1 H), 3.89 (dt, $J = 9.3, 7.1$ Hz, 1 H), 3.12 (t, $J = 6.9$ Hz, 2H), 2.66 (t, $J = 6.8$ Hz, 2 H), 1.62-1.47 (m, 6 H); ^{13}C NMR (125 MHz, CDCl_3) δ 137.55, 137.49, 137.21, 135.11, 127.34, 127.27, 127.20, 126.96, 126.80, 126.67, 126.49, 126.45, 121.18, 120.75, 118.09, 117.59, 111.69, 110.04, 102.60, 83.61, 81.28, 76.09, 74.59, 73.90, 73.73, 73.63, 70.42, 68.88, 68.51, 40.90, 29.29, 26.00, 24.69.

K. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3,4-Tri-O-benzyl-6-O-(6-azidohexyl)- β -D-glucopyranoside (III-19c).

15 A stirred solution of 6-azido-1-hexanol (0.087 g, 0.61 mmol), prepared in a manner similar to 5-azido-1-pentanol above, and 2,6-di-tert-butyl-4-methylpyridine (0.125 g, 0.061 mmol) in dichloromethane (5 ml) was treated with triflic anhydride (0.1 ml, 0.61 mmol) at room temperature. After 15
20 min the solution was diluted with dichloromethane (20 ml) and poured into saturated aqueous sodium bicarbonate (20 ml). The organic phase was washed with brine, dried over magnesium sulfate, filtered, and concentrated, to afford a white semisolid which was used without purification. A solution of
25 the alcohol III-17 (0.3 g, 0.41 mmol) and the crude triflate in dichloromethane (3 ml) was treated with sodium hydride (0.024g, 0.6 mmol, 66% dispersion in oil) followed by 15-crown-5 (10 mg). The mixture was then stirred at ambient temperature for 48 h, diluted with dichloromethane (25 ml),
30 and poured into saturated aqueous ammonium chloride (20 ml). The aqueous phase was extracted with dichloromethane (2 x 20 ml) and the combined organic solutions were washed with brine (25 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (15% ethyl
35 acetate/hexane) furnished III-19c (302 mg, 86% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -4.8^\circ$ (c 1.06, CH_2Cl_2); IR (solvent?) 3030 (m), 2991 (w), 2920 (m), 2832 (m), 2110 (s), 1720 (w),

- 116 -

1609 (w), 1450 (s), 1372 (s), 1252 (s), 1212 (w), 1180 (s),
1122 (s), 1091 (s), 1071 (s), 892 (w), 692 (br), 600 (s), 573
(s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.97 (d, J = 7.8 Hz, 1 H),
7.83 (dd, J = 8.5, 1.1 Hz, 2 H), 7.50-7.16 (m, 22 H), 4.91 (d,
5 J = 10.9 Hz, 1 H), 4.85 (d, J = 10.9 Hz, 1 H), 4.78 (d, J =
10.9 Hz, 1 H), 4.73 (d, J = 11.0 Hz, 1 H), 4.61 (d, J = 10.9
Hz, 1 H), 4.41 (d, J = 7.7 Hz, 1 H), 4.20 (dt, J = 9.4, 7.1
Hz, 1 H), 3.83 (dt, J = 9.4, 7.5 Hz, 1 H), 3.69-3.56 (m, 4 H),
3.53-3.48 (m, 1 H), 3.43-3.40 (m, 3 H), 3.19 (t, J = 6.9 Hz,
10 2 H), 3.01 (t, J = 7.0 Hz, 2 H), 1.63-1.20 (m, 8 H); ^{13}C NMR
(125 MHz, CDCl_3) δ 138.56, 138.37, 138.27, 138.23, 135.17,
133.59, 130.96, 129.12, 128.41, 128.33, 128.27, 127.99,
127.84, 127.75, 127.57, 127.56, 126.66, 124.72, 123.46,
123.11, 119.64, 119.44, 113.70, 103.74, 84.64, 82.24, 77.93,
15 75.66, 74.96, 74.89, 74.73, 71.52, 69.65, 68.75, 51.33, 29.48,
28.72, 26.52, 25.73, 25.71; high resolution mass spectrum
(FAB, m-nitrobenzyl alcohol) m/z 881.3538 [(M+ Na) $^+$; calcd for
 $\text{C}_{29}\text{H}_{34}\text{N}_4\text{O}_6\text{S}$: 881.3560].

20 L. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-O-
(6-aminohexyl)- β -D-glucopyranoside (III-4c).

A solution of azide III-19c (0.234 g, 0.272 mmol) in
THF (15 ml) was treated sequentially with water (0.12 ml, 6.67
mmol) and triphenylphosphine (0.142 g) and then heated to 60°C
for 4 h. The mixture was then cooled and concentrated to a
25 gum. Flash chromatography (10% methanol/dichloromethane)
yielded the requisite amine (190 mg, 84% yield) as a colorless
oil: $[\alpha]_{\text{D}}^{25}$ -1.7° (c 0.52, CHCl_3); IR (CH_2Cl_2) 3730 (w), 3045
(m), 2940 (m), 1610 (w), 1450 (m), 1426 (s), 1372 (m), 1271
(s), 1183 (s), 1180 (s), 1115 (s), 1091 (s), 1076 (s), 900
30 (s), 730 (br s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.97 (d, J =
8.3 Hz, 1 H), 7.83 (apparent d, J = 7.4 Hz, 2 H), 7.49-7.44
(m, 3 H), 7.37-7.14 (m, 17 H), 4.90 (d, J = 10.9 Hz, 1 H),
4.85 (d, J = 10.9 Hz, 1 H), 4.78 (d, J = 10.9 Hz, 1 H), 4.72
(d, J = 11.0 Hz, 1 H), 4.61 (d, J = 10.9 Hz, 1 H), 4.59 (d,
35 J = 11.0 Hz, 1 H), 4.41 (d, J = 7.8 Hz, 1 H), 4.20 (dt, J =
9.6, 6.9 Hz, 1 H), 3.83 (dt, J = 9.6, 7.2 Hz, 1 H), 3.67
(apparent t, J = 9.0 Hz, 2 H), 3.63-3.60 (m, 1 H), 3.58

- 117 -

(apparent t, $J = 9.0$ Hz, 2 H), 3.49 (dt, $J = 9.4, 6.5$ Hz, 1 H), 3.41 (t, $J = 6.7$ Hz, 2 H), 3.39-3.37 (m, 1 H), 3.00 (t, $J = 6.9$ Hz, 2 H), 2.99-2.97 (br, 2 H), 1.57-1.25 (m, 8 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.24, 133.60, 129.13, 128.41, 128.34, 128.27, 128.00, 127.85, 127.84, 127.56, 126.67, 124.74, 123.47, 123.12, 119.68, 113.71, 103.73, 84.65, 82.25, 77.95, 75.65, 74.97, 74.90, 74.74, 71.64, 69.65, 68.76, 29.55, 26.60, 25.88, 25.71.

A solution of the above amine (0.248 g, 0.30 mmol) in ethanol (22.5 ml) was treated with 5 M aqueous potassium hydroxide (4.5 ml) and heated to reflux. After 5 h the mixture was cooled, diluted with saturated aqueous ammonium chloride (30 ml), and poured into dichloromethane (30 ml). The aqueous phase was extracted with dichloromethane and the combined organic solutions were washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) furnished III-4c (179 mg, 87% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} +9.4^\circ$ (c 0.25, CHCl_3); IR (CH_2Cl_2) 3700 (br), 3026 (s), 2980 (s), 2925 (m), 2860 (m), 2085 (m), 1610 (w), 1440 (s), 1421 (s), 1365 (s), 1255 (s), 1175 (s), 1120 (s), 1085 (s), 1075 (s), 980 (w), 890 (s), 700 (br) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.49 (br s, 1 H), 7.49 (d, $J = 7.8$ Hz, 1 H), 7.26-7.15 (m, 16 H) 7.07 (t, $J = 8.0$ Hz, 1 H), 7.00 (t, $J = 7.1$ Hz, 1 H), 6.92 (s, 1 H), 4.84 (d, $J = 11.0$ Hz, 1 H), 4.77 (d, $J = 10.9$ Hz, 1 H), 4.76 (d, $J = 10.9$ Hz, 1 H), 4.70 (d, $J = 10.9$ Hz, 1 H), 4.59 (d, $J = 11.0$ Hz, 1 H), 4.49 (d, $J = 11.0$ Hz, 1 H), 4.38 (d, $J = 7.8$ Hz, 1 H), 4.08 (dt, $J = 9.3, 6.9$ Hz, 1 H), 3.77 (dt, $J = 9.3, 7.1$ Hz, 1 H), 3.62-3.28 (m, 8 H), 3.03 (t, $J = 7.3$ Hz, 2 H), 2.67 (t, $J = 7.5$ Hz, 2 H), 1.48-1.37 (m, 4 H), 1.17-1.13 (m, 4 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.54, 138.48, 136.15, 136.11, 128.39, 128.32, 128.27, 128.03, 127.84, 127.75, 127.54, 127.47, 122.20, 121.79, 119.12, 118.64, 112.16, 111.23, 103.68, 84.65, 82.29, 78.09, 75.62, 74.91, 74.83, 74.68, 71.37, 70.26, 69.77, 39.74, 29.35, 27.37, 26.13, 25.83, 25.42.

- 118 -

M. 5-Trifluoroacetamido-1-pentanol (III-18a).

A solution of 5-amino-1-pentanol (1.00 g, 9.69 mmol) in methanol (8 ml) was cooled to 0°C and treated with triethylamine (3.28 ml, 23.5 mmol), followed by dropwise addition of trifluoroacetic anhydride (1.88 ml, 13.4 mmol). The reaction mixture was stirred at room temperature for 16 h. Concentration and flash chromatography (60% ethyl acetate/petroleum ether) then furnished III-18a (1.7 g, 89% yield) as an oil: IR (film) 3300 (s), 3100 (m), 2950 (s), 2875 (m), 1705 (s), 1563 (m), 1450 (w), 1375 (w), 1345 (w), 1210 (s), 1185 (s), 1160 (s), 1075 (w), 1055 (m), 1028 (w), 1003 (w), 970 (w), 875 (w), 720 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 6.72 (s, 1 H), 3.66 (m, 2 H), 3.37 (dd, $J = 13.3, 6.8$ Hz, 2 H), 1.77 (s, 1 H), 1.66-1.58 (m, 4 H), 1.47-1.41 (m, 2 H); high resolution mass spectrum (Cl , CH_4) m/z 200.0901 [$(\text{M}+\text{H})^+$; calcd for $\text{C}_7\text{H}_{13}\text{F}_3\text{NO}_2$: 200.0696].

N. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-amino-6-deoxy-6-N-(5-hydroxypentyl)- β -D-glucopyranoside (III-4e).

A stirred solution of III-17 (196 mg, 0.27 mmol) in dry dichloromethane (2.7 ml) was cooled to -78°C and treated with 2,6-di-tert-butyl-4-methylpyridine (880 mg, 0.427 mmol) followed by triflic anhydride (58 ml, 0.347 mmol). The mixture was stirred for 15 min at -78°C, warmed to room temperature over 20 min, and then poured into saturated aqueous sodium bicarbonate (20 ml) and extracted with ethyl acetate (60 ml). The organic layer was washed with saturated aqueous sodium bicarbonate (3 x 20 ml) and brine (20 ml) and dried over magnesium sulfate. Filtration and concentration in vacuo provided crude triflate which was used without purification.

A solution of 5-trifluoroacetamido-1-pentanol (III-18a) (265 mg, 1.3 mmol) in THF (10 ml) was added to a stirred suspension of sodium hydride (123 mg, 3.07 mmol, 60% oil dispersion) in THF (17 ml) at 0°C. After 10 min the suspension was warmed to room temperature, stirred for 1 h, and re-cooled to 0°C and a solution of the above triflate

- 119 -

(0.574 mmol) in dichloromethane (25 ml) was added dropwise. The reaction was stirred at 0°C for 30 min and then at room temperature for 24 h, cooled to 0°C, quenched with saturated aqueous ammonium chloride (10 ml), and extracted with ethyl acetate (2 x 150 ml). The combined extracts were washed with water (50 ml) and brine (50 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (2% methanol/dichloromethane) afforded an inseparable mixture of compounds, presumably III-19d and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (6 ml) was treated with 5 M aqueous NaOH (2 ml, 10 mmol) and the reaction mixture was heated to reflux for 2 h, cooled, and concentrated in vacuo. The residue was dissolved in ethyl acetate (40 ml) and the solution was washed with water (15 ml) and brine (15 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (5% methanol/dichloromethane) afforded III-4e (150 mg, 83% yield for 3 steps) as a pale yellow oil: $[\alpha]_D^{25} +3.2^\circ$ (c 0.31, acetonitrile); UV (1.14×10^{-4} M, acetonitrile) λ_{\max} 289.6 (ϵ 4.17×10^3), 280.8 (4.97×10^3), 220.0 (2.4×10^4) nm; IR (film) 3420 (w), 3300 (w), 3063 (w), 3033 (w), 2938 (m), 2860 (m), 1495 (w), 1455 (m), 1360 (m), 1210 (w), 1072 (s), 1026 (m), 910 (w), 538 (s), 495 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.98 (s, 1 H), 7.59 (d, $J = 7.9$ Hz, 1 H), 7.33-7.04 (m, 1.9 H), 4.90 (d, $J = 10.9$ Hz, 1 H), 4.85 (d, $J = 11.1$ Hz, 1 H), 4.80 (d, $J = 11.0$ Hz, 1 H), 4.77 (d, $J = 10.9$ Hz, 1 H), 4.64 (d, $J = 11.0$ Hz, 1 H), 4.60 (d, $J = 11.1$ Hz, 1 H), 4.48 (d, $J = 7.8$ Hz, 1 H), 4.21 (ddd, $J = 9.4, 6.7, 6.7$ Hz, 1 H), 3.89 (ddd, $J = 9.4, 7.3, 7.3$ Hz, 1 H), 3.64 (dd, $J = 9.0, 9.0$ Hz, 1 H), 3.56 (t, $J = 6.4$ Hz, 2 H), 3.51-3.47 (m, 1 H), 3.42 (t, $J = 9.2$ Hz, 2 H), 3.11 (t, $J = 7.0$ Hz, 2 H), 2.96 (dd, $J = 12.3, 2.6$ Hz, 1 H), 2.66 (dd, $J = 12.3, 7.8$ Hz, 1 H), 2.62-2.54 (m, 2 H), 1.93 (s, 2 H), 1.54-1.44 (m, 4 H), 1.38-1.32 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.57, 138.49, 138.14, 136.17, 128.43, 128.36, 128.29, 128.02, 127.88,

- 120 -

127.82, 127.60, 127.56, 127.50, 122.14, 121.96, 119.30, 118.68, 112.60, 111.13, 103.67, 84.61, 82.45, 79.70, 77.20, 75.68, 74.99, 74.73, 73.82, 70.25, 62.63, 50.52, 49.59, 32.36, 29.28, 25.86, 23.31; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 679.3700 $[(M+H)^+]$; calcd for $C_{42}H_{50}N_2O_6$: 679.3747].

O. 4-Trifluoroacetamido-1-butanol (III-18b).

Trifluoroacetylation of 4-amino-1-butanol (0.700 g, 7.86 mmol) as described for III-18a followed by flash chromatography (55% ethyl acetate/hexane) afforded III-18b (1.32 g, 85% yield) as an oil: IR (film) 3310 (s), 3100 (m), 2950 (m), 2890 (m), 1710 (s), 1568 (m), 1450 (w), 1380 (w), 1348 (w), 1215 (s), 1186 (s), 1160 (s), 1073 (m), 1053 (m), 1028 (w), 900 (w), 880 (w), 857 (w), 723 (m) cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 7.28 (s, 1 H), 3.72 (dd, J = 10.2, 5.8 Hz, 2 H), 3.40 (dd, J = 12.6, 6.3 Hz, 2 H), 1.99 (t, J = 4.2 Hz, 1 H), 1.78-1.70 (m, 2 H), 1.68-1.62 (m, 2 H); high resolution mass spectrum (Cl , CH_4) m/z 186.0732 $[(M+H)^+]$; calcd for $C_6H_{11}F_3NO_2$: 186.0742].

20 P. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-amino-6-deoxy-6-N-(4-hydroxybutyl)- β -D-glucopyranoside (III-4f).

A solution of 4-trifluoroacetamido-1-butanol (III-18b) (425 mg, 2.29 mmol) in THF (10 ml) was added to a stirred suspension of sodium hydride (60% dispersion in oil, 210 mg, 5.27 mmol) in THF (28 ml) at 0°C. After 10 min the suspension was warmed to room temperature, stirred for 1 h, and recooled to 0°C. Crude triflate (0.27 mmol), prepared as described for III-4e, was dissolved in dichloromethane (16 ml) and added dropwise. The reaction was stirred at 0°C for 1 h and then at room temperature for 24 h, cooled to 0°C, quenched with saturated aqueous ammonium chloride (10 ml), and extracted with ethyl acetate (2 x 150 ml). The combined extracts were washed with water (50 ml) and brine (50 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (3% methanol/dichloromethane) afforded an inseparable mixture of compounds, presumably III-

- 121 -

19e and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (11 ml) was treated with 2.5 M aqueous NaOH (7.0 ml, 17.5 mmol) and the reaction mixture was heated to reflux for 2 h, cooled to room temperature, and concentrated in vacuo. The residue was taken up in dichloromethane (60 ml) and the solution was washed with brine (20 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (5% methanol/dichloromethane) provided III-4f (148 mg, 39%) as a pale yellow oil: IR (film) 3435 (w), 3310 (w), 2930 (m), 2870 (m), 1502 (w), 1460 (m), 1364 (m), 1215 (w), 1075 (s), 1032 (sh), 1012 (sh), 913 (m), 815 (w), 740 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.98 (s, 1 H), 7.59 (d, J = 7.9 Hz, 1 H), 7.33-7.21 (m, 1.5 H), 7.19-7.16 (m, 2 H), 7.12-7.09 (m, 1 H), 7.04 (d, J = 2.1 Hz, 1 H), 4.90 (d, J = 10.9 Hz, 1 H), 4.86 (d, J = 11.1 Hz, 1 H), 4.78 (d, J = 11.1 Hz, 1 H), 4.76 (d, J = 10.9 Hz, 1 H), 4.63 (d, J = 11.0 Hz, 1 H), 4.58 (d, J = 11.1 Hz, 1 H), 4.46 (d, J = 7.8 Hz, 1 H), 4.20 (ddd, J = 9.5, 6.7, 6.7 Hz, 1 H), 3.89 (ddd, J = 9.5, 7.3, 7.3 Hz, 1 H), 3.62 (apparent t, J = 9.0 Hz, 1 H), 3.53 (t, J = 5.3 Hz, 2 H), 3.46 (ddd, J = 9.5, 4.4, 2.9 Hz, 1 H), 3.41 (dd, J = 9.1, 7.9 Hz, 1 H), 3.36 (apparent t, J = 9.2 Hz, 1 H), 3.11 (t, J = 6.9 Hz, 2 H), 2.93 (dd, J = 12.3, 2.9 Hz, 1 H), 2.63 (dd, J = 12.3, 7.9 Hz, 1 H), 2.59 (t, J = 5.7 Hz, 2 H), 1.61 (m, 2 H), 1.55 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.50, 138.46, 138.07, 136.15, 128.43, 128.35, 128.28, 127.98, 127.87, 127.82, 127.59, 127.53, 127.46, 122.15, 121.95, 119.29, 118.67, 112.60, 111.14, 103.61, 84.58, 82.38, 79.73, 75.66, 74.97, 74.69, 73.36, 70.20, 62.54, 50.32, 49.49, 32.11, 28.10, 25.85; high resolution mass spectrum (Cl , NH_3) m/z 665.3640 $[(\text{M}+\text{H})^+]$; calcd for $\text{C}_{41}\text{H}_{49}\text{N}_2\text{O}_6$: 665.3590].

- 122 -

Q. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl-
6-amino-6-deoxy-6-N-(6-hydroxyhexyl)- β -D-glucopyranoside (III-4g).

A solution of 6-trifluoroacetamido-1-hexanol (III-18c) (145.0 mg, 0.680 mmol) in THF (2 ml) was added to a suspension of sodium hydride (60.0 mg, 1.50 mmol, 60% dispersion in oil) in THF (2 ml) at 0°C. The mixture was stirred at room temperature for 1.5 h, cooled to 0°C, and treated with a solution of the triflate derived from III-17 (0.136 mmol), prepared as described for the synthesis of III-4e, in dichloromethane (4 ml). The reaction mixture was then stirred at room temperature for 48 h, cooled to 0°C, quenched with saturated aqueous ammonium chloride (10 ml), and extracted with ethyl acetate (3 x 10 ml). The combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (5% methanol/dichloromethane) afforded an inseparable mixture of compounds, presumably III-19f and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (6 ml) was treated with 5 N aqueous sodium hydroxide (2 ml) and heated to reflux for 2 h. Cooling followed by concentration in vacuo gave an oily residue which was taken up in water (5 ml) and extracted with dichloromethane (3 x 5 ml). The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (6% methanol/dichloromethane) furnished III-4g as a colorless oil (36.4 mg, 54% yield): $[\alpha]_D^{25} -18^\circ$ (c 0.18, acetonitrile); UV (1.72 x 10⁻⁴ M, acetonitrile) λ_{\max} 290.0 (ϵ 1.02 x 10³), 281.2 (1.13 x 10³), 228.4 (1.39 x 10³) nm; IR (film) 3440 (m), 3310 (m), 3060 (m), 3030 (m), 2930 (s), 2860 (s), 2240 (w), 1497 (w), 1455 (s), 1360 (m), 1210 (w), 1070 (s), 910 (s), 740 (s), 700 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.17 (br s, 1 H), 7.59 (d, J = 7.9 Hz, 1 H), 7.33-7.00 (m, 19 H), 4.91 (d, J = 10.9 Hz, 1 H), 4.86 (d, J = 11.1 Hz, 1 H), 4.80 (d, J = 11.3 Hz, 1 H), 4.78 (d, J = 11.1 Hz, 1 H), 4.65 (d, J = 11.0 Hz, 1 H), 4.60 (d, J = 11.1 Hz, 1 H), 4.47 (d, J = 7.8 Hz, 1 H), 4.21

- 123 -

(dt, $J = 9.4, 6.8$ Hz, 1 H), 3.86 (dt, $J = 9.4, 7.6$ Hz, 1 H), 3.64 (t, $J = 9.0$ Hz, 1 H), 3.55 (t, $J = 6.6$ Hz, 2 H), 3.51-3.40 (m, 3 H), 3.12 (t, $J = 7.2$ Hz, 2 H), 2.96-2.13 (dd, $J = 12.2, 2.6$ Hz, 1 H), 2.68-2.51 (m, 3 H), 1.87 (br s, 2 H), 1.51-1.41 (m, 4 H), 1.33-1.25 (m, 4 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.47, 138.39, 138.05, 136.11, 128.39, 128.34, 128.27, 128.02, 127.96, 127.88, 127.80, 127.59, 127.55, 127.40, 122.10, 121.87, 119.21, 118.62, 112.32, 111.13, 103.61, 84.55, 82.38, 79.77, 75.69, 75.00, 74.72, 73.91, 70.25, 62.67, 50.64, 49.61, 32.55, 29.78, 26.97, 25.81, 25.55; high resolution mass spectrum (Cl , CH_4) m/z 693.3946 (M^+ ; calcd for $\text{C}_{43}\text{H}_{50}\text{N}_2\text{O}_6$: 693.3903).

R. 5-Acetamido-1-pentanol (III-20).

A solution of 5-amino-1-pentanol (0.650 g, 6.31 mmol) in methanol (15 ml) was cooled to 0°C and treated with triethylamine (1.62 ml, 11.6 mmol) followed by acetic anhydride (0.891 ml, 9.45 mmol). The reaction mixture was stirred at room temperature overnight. TLC analysis (8% methanol/dichloromethane) then revealed some unreacted material, so additional triethylamine (1.6 ml, 11.6 mmol) and acetic anhydride (0.9 ml, 9.5 mmol) were added at room temperature and the solution was stirred 16 h further. Concentration in vacuo and flash chromatography (7% methanol/dichloromethane) afforded III-20 (1 g, 94% yield) as a pale yellow oil: IR (film) 3300 (s), 3100 (m), 2940 (s), 2870 (m), 1650 (s), 1560 (s), 1439 (m), 1372 (m), 1295 (m), 1220 (w), 1180 (w), 1050 (m), 1010 (w) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 6.21 (s, 1 H), 3.62 (t, $J = 6.4$ Hz, 2 H), 3.23 (dd, $J = 12.9, 7.0$ Hz, 2 H), 2.87 (s, 1 H), 1.97 (s, 3 H), 1.60-1.50 (m, 4 H), 1.43-1.37 (m, 2 H); high resolution mass spectrum (Cl , CH_4) m/z 146.1164 [$(\text{M}+\text{H})^+$; calcd for $\text{C}_7\text{H}_{16}\text{NO}_2$: 146.1181].

S. 2-(N-Phenylsulfonylindol-3-yl)ethyl 2,3,4-Tri-O-benzyl-6-O-(5-acetamidopentyl)- β -D-glucopyranoside (III-4d).

A solution of 5-acetamido-1-pentanol (177 mg, 1.22 mmol) in THF (8 ml) was added to a stirred suspension of

- 124 -

sodium hydride (60% dispersion in oil, 108 mg, 2.70 mmol) in THF (20 ml) at 0°C. After 10 min the mixture was stirred at room temperature for 1.5 h and cooled to 0°C. The triflate derived from III-17 (0.245 mmol), prepared as described for the synthesis of III-4a, was dissolved in dichloromethane (20 ml) and slowly added dropwise. The reaction was stirred at 0°C for 1 h and at room temperature for 24 h, and then was cooled to 0°C, quenched with saturated aqueous ammonium chloride (10 ml) and diluted with ethyl acetate (150 ml). The organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (3% methanol/dichloromethane) afforded an inseparable mixture of compounds, presumably III-21 and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (4 ml) was treated with 5 N aqueous NaOH (2 ml, 10 mmol) and then heated to reflux for 2 h, cooled, and concentrated in vacuo. The residue was dissolved in ethyl acetate (40 ml) and the resultant solution was washed with water and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (4% methanol/dichloromethane) provided III-4d (88 mg, 50% yield) as a colorless oil: $[\alpha]_D^{25} +14.5^\circ$ (c 0.53, CHCl₃); IR (film) 3300 (s), 3090 (w), 3065 (m), 3035 (m), 2940 (s), 2870 (s), 1960 (w), 1885 (w), 1815 (w), 1662 (s), 1550 (m), 1500 (m), 1458 (s), 1369 (s), 1285 (m), 1213 (m), 1070 (s), 914 (w), 810 (w), 742 (s), 700 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.26 (s, 1 H), 7.59 (d, J = 8.1 Hz, 1 H), 7.34-7.21 (m, 1.6 H), 7.19-7.16 (m, 1 H), 7.12-7.08 (m, 1 H), 7.03 (d, J = 2.2 Hz, 1 H), 5.41 (s, 1 H), 4.92 (d, J = 10.9 Hz, 1 H), 4.85 (d, J = 11.0 Hz, 1 H), 4.83 (d, J = 11.0 Hz, 1 H), 4.78 (d, J = 11.0 Hz, 1 H), 4.66 (d, J = 11.0 Hz, 1 H), 4.59 (d, J = 11.0 Hz, 1 H), 4.45 (d, J = 7.8 Hz, 1 H), 4.22 (ddd, J = 9.4, 6.9, 6.9 Hz, 1 H), 3.86 (ddd, J = 9.4, 7.5, 7.5 Hz, 1 H), 3.68 (dd, J = 10.9, 1.8 Hz, 1 H), 3.64 (apparent t, J = 9.0 Hz, 1 H), 3.59 (dd, J = 10.9, 5.1 Hz, 1 H), 3.55 (apparent t, J = 9.0 Hz, 1 H), 3.51-3.39 (m, 4 H), 3.17-3.13 (m, 2 H), 3.12

- 125 -

(t, $J = 7.2$ Hz, 2 H), 1.91 (s, 3 H), 1.58-1.53 (m, 2 H), 1.48-1.42 (m, 2 H), 1.38-1.32 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.06, 138.57, 138.22, 136.21, 128.41, 128.35, 128.27, 128.03, 127.87, 127.84, 127.76, 127.59, 127.55, 127.49, 122.18, 121.81, 119.14, 118.61, 112.46, 111.19, 103.68, 84.68, 82.33, 78.04, 77.20, 75.67, 74.93, 74.83, 74.67, 71.42, 70.06, 69.71, 39.56, 29.29, 25.76, 23.61, 23.27; high resolution mass spectrum (Cl , NH_3) m/z 721.3790 [$(\text{M}+\text{H})^+$; calcd for $\text{C}_{44}\text{H}_{53}\text{N}_2\text{O}_7$: 721.3852].

10 **T. 1,2,4,6-Tetra-O-acetyl-3-deoxy- β -D-glucopyranoside (III-23).**

A solution of 3-deoxydiacetone-D-glucose (III-22) (27.5 g, 113 mmol) in 60% aqueous acetic acid (200 ml) was heated at 90°C for 1 h, cooled, and concentrated in vacuo, and the residue was azeotroped with dry benzene (4 x 20 ml). A solution of the concentrate in dry pyridine (250 ml) was treated with acetic anhydride (107 ml, 1.13 mol) and DMAP (2 mol%, 275 mg) and stirred at room temperature for 30 min. After concentration in vacuo the residue was diluted with water (40 ml) and extracted with dichloromethane (3 x 40 ml), and the combined extracts were then washed with brine (40 ml), dried over sodium sulfate, filtered, and concentrated in vacuo. Recrystallization from ether afforded the pure β -anomer (11.3 g) as a fine white powder. Concentration of the filtrate and flash chromatography (45% ethyl acetate/hexane) gave a mixture of α - and β -anomers as a colorless gum (23.0 g, total yield 91.7%). β -Anomer III-23: $[\alpha]_{\text{D}}^{25}$ 17.1° (c 1.05, CH_3OH); IR (CHCl_3) 3010 (m), 2940 (w), 2870 (w), 1745 (s), 1510 (w), 1365 (m), 1230 (s), 1210 (s), 1030 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 5.67 (d, $J = 7.9$ Hz, 1 H), 4.89-4.81 (m, 2 H), 4.21 (dd, $J = 5.1, 12.3$ Hz, 1 H), 4.12 (dd, $J = 2.5, 12.2$ Hz, 1 H), 3.81-3.79 (m, 1 H), 2.60 (ddd, $J = 5.0, 5.0, 12.3$ Hz, 1 H), 2.10 (s, 3 H), 2.06 (s, 3 H), 2.03 (s, 3 H), 2.02 (s, 3 H), 1.64 (apparent q, $J = 11.0$, 1 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 170.69, 169.43, 169.31, 169.19, 93.06, 75.68, 67.33, 65.00, 62.07, 32.69, 20.92, 20.77; high resolution mass spectrum (Cl , NH_3) m/z 350.1412

- 126 -

[(M+NH₄)⁺; calcd for C₁₄H₂₀O₉Cl: 350.1450]. Anal. Calcd for C₁₄H₂₀O₉: C, 50.60; H, 6.07. Found: C, 50.65; H, 6.16.

U. 2 - (N-Phenylsulfonylindol-3-yl) ethyl
2,4,6-Tri-O-acetyl-3-deoxy-β-D-glucopyranoside
(III-24).

Hydrobromic acid (30% in acetic acid, 3 ml, 14.0 mmol) was added to III-23 (750 mg, 2.26 mmol) at 0°C. After 10 min, the solution was warmed to room temperature, stirred for 30 min, diluted with ether (20 ml), and poured into a mixture of ice and saturated aqueous sodium bicarbonate (25 ml). An additional 30 ml of ether was added and the layers were separated. The organic layer was washed with saturated aqueous sodium bicarbonate (3 x 25 ml), water, and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The crude bromide was used without purification in the next step: high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 370.0470 [(M+NH₄)⁺; calcd for C₁₂H₁₇BrO₃: 370.0494].

A solution of N-(benzenesulfonyl)tryptophol (III-12) (1.20 g, 4.0 mmol) in dry benzene (4 ml) was added to a stirred suspension of activated, powdered 4 Angstrom molecular sieves (1.33 g) in dry hexane (11 ml) at room temperature. A solution of the bromide (2.26 mmol) in dry benzene (4 ml) was introduced, followed by silver(I) oxide (523 mg, 2.26 mmol). The reaction vessel was covered with aluminum foil, and the mixture was stirred for 3 days and then filtered through Celite. Concentration and flash chromatography (10:1 dichloromethane/ether) provided pure III-24 (781 mg, 60% yield) as a white foam: mp 49-51°C; [α]_D²⁵ -12° (c 0.21, acetonitrile); UV (8.3 x 10⁻⁵ M, acetonitrile) λ_{max} 253.6 (ε 1.12 x 10⁴), 214.0 (2.43 x 10⁴) nm; IR (film) 3045 (w), 2970 (w), 2895 (w), 1745 (s), 1449 (m), 1370 (s), 1230 (s), 1167 (m), 1120 (w), 1083 (w), 1035 (m), 908 (w), 853 (w), 748 (s), 720 (w), 682 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.97 (d, J = 8.3 Hz, 1 H), 7.86-7.84 (m, 2 H), 7.53-7.41 (m, 5 H), 7.32-7.29 (m, 1 H), 7.25-7.22 (t, J = 7.6 Hz, 1 H), 4.84 (ddd, J = 10.7, 9.6, 4.9 Hz, 1 H), 4.77 (ddd, J = 12.8, 7.6, 5.2 Hz,

- 127 -

1 H), 4.49 (d, $J = 7.6$ Hz, 1 H), 4.24-4.14 (m, 3 H), 3.76 (ddd, $J = 9.4, 6.9, 6.9$ Hz, 1 H), 3.68 (ddd, $J = 9.2, 5.0, 3.0$ Hz, 1 H), 2.96 (t, $J = 7.1$ Hz, 2 H), 2.55 (ddd, $J = 12.2, 5.0, 3.0$ Hz, 1 H), 2.06 (s, 3 H), 2.04 (s, 3 H), 1.93 (s, 3 H), 1.56 (apparent q, $J = 11.5$ Hz, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.80, 169.47, 133.68, 131.06, 129.20, 126.72, 124.73, 123.56, 123.16, 119.84, 119.50, 113.66, 106.62, 102.09, 75.03, 68.46, 68.38, 65.83, 62.65, 32.92, 25.37, 20.87, 20.79; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.1623 (M^+ ; calcd for $\text{C}_{28}\text{H}_{31}\text{NO}_{10}\text{S}$: 573.1669).

V. 2-(N-Phenylsulfonylindol-3-yl)ethyl 3-Deoxy- β -D-glucopyranoside (III-25).

Sodium methoxide (55.2 mg, 1.02 mmol) was added to a suspension of III-24 (735 mg, 1.28 mmol) in methanol (6.4 ml). The mixture was stirred at room temperature for 90 min, diluted with methanol (6 ml), and neutralized with Amberlyst[®] 15 ion exchange resin. The resin was quickly filtered. Concentration in vacuo and flash chromatography (12:1:1 dichloromethane/acetone/methanol) afforded pure III-25 (498 mg, 87% yield) as a white solid: mp 55-57°C; $[\alpha]_{\text{D}}^{25} -26^\circ$ (c 0.25, methanol); UV (1.39×10^{-4} M, acetonitrile) λ_{max} 254.0 (ϵ 1.24×10^4), 216.0 (2.02×10^4) nm; IR (film) 3415 (s), 3070 (w), 3025 (w), 2945 (m), 2890 (m), 1605 (w), 1449 (s), 1366 (s), 1279 (w), 1215 (w), 1173 (s), 1125 (m), 1078 (s), 1028 (s), 975 (w), 741 (s), 720 (m), 681 (m) cm^{-1} ; ^1H NMR (500 MHz, CD_3OD) δ 7.96-7.94 (m, 1 H), 7.91-7.89 (m, 1 H), 7.61 (s, 1 H), 7.59-7.54 (m, 2 H), 7.49-7.45 (m, 2 H), 7.31-7.28 (m, 1 H), 7.24-7.21 (m, 1 H), 4.30 (d, $J = 7.6$ Hz, 1 H), 4.18 (ddd, $J = 9.6, 7.0, 7.0$ Hz, 1 H), 3.88-3.82 (m, 2 H), 3.66 (dd, $J = 11.8, 6.2$ Hz, 1 H), 3.50 (ddd, $J = 11.2, 9.4, 4.8$ Hz, 1 H), 3.40 (ddd, $J = 12.4, 7.6, 5.0$ Hz, 1 H), 3.31 (s, 2 H), 3.27 (ddd, $J = 9.2, 6.1, 2.5$ Hz, 1 H), 3.00 (t, $J = 6.8$ Hz, 2 H), 2.31 (ddd, $J = 12.2, 4.9, 4.9$ Hz, 1 H), 1.50 (apparent q, $J = 11.8$ Hz, 1 H); ^{13}C NMR (500 MHz, CD_3OD) δ 139.40, 136.57, 135.02, 132.62, 130.38, 127.89, 125.65, 125.30, 124.36, 121.74, 120.63, 114.70, 106.49, 81.82, 69.41, 69.37, 66.27, 62.95, 40.72, 26.32; high

- 128 -

resolution mass spectrum (CI, NH₃) m/z 465.1627 [(M+NH₄)⁺];
calcd for C₂₂H₂₅NO₇S: 465.1685].

W. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
3-Deoxy-6-O-tert-butyldiphenylsilyl- β -D-
glucopyranoside (III-26).

5 A stirred solution of III-25 (779 mg, 1.74 mmol) in dry
DMF (17 ml, 0.1 M) was treated with imidazole (260 mg, 3.83
mmol) followed by tert-butyldiphenylsilyl chloride (0.541 ml,
2.09 mmol). The solution was heated at 50°C for 24 h, cooled,
10 diluted with ethyl acetate (250 ml), and washed with water and
brine. The organic phase was dried over magnesium sulfate,
filtered, and concentrated in vacuo. Flash chromatography (3%
methanol/dichloromethane) provided pure III-26 (1.04 g, 87%
yield) as a white foam: $[\alpha]_D^{25}$ -24° (c 0.46, acetonitrile);
15 UV (1.68 x 10⁻⁴ M, acetonitrile) λ_{max} 254.0 (ϵ 1.11 x 10⁴),
220.4 (1.90 x 10⁴) nm; IR (film) 3430 (s), 3080 (w), 3060 (w),
3020 (w), 2940 (s), 2865 (s), 1668 (m), 1449 (s), 1428 (m),
1370 (s), 1275 (w), 1213 (w), 1112 (s), 1070 (s), 855 (w), 820
(w), 740 (s), 720 (w), 700 (m), 680 (w) cm⁻¹; ¹H NMR (500 MHz,
20 acetone-d₆) δ 7.98-7.96 (m, 1 H), 7.84-7.82 (m, 2 H),
7.68-7.65 (m, 4 H), 7.51-7.36 (m, 1 H), 7.31-7.28 (m, 1 H),
7.21-7.18 (m, 1 H), 4.19 (d, J = 7.4 Hz, 1 H), 4.09 (ddd, J
= 9.5, 6.2, 6.2 Hz, 1 H), 3.92 (dd, J = 10.3, 5.0 Hz, 1 H),
3.84 (dd, J = 10.4, 7.3 Hz, 1 H), 3.82-3.77 (m, 1 H), 3.68
25 (ddd, J = 9.5, 7.1, 7.1 Hz, 1 H), 3.48-3.40 (m, 2 H), 3.29 (d,
 J = 2.3 Hz, 1 H), 2.97-2.89 (m, 2 H), 2.37 (ddd, J = 12.4,
4.8, 4.8 Hz, 1 H), 2.10 (d, J = 2.5 Hz, 1 H), 1.53 (apparent
q, J = 11.5 Hz, 1 H), 1.06 (s, 9 H); ¹³C NMR (125 MHz,
acetone-d₆) δ 138.24, 135.54, 135.51, 135.14, 133.65, 132.46,
30 132.38, 130.97, 130.00, 129.17, 128.30, 127.86, 126.65,
124.79, 123.42, 123.13, 119.67, 119.34, 113.73, 104.73, 77.34,
68.83, 68.58, 68.28, 66.11, 37.34, 26.77, 25.45, 19.09; high
resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z
686.2651 [(M+H)⁺; calcd for C₃₈H₄₃NO₇SSi: 686.2607]. Anal. Calcd
35 for C₃₈H₄₃O₇NSSi: C, 66.54; H, 6.32. Found: C, 66.18; H,
6.14.

- 129 -

X. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
3-Deoxy-2,4-di-O-benzyl-6-O-tert-butyldiphenyl-
silyl- β -D-glucopyranoside (III-27).

A stirred suspension of sodium hydride (4.63 mmol, 185
5 mg, 60% oil dispersion) in THF (5 ml) was cooled to 0°C and
a solution of III-26 (1.27 g, 1.85 mmol) in THF (10 ml) was
added. After 10 min the reaction mixture was warmed to room
temperature, stirred for 1 h, re-cooled to 0°C and treated with
benzyl bromide (5.55 mmol, 0.660 ml) followed by
10 tetrabutylammonium iodide (68 mg, 0.185 mmol). The reaction
was then warmed to room temperature, stirred for 3 days, and
quenched with saturated aqueous ammonium chloride (3 ml) at
0°C. The mixture was diluted with ether (80 ml), washed with
water (2 x 30 ml) and brine (30 ml), dried over magnesium
15 sulfate, filtered, and concentrated in vacuo. Flash
chromatography (25% ether/petroleum ether) furnished pure III-
27 (760 mg, 47% yield) as a white foam: $[\alpha]_D^{25}$ -2.7° (c 0.66,
acetonitrile); UV (1.9×10^{-4} M, acetonitrile) λ_{\max} 254.0 (ϵ
1.19 x 10⁴), 220.8 (1.71 x 10⁴) nm; IR (film) 3080 (m), 3040
20 (m), 2945 (s), 2870 (s), 1585 (w), 1494 (w), 1445 (s), 1425
(m), 1369 (s), 1330 (w), 1307 (w), 1275 (m), 1205 (m), 1171
(s), 1109 (s), 1100 (s), 1025 (s), 972 (m), 935 (w), 905 (w),
849 (w), 817 (m), 739 (s), 695 (s) cm⁻¹; ¹H NMR (500 MHz,
CDCl₃) δ 7.99-7.97 (m, 1 H), 7.83-7.80 (m, 2 H), 7.71-7.67 (m,
25 4 H), 7.51-7.18 (m, 2-3 H), 4.70 (d, J = 12.0 Hz, 1 H), 4.59
(d, J = 11.4 Hz, 1 H), 4.56 (d, J = 12.0 Hz, 1 H), 4.44 (d,
 J = 11.5 Hz, 1 H), 4.42 (d, J = 7.5 Hz, 1 H), 4.19 (ddd, J =
9.6, 6.7, 6.7 Hz, 1 H), 3.95 (dd, J = 11.2, 1.9 Hz, 1 H), 3.88
(dd, J = 11.2, 5.0 Hz, 1 H), 3.80 (ddd, J = 9.6, 7.3, 7.3 Hz,
30 1 H), 3.55 (ddd, J = 11.0, 9.4, 4.6 Hz, 1 H), 3.41 (ddd, J =
9.2, 4.9, 1.8 Hz, 1 H), 3.32 (m, 1 H), 3.04 (t, J = 7.2 Hz,
2 H), 2.52 (ddd, J = 12.3, 4.9, 4.9 Hz, 1 H), 1.55 (apparent
q, J = 11.6 Hz, 1 H), 1.03 (s, 9 H); ¹³C NMR (125 MHz, CDCl₃)
 δ 138.68, 138.32, 138.08, 135.72, 135.56, 135.18, 133.74,
35 133.54, 133.49, 131.06, 129.52, 129.10, 128.36, 128.30,
127.66, 127.63, 127.59, 127.51, 127.45, 126.63, 124.69,
123.47, 123.12, 119.94, 119.57, 113.67, 105.11, 79.10, 75.27,

- 130 -

72.68, 72.06, 71.37, 68.18, 63.23, 34.99, 26.77, 25.80, 19.29; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 865.3419 (M^+ ; calcd for $C_{22}H_{25}NO_7Si$: 865.3468).

Y. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
5 3-Deoxy-2,4-di-O-benzyl-b-D-glucopyranoside (III-28).

Tetrabutylammonium fluoride (1.0 M in THF, 1.17 mmol, 1.17 ml) was added to a stirred solution of III-27 (675 mg, 0.780 mmol) in THF (10 ml). The solution was stirred for 2
10 h, diluted with ethyl acetate, washed with water and brine, and dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (60% ether/petroleum ether) afforded pure III-28 (445 mg, 91% yield) as a pale yellow oil:
15 $[\alpha]_D^{25} +2.5^\circ$ (c 0.44, acetonitrile); UV (9.97×10^{-5} M, acetonitrile) λ_{max} 254.0 (ϵ 1.06×10^4), 210.0 (2.88×10^4) nm; IR (film) 3485 (m), 3080 (w), 3045 (w), 2945 (m), 2890 (m), 1603 (w), 1484 (w), 1447 (s), 1369 (s), 1277 (w), 1206 (w), 1173 (s), 1118 (m), 1082 (s), 1039 (m), 1025 (m), 948 (w), 900 (w), 745 (s), 717 (m), 693 (m), 678 (m) cm^{-1} ; 1H NMR
20 (500 MHz, $CDCl_3$) δ 7.97 (d, $J = 8.3$ Hz, 1 H), 7.84-7.82 (m, 2 H), 7.53 (s, 1 H), 7.49-7.44 (m, 2 H), 7.37-7.21 (m, 1.4 H), 4.67 (d, $J = 12.0$ Hz, 1 H), 4.60 (d, $J = 11.4$ Hz, 1 H), 4.54 (d, $J = 12.0$ Hz, 1 H), 4.47 (d, $J = 11.6$ Hz, 1 H), 4.45 (d, $J = 7.5$ Hz, 1 H), 4.19 (ddd, $J = 9.5, 6.8, 6.8$ Hz, 1 H),
25 3.89-3.84 (m, 2 H), 3.73 (dd, $J = 11.9, 4.6$ Hz, 1 H), 3.47 (ddd, $J = 11.0, 9.3, 4.6$ Hz, 1 H), 3.39 (ddd, $J = 9.1, 4.5, 3.1$ Hz, 1 H), 3.26 (ddd, $J = 11.7, 9.2, 5.1$ Hz, 1 H), 2.99 (m, 2 H), 2.51 (ddd, $J = 12.3, 4.8, 4.8$ Hz, 1 H), 1.89 (s, 1 H), 1.55 (dd, $J = 23.4, 11.7$ Hz, 1 H); ^{13}C NMR (125 MHz, $CDCl_3$) δ
30 138.44, 138.30, 137.82, 135.15, 133.63, 131.03, 129.15, 128.49, 128.35, 127.89, 127.79, 127.63, 127.58, 126.68, 124.75, 123.65, 123.15, 119.80, 119.38, 113.73, 105.19, 78.18, 75.02, 72.71, 72.23, 71.29, 68.38, 62.38, 34.83, 25.61; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z
35 627.2370 (M^+ ; calcd for $C_{36}H_{37}NO_7S$: 627.2291).

- 131 -

2. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,4-Di-O-benzyl-3-deoxy-6-O-(5-azidopentyl)- β -D-
glucopyranoside (III-29a).

A stirred solution of 5-bromo-1-pentanol (0.79 g, 4.7
5 mmol) in DMSO (15 ml) was treated with sodium azide (1.83 g,
28.2 mmol). The resultant mixture was stirred at room
temperature for 2.5 h, diluted with water, and extracted with
diethyl ether. The combined organic layers were washed with
saturated aqueous sodium bicarbonate and brine, dried over
10 magnesium sulfate, filtered, and concentrated in vacuo. The
azide was used without purification in the next step.

Crude 5-azido-1-pentanol (280 mg, equivalent to 2.17
mmol) and 2,6-di-tert-butyl-4-methylpyridine (441 mg, 2.17
mmol) were dissolved in dichloromethane (9 ml) and triflic
15 anhydride (0.36 ml, 2.17 mmol) was added dropwise. After 10
min the mixture was poured into brine (40 ml) and extracted
with dichloromethane (2 x 40 ml). The combined organic layers
were dried over magnesium sulfate, filtered, and concentrated
in vacuo. The triflate was used without purification in the
20 next step.

Sodium hydride (16 mg, 0.40 mmol, 60% dispersion in
oil) was added to a solution of alcohol III-28 (120 mg, 0.198
mmol) and azido triflate (105 mg, equivalent to 0.40 mmol) in
dichloromethane (3 ml) at room temperature. The mixture was
25 stirred for 24 h, diluted with dichloromethane (40 ml) and
poured into saturated ammonium chloride (40 ml). The aqueous
phase was extracted with dichloromethane and the combined
organic solutions were washed with brine, dried over magnesium
sulfate, filtered, and concentrated in vacuo. Flash
30 chromatography (15% ethyl acetate/hexane) afforded III-29a
(121 mg, 83% yield) as a colorless oil: $[\alpha]_D^{25} +4.0^\circ$ (c 0.24,
CHCl₃); IR (CHCl₃) 3022 (s), 2940 (s), 2880 (m), 2105 (s),
1455 (s), 1375 (s), 1270 (s), 1210 (m), 1180 (m), 1125 (m),
1090 (m), 725 (s), 599 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.96
35 (d, J = 8.1 Hz, 1 H), 7.82 (dd, J = 8.2, 0.9 Hz, 2 H),
7.50-7.43 (m, 3 H), 7.29-7.19 (m, 14 H), 4.65 (d, J = 12.0 Hz,
1 H), 4.58 (d, J = 11.4 Hz, 1 H), 4.52 (d, J = 12.0 Hz, 1 H),

- 132 -

4.42 (d, $J = 11.5$ Hz, 1 H), 4.18 (dt, $J = 9.5, 6.7$ Hz, 1 H),
3.81 (dt, $J = 9.5, 7.1$ Hz, 1 H), 3.71 (d, $J = 10.6$ Hz, 1 H),
3.57 (dd, $J = 10.8, 4.7$ Hz, 1 H), 3.51-3.38 (m, 4 H),
3.31-3.21 (m, 1 H), 3.16 (t, $J = 6.9$ Hz, 2 H), 3.00 (t, $J =$
5 6.9 Hz, 2 H), 2.50-2.46 (dt, $J = 12.1, 4.5$ Hz, 1 H), 1.63-1.50
(m, 5 H), 1.48-1.32 (m, 3 H); ^{13}C NMR (62.5 MHz, CDCl_3) δ
138.52, 138.23, 137.00, 135.07, 133.59, 131.09, 129.14,
128.43, 128.31, 127.78, 127.68, 127.50, 126.70, 126.69,
124.70, 123.54, 123.09, 119.71, 119.48, 113.70, 105.26, 78.01,
10 74.92, 72.67, 72.25, 71.38, 71.24, 69.96, 68.41, 34.97, 29.62,
29.15, 28.66, 25.65, 23.39; high resolution mass spectrum
(FAB, *m*-nitrobenzyl alcohol) m/z 761.2973 (M^+ ; calcd for
 $\text{C}_{41}\text{H}_{46}\text{N}_4\text{O}_7\text{S}$: 761.2985).

AA. 2-(1*H*-Indol-3-yl)ethyl 2,4-Di-O-benzyl-
15 3-deoxy-6-O-(5-aminopentyl)- β -D-glucopyranoside
(III-5a).

A stirred solution of azide III-29a (80 mg, 0.109 mmol)
in THF (5.2 ml) and water (0.083 ml) was treated with
triphenylphosphine (65 mg, 0.248 mmol), heated at reflux for
20 2.5 h, cooled, and concentrated in vacuo. Flash
chromatography (10% methanol/dichloromethane) furnished the
corresponding amine (70 mg, 90% yield) as a colorless oil:
IR (CHCl_3) 3028 (m), 2940 (s), 2875 (m), 1450 (s), 1370 (s),
1280 (w), 1178 (s), 1122 (m), 1070 (m), 695 (w), 597 (w) cm^{-1} ;
25 ^1H NMR (500 MHz, CDCl_3) δ xxx; ^{13}C NMR (125 MHz, CDCl_3) δ
138.51, 138.25, 138.00, 135.13, 133.58, 131.05, 129.11,
128.40, 128.27, 127.76, 127.69, 127.62, 127.46, 126.65,
124.68, 123.54, 123.09, 119.91, 119.48, 133.66, 105.21, 77.97,
74.96, 72.64, 72.18, 71.34, 71.21, 69.94, 68.39, 39.70, 34.94,
30 28.89, 25.59, 23.44, 23.26; high resolution mass spectrum
(FAB, *m*-nitrobenzyl alcohol) m/z 713.3241 (M^+ ; calcd for
 $\text{C}_{41}\text{H}_{48}\text{N}_2\text{O}_7\text{S}$: 713.3260).

The above amine (14 mg, 0.020 mmol) was dissolved in
ethanol (2.2 ml) and treated with 5 M aqueous sodium hydroxide
35 (0.36 ml). The resultant mixture was heated at reflux for 3
h, cooled, diluted with brine, and poured into
dichloromethane. The aqueous layer was extracted with

- 133 -

dichloromethane (2 x 40 ml) and the combined organic solutions were dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) afforded III-5a (7 mg, 61% yield) as a colorless oil: $[\alpha]_D^{25}$ -12° (c 0.11, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 9.05 (br s, 1 H), 7.58 (d, J = 7.8 Hz, 1 H), 7.34-7.25 (m, 11 H), 7.14 (t, J = 7.5 Hz, 1 H), 7.07 (t, J = 7.5 Hz, 1 H), 7.04 (s, 1 H), 4.77 (d, J = 11.8 Hz, 1 H), 4.60 (d, J = 12.0 Hz, 1 H), 4.57 (d, J = 11.6 Hz, 1 H), 4.44 (d, J = 7.5 Hz, 1 H), 4.39 (d, J = 11.5 Hz, 1 H), 4.16 (dt, J = 9.3, 7.3 Hz, 1 H), 3.85 (dt, J = 9.3, 7.2 Hz, 1 H), 3.70 (d, J = 10.4 Hz, 1 H), 3.51 (dd, J = 10.6, 5.8 Hz, 1 H), 3.46-3.36 (m, 4 H), 3.35-3.29 (m, 1 H), 3.11 (t, J = 7.2 Hz, 2 H), 2.68 (br t, J = 7.1 Hz, 2 H), 2.53-2.49 (dt, J = 12.3, 4.7 Hz, 1 H), 1.56-1.42 (m, 5 H), 1.36-1.25 (m, 4 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.53, 137.86, 136.12, 128.45, 128.38, 127.86, 127.82, 127.72, 127.62, 127.12, 123.06, 122.02, 119.32, 118.62, 112.26, 111.63, 105.43, 77.49, 75.28, 72.79, 71.34, 71.19, 71.05, 70.39, 68.85, 39.21, 34.65, 27.54, 26.16, 25.72, 22.51; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.3313 (M⁺; calcd for C₃₅H₄₄N₂O₅: 573.3328).

**AB. 2-(N-Phenylsulfonylindol-3-yl)ethyl
2,4-Di-O-benzyl-3-deoxy-6-O-(6-azidoheptyl)-β-D-
glucopyranoside (III-29b).**

A solution of alcohol III-28 (0.21 g, 0.317 mmol) and benzyl bromide (0.307 g, 1.79 mmol) in THF (4 ml) was sequentially treated with sodium hydride (0.016 g, 0.4 mmol, 60% dispersion in oil) and tetra-n-butylammonium iodide (0.01 g). The mixture was then stirred for 36 h, diluted with saturated aqueous ammonium chloride (10 ml), and poured into ethyl acetate (30 ml). The aqueous phase was extracted with ethyl acetate (3 x 20 ml) and the combined organic solutions were, washed with brine (20 ml), dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (15% ethyl acetate/hexane) furnished III-29b (192 mg, 81% yield) as a colorless oil: $[\alpha]_D^{25}$ +6.2° (c 0.45, CH₂Cl₂); IR (CH₂Cl₂) 3041 (s), 2980 (m), 2940 (m), 2865 (m), 2100 (s),

- 134 -

1610 (m), 1450 (s), 1375 (s), 1262 (s), 1190 (s), 1178 (s),
680 (br) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.97 (dd, $J = 6.4, 0.5$
Hz, 1 H), 7.83 (apparent t, $J = 7.5$ Hz, 2 H), 7.51-7.45 (m,
3 H), 7.38-7.20 (m, 14 H), 4.66 (d, $J = 12.0$ Hz, 1 H), 4.59
5 (d, $J = 11.4$ Hz, 1 H), 4.53 (d, $J = 12.0$ Hz, 1 H), 4.43 (d,
 $J = 11.4$ Hz, 1 H), 4.41 (d, $J = 7.6$ Hz, 1 H), 4.19 (dt, $J =$
9.5, 6.8 Hz, 1 H), 3.82 (dt, $J = 9.5, 7.1$ Hz, 1 H), 3.72 (d,
 $J = 10.9$ Hz, 1 H), 3.59 (dd, $J = 10.9, 4.9$ Hz, 1 H), 3.51-3.39
(m, 4 H), 3.30-3.25 (m, 1 H), 3.18 (t, $J = 6.9$ Hz, 2 H), 3.01
10 (t, $J = 6.9$ Hz, 2 H), 2.49 (dt, $J = 12.2, 4.4$ Hz, 1 H),
1.56-1.49 (m, 5), 1.36-1.31 (m, 4 H); ^{13}C NMR (125 MHz, CDCl_3)
 δ 138.54, 138.32, 138.04, 135.16, 133.58, 131.06, 129.13,
128.41, 128.30, 127.78, 127.69, 127.66, 127.49, 126.69,
124.69, 123.54, 123.09, 119.87, 119.48, 113.69, 105.26, 78.03,
15 74.96, 72.67, 72.29, 71.52, 71.27, 69.94, 68.41, 51.35, 34.99,
29.48, 28.75, 26.53, 25.72, 25.66; high resolution mass
spectrum (FAB, m-nitrobenzyl alcohol) m/z 775.3132 [(M+ Na) $^+$;
calcd for $\text{C}_{42}\text{H}_{48}\text{N}_4\text{O}_7\text{S}$: 775.3142].

20 **AC. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-benzyl-3-deoxy-
6-O-(6-aminohexyl)- β -D-glucopyranoside (III-5b).**

A solution of azide III-29b (0.16 g, 0.21 mmol) in THF
(13.3 ml) was treated sequentially with water (0.093 ml, 5.16
mmol) and triphenylphosphine (0.112 g, 0.43 mmol). The
mixture was then heated at 60°C for 5 h, cooled to room
25 temperature, and concentrated in vacuo. Flash chromatography
(10% methanol/dichloromethane) yielded the corresponding amine
(142.3 mg, 92% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} +7.0^\circ$ (c
1.7, CHCl_3); IR (CH_2Cl_2) 3680 (w), 3045 (m), 2938 (s), 2880
(s), 1606 (m), 1582 (m), 1450 (s), 1370 (s), 1260 (s), 1208
30 (m), 1180 (s), 1090 (s), 1075 (s), 590 (m), 570 (m) cm^{-1} ; 1NMR
(500 MHz, CDCl_3) δ 7.90 (d, $J = 8.4$ Hz, 1 H), 7.76 (d, $J = 7.9$
Hz, 1 H), 7.76 (d, $J = 8.4$ Hz, 1 H), 7.43-7.13 (m, 17 H), 4.58
(d, $J = 12.0$ Hz, 1 H), 4.52 (d, $J = 11.5$ Hz, 1 H), 4.45 (d,
 $J = 12.0$ Hz, 1 H), 4.36 (d, $J = 11.5$ Hz, 1 H), 4.33 (d, $J =$
35 7.5 Hz, 1 H), 4.13 (dt, $J = 9.5, 6.8$ Hz, 1 H), 3.75 (dt, $J =$
9.51, 7.2 Hz, 1 H), 3.65 (d, $J = 10.4$ Hz, 1 H), 3.51 (dd, J
= 10.7, 4.7 Hz, 1 H), 3.44-3.32 (m, 4 H), 3.20 (m, 1 H), 2.93

- 135 -

(t, $J = 6.9$ Hz, 2 H), 2.55 (t, $J = 7.0$ Hz, 2 H), 2.41 (dt, $J = 12.3, 4.2$ Hz, 1 H), 1.53-1.42 (m, 7 H), 1.34-1.18 (m, 6 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.30, 138.06, 137.81, 134.93, 133.32, 130.82, 128.87, 128.15, 128.04, 127.51, 127.45, 5 127.40, 127.23, 126.43, 124.43, 123.29, 122.84, 119.63, 119.24, 113.43, 105.01, 76.49, 72.41, 72.05, 71.42, 71.03, 69.66, 68.14, 44.72, 41.80, 34.77, 33.26, 29.34, 26.45, 25.75, 25.37.

A solution of the above amine (0.119 g, 0.16 mmol) in 10 ethanol (15 ml) was treated with 5 M aqueous potassium hydroxide (3 ml) and then heated to reflux. After 5 h the mixture was cooled, diluted with saturated aqueous ammonium chloride (25 ml), and poured into dichloromethane (30 ml). The aqueous phase was extracted with dichloromethane (4 x 10 15 ml) and the combined organic solutions were dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (15% methanol/dichloromethane) furnished III-5b (80.9 mg, 73% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} +11.8^\circ$ (c 0.43, CH_2Cl_2); IR, 3681 (w), 3436 (m), 3025 (m), 2918 (s), 20 2862 (s), 1729 (m), 1609 (m), 1458 (s), 1251 (m), 1098 (s), 1076 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.64 (br s, 1 H), 7.49 (d, $J = 8.6$ Hz, 1 H), 7.27-7.16 (m, 11 H), 7.05 (apparent t, $J = 7.1$ Hz, 1 H), 6.98 (apparent t, $J = 5.9$ Hz, 1 H), 6.93 (s, 1 H), 4.67 (d, $J = 11.8$ Hz, 1 H), 4.51 (d, $J = 11.8$ Hz, 1 H), 25 4.49 (d, $J = 11.4$ Hz, 1 H), 4.36 (d, $J = 7.6$ Hz, 1 H), 4.31 (d, $J = 11.4$ Hz, 1 H), 4.07 (dt, $J = 9.5, 7.3$ Hz, 1 H), 3.75 (dt, $J = 9.5, 7.5$ Hz, 1 H), 3.44-3.21 (m, 6 H), 3.02 (t, $J = 7.4$ Hz, 2 H), 2.63 (br t, $J = 6.9$ Hz, 2 H), 2.42 (dt, $J = 12.3, 4.7$ Hz, 1 H), 1.49-1.35 (m, 6 H), 1.18-1.1 (m, 5 H); ^{13}C 30 NMR (125 MHz, CDCl_3) δ 138.70, 138.04, 136.20, 128.42, 128.31, 127.75, 127.71, 127.58, 127.50, 122.27, 121.78, 119.11, 118.71, 112.42, 111.22, 105.30, 77.92, 75.09, 72.70, 72.40, 71.31, 71.09, 70.00, 69.93, 39.76, 34.91, 29.29, 27.37, 26.09, 25.82, 25.42; high resolution mass spectrum (FAB *m*-nitrobenzyl 35 alcohol) m/z 609.3332 $[(M+\text{Na})^+]$; calcd for $\text{C}_{36}\text{H}_{46}\text{N}_2\text{O}_5$: 609.3305].

- 136 -

AD. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-benzyl-3,6-dideoxy-6-amino-6-N-(5-hydroxypentyl)- β -D-glucopyranoside (III-5c).

Triflic anhydride (126 ml, 0.748 mmol) was added to a stirred solution of III-28 (360 mg, 0.575 mmol) and 2,6-di-tert-butyl-4-methylpyridine (189 mg, 0.92 mmol) in dichloromethane (3 ml) at -78°C. After 20 min at -78°C, the mixture was allowed to warm to room temperature over 20 min. The resultant suspension was poured into saturated aqueous sodium bicarbonate and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The oily crude triflate was used without purification in the next step.

A solution of 5-trifluoroacetamido-1-pentanol (III-18a) (687 mg, 3.45 mmol) in THF (16 ml) was added to a stirred suspension of sodium hydride (8.63 mmol, 345 mg, 60% dispersion in oil) in THF (20 ml) at 0°C. After 10 min the mixture was allowed to warm to room temperature, stirred for 90 min, recooled to 0°C, and treated with a solution of crude triflate (0.575 mmol) in dichloromethane (22 ml). The suspension was stirred for 30 min at 0°C and then at room temperature for an additional 24 h. The reaction was quenched at 0°C with saturated aqueous ammonium chloride (10 ml) and extracted with ethyl acetate, and the extracts were washed with water and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (gradient elution, 1% to 2% methanol/dichloromethane) afforded an inseparable mixture of compounds, presumably III-29c and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (6 ml) was treated with 5 N NaOH (1 ml, 5 mmol), heated at reflux for 2 h, cooled, and concentrated in vacuo. The residue was taken up in dichloromethane and the resultant solution washed with 2 N HCl. The aqueous layer was extracted with dichloromethane and the combined organic layers were washed

- 137 -

with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (8% methanol/dichloromethane) yielded pure III-5c (172 mg, 52% yield for 3 steps) as a colorless oil: $[\alpha]_D^{25} +17^\circ$ (c 0.15, acetonitrile); UV (6.5×10^{-5} M, acetonitrile) λ_{\max} 281.2 (ϵ 6.2×10^3), 218.8 (3.62×10^4) nm; IR (film) 3325 (m), 3065 (w), 3035 (w), 3015 (w), 2940 (s), 2870 (s), 1500 (w), 1458 (m), 1354 (w), 1220 (w), 1076 (s), 1030 (m), 745 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.44 (s, 1 H), 7.57 (d, $J = 7.7$ Hz, 1 H), 7.31-7.23 (m, 10 H), 7.17-7.14 (m, 1 H), 7.11-7.07 (m, 1 H), 7.04 (d, $J = 2.0$ Hz, 1 H), 4.71 (d, $J = 11.8$ Hz, 1 H), 4.57 (d, $J = 11.7$ Hz, 1 H), 4.56 (d, $J = 11.9$ Hz, 1 H), 4.46 (d, $J = 7.5$ Hz, 1 H), 4.40 (d, $J = 11.5$ Hz, 1 H), 4.20 (ddd, $J = 13.8, 9.4, 6.8$ Hz, 1 H), 3.87 (ddd, $J = 14.9, 9.3, 7.4$ Hz, 1 H), 3.55-3.50 (m, 3 H), 3.32-3.26 (m, 2 H), 3.11 (t, $J = 7.2$ Hz, 2 H), 3.02 (dd, $J = 12.4, 2.9$ Hz, 1 H), 2.68 (dd, $J = 12.4, 8.1$ Hz, 1 H), 2.67-2.57 (m, 2 H), 2.50 (ddd, $J = 12.3, 4.8, 4.8$ Hz, 1 H), 2.20 (s, 3 H), 1.57-1.44 (m, 5 H), 1.36-1.30 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.61, 137.92, 136.14, 128.41, 128.27, 127.79, 127.70, 127.53, 127.49, 122.18, 121.84, 119.18, 118.67, 112.56, 111.12, 105.22, 105.18, 76.53, 75.14, 74.28, 72.69, 70.99, 69.91, 62.45, 50.69, 49.49, 34.86, 32.28, 29.16, 25.80, 23.27; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.3314 [(M+H) $^+$; calcd for $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_5$: 573.3328].

AE. 2-(1H-Indol-3-yl)ethyl 2,4-Di-O-benzyl-3,6-dideoxy-6-amino-6-N-(6-hydroxyhexyl)- β -D-glucopyranoside (III-5d).

A solution of 6-trifluoroacetamido-1-hexanol (III-18c) (147 mg, 0.690 mmol) in THF (1 ml) was added to a suspension of sodium hydride (60% oil dispersion, 69.0 mg, 1.73 mmol) in THF (3 ml) at 0°C . The mixture was stirred at room temperature for 1 h, recooled to 0°C , and treated with a solution of the crude triflate derived from 28 (0.115 mmol), prepared as described for the synthesis of III-5c, in dry dichloromethane (5 ml). The reaction mixture was then warmed to room temperature, stirred for 48 h, and quenched at 0°C

- 138 -

with saturated ammonium chloride solution. The mixture was extracted with ethyl acetate and the combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuo, affording an inseparable mixture of compounds, presumably III-29d and its benzenesulfonamide deprotected counterpart, which was used directly in the next step.

A stirred solution of the above mixture in ethanol (6 ml) was treated with 5 N sodium hydroxide (2 ml), heated to reflux for 2 h, cooled, and concentrated in vacuo. The oily residue was taken up in water and extracted with dichloromethane, and the organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (5% methanol/dichloromethane) yielded III-5d (56 mg, 64% yield for 2 steps) as a colorless oil: $[\alpha]_D^{25} +13^\circ$ (c 0.12, acetonitrile); UV (1.23×10^{-4} M, acetonitrile) λ_{\max} 289.6 (ϵ 1.78×10^3), 280.8 (1.37×10^3), 228.0 (2.63×10^3) nm; IR (film) 3300 (br), 3060 (w), 3030 (w), 2930 (s), 2860 (m), 1450 (m), 1350 (w), 1070 (s), 740 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.16 (br s, 1 H), 7.60 (d, $J = 7.8$ Hz, 1 H), 7.35-7.04 (m, 4 H), 4.71 (d, $J = 11.8$ Hz, 1 H), 4.60 (d, $J = 11.6$ Hz, 1 H), 4.57 (d, $J = 11.9$, 1 H), 4.47 (d, $J = 7.6$ Hz, 1 H), 4.41 (d, $J = 11.5$ Hz, 1 H), 4.20 (dt, $J = 9.4, 6.8$ Hz, 1 H), 3.87 (dt, $J = 9.3, 7.6$ Hz, 1 H), 3.56 (t, $J = 10.0$ Hz, 1 H), 3.52 (m, 1 H), 3.12 (t, $J = 6.9$ Hz, 2 H), 3.04 (d, $J = 2.8$ Hz, 1 H), 3.02 (d, $J = 2.8$ Hz, 1 H), 2.70-2.48 (m, 4 H), 2.05 (br s, 2 H), 1.54 (q, $J = 11.6$ Hz, 1 H), 1.48-1.26 (m, 8 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.61, 137.95, 136.14, 128.40, 128.27, 127.77, 127.69, 127.53, 127.49, 122.12, 121.85, 119.19, 118.68, 112.54, 111.10, 105.24, 76.87, 76.74, 75.17, 74.37, 72.70, 71.00, 69.92, 62.71, 50.81, 49.58, 34.90, 32.53, 29.67, 26.94, 25.81, 25.53; high resolution mass spectrum (Cl, CH_4) m/z 587.3557 [(M+H) $^+$; calcd for $\text{C}_{36}\text{H}_{47}\text{N}_2\text{O}_5$: 587.3485].

AF. 5-Phthalimido-1-pentanol (III-33).

A solution of 5-amino-1-pentanol (5.00 g, 48.5 mmol) in benzene (150 ml) was treated with N-carboethoxyphthalimide (11.0 g, 50.2 mmol) and stirred at room temperature for 5 h.

- 139 -

Concentration in vacuo and flash chromatography (25% ethyl acetate/petroleum ether) yielded III-33 (9.6 mg, 84% yield) as a clear, colorless oil: UV (9.65×10^{-4} M, acetonitrile) λ_{max} 292.0 (ϵ 212), 242.4 (226) nm; IR (CHCl_3) 3460 (br), 2940 (s), 2860 (s), 1770 (s), 1710 (s), 1610 (s), 1470 (s), 1440 (s), 1400 (s), 1370 (s), 1190 (m), 1170 (m), 1130 (m), 1050 (s), 960 (m), 890 (m), 875 (m), 790 (m), 720 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.72-7.70 (m, 2 H), 3.69 (t, $J = 7.2$ Hz, 1 H), 3.64 (t, $J = 6.5$ Hz, 1 H), 2.17 (br s, 1 H), 1.74-1.59 (m, 2 H), 1.46-1.40 (m, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 169.39, 133.78, 131.96, 123.05, 62.34, 37.74, 32.03, 28.22, 22.93; high resolution mass spectrum (Cl , NH_3) m/z 234.1108 [$(\text{M}+\text{H})^+$; calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_3$: 234.1129].

15 AG. 3,4-Di-O-Benzyl-6-O-(5-phthalimidopentyl)-D-glucal (III-34).

5-Phthalimidopentyl triflate was prepared as follows: A stirred solution of 5-phthalimido-1-pentanol (III-33) (1.32 g, 4.67 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.960 g, 4.67 mmol) in dry dichloromethane (10 ml) was treated with triflic anhydride (0.784 ml, 4.67 mmol). After 10 min at room temperature, the mixture was diluted with water (100 ml) and extracted with dichloromethane (2 x 200 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction. Sodium hydride (60% dispersion in oil, 0.20 g, 5.06 mmol) was added to a solution of alcohol III-32 (1.27 g, 3.89 mmol), 5-phthalimidopentyl triflate (4.67 mmol), and 15-crown-5 (20 mg, 2.3 mol %), in dichloromethane (100 ml) at 0°C . After stirring for 24 h at room temperature, the mixture was poured into water. The aqueous layer was extracted with dichloromethane (3 x 50 ml) and the combined extracts were washed with water, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography (3% ether/dichloromethane) provided III-34 (1.82 g, 86% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -8.2^\circ$ (c 0.70, CHCl_3); IR (CHCl_3) 3080 (w), 3020 (m), 3009 (m), 2959 (m), 2880 (m), 1780 (m), 1719

- 140 -

(s), 1652 (m), 1500 (w), 1470 (w), 1457 (m), 1440 (m), 1400 (s), 1365 (m), 1235 (m), 1110 (br, s), 1058 (br, s), 908 (w), 692 (m), cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.80 (m, 2 H), 7.68 (m, 2 H), 7.25-7.34 (m, 10 H), 6.38, (dd, $J = 6.1, 1.2$ Hz, 1 H), 4.84 (m, 2 H), 4.66 (d, $J = 11.4$ Hz, 1 H), 4.63 (d, $J = 11.7$ Hz, 1 H), 4.55 (d, $J = 11.7$ Hz, 1 H), 4.19 (m, 1 H), 4.00 (m, 1 H), 3.81 (dd, $J = 8.7, 6.2$ Hz, 1 H), 3.64-3.74 (m, 4 H), 3.40-3.50 (m, 2 H), 1.60-1.70 (m, 4 H), 1.40 (m, 2 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 168.4, 144.8, 138.4, 138.3, 133.9, 132.2, 128.4, 127.9, 127.8, 127.6, 123.2, 99.9, 76.8, 75.8, 74.5, 73.8, 71.4, 70.5, 69.2, 37.9, 29.2, 28.5, 23.5; high resolution mass spectrum (Cl, NH_3) m/z 541.2483 (M^+ ; calcd for $\text{C}_{33}\text{H}_{35}\text{NO}_6$: 541.2464).

15 **AH. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
3,4-Di-O-benzyl-6-O-(5-phthalimidopentyl)- β -D-glucopyranoside (III-35).**

A solution of dimethyldioxirane in acetone (1.2 equiv, ca. 0.05 M) was added dropwise to glycal III-34 (1.53 g, 2.80 mmol) in dichloromethane (26 ml) at 0°C . The mixture was stirred at 0°C for 1 h and concentrated in vacuo. To a solution of the crude epoxide and III-12 (1.15 g, 3.82 mmol) in THF (12 ml) at -78°C was added ZnCl_2 (1.0 M in ether, 5.6 ml, 5.6 mmol) and the mixture was allowed to stir at -78°C for 1 h. The solution was then slowly warmed to room temperature and stirred 18 h. The mixture was poured into saturated aqueous sodium bicarbonate (50 ml) and extracted with ethyl acetate (3 x 50 ml) and the combined extracts were washed with water, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography (45% ethyl acetate/hexane) yielded III-35 (1.05 g, 44% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -8.1^\circ$ (c 1.8 CHCl_3); IR (CHCl_3) 3069 (w), 3039 (m), 3019 (m), 2955 (m), 2879 (m), 1780 (m), 1719 (s), 1612 (w), 1472 (w), 1451 (s), 1401 (s), 1370 (s), 1175 (s), 1121 (s), 1068 (s), 695 (w), 680 (w), 596 (m), 570 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.96 (dd, $J = 8.1, 0.6$ Hz, 1 H), 7.85 (dd, $J = 8.2, 0.9$ Hz, 2 H), 7.78 (m, 2 H), 7.66 (m, 2 H), 7.20-7.50 (m, 17 H), 4.89 (d, $J = 11.3$ Hz, 1 H), 4.86 (d, $J = 11.0$ Hz, 1 H),

- 141 -

4.83 (d, $J = 11.4$ Hz, 1 H), 4.60 (d, $J = 10.9$ Hz, 1 H), 4.24 (d, $J = 7.6$ Hz, 1 H), 4.20 (dt, $J = 9.5, 6.4$ Hz, 1 H), 3.76 (dt, $J = 9.5, 7.2$ Hz, 1 H), 3.37-3.68 (m, 10 H), 2.98 (m, 2 H), 2.13 (br s, 1 H), 1.57-1.68 (m, 4 H), 1.38 (m, 2 H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 168.4, 138.6, 138.2, 135.1, 133.8, 133.7, 132.1, 131.0, 129.1, 128.4, 127.9, 127.8, 127.7, 126.7, 124.7, 123.5, 123.1, 119.7, 119.4, 113.7, 102.8, 84.4, 76.5, 75.1, 71.5, 69.6, 68.7, 37.8, 29.2, 28.4, 25.4, 23.5; high resolution mass spectrum (Cl , NH_3) m/z 662.2774 (M^+ ; calcd for $\text{C}_{35}\text{H}_{42}\text{SO}_7$: 662.2775).

AI. 2-Deoxy-3,4-di-O-benzyl-6-O-(5-phthalimido-pentyl)- β -D-glucopyranoside (III-36).

A solution of III-35 (0.455 g, 0.530 mmol) in THF (10 ml) was cooled to -78°C and treated with carbon disulfide (27 ml, 0.583 mmol) followed by sodium bis(trimethylsilyl)amide (0.6 M in toluene, 0.953 ml, 0.572 mmol). After 20 min, methyl iodide (59 ml, 0.640 mmol) was added and the solution was stirred for 5 min at -78°C and then at room temperature for 45 min. The reaction mixture was quenched with water (50 ml) and extracted with ethyl acetate (3 x 50 ml). The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording the crude xanthate as a pale yellow oil (0.462 g, 92% yield) which was used without purification in the next step.

To a solution of the crude xanthate (0.462 g, 0.487 mmol) and AIBN (10 mg) in toluene (8 ml) was added tributyltin hydride (0.214 ml, 0.795 mmol) and the reaction mixture heated to reflux for 4 h, cooled, and concentrated in vacuo. The residue was taken up in acetonitrile (30 ml) and washed with petroleum ether (5 x 10 ml), dried over sodium sulfate, filtered, and concentrated in vacuo to an oil. Flash chromatography (20% ethyl acetate/petroleum ether) yielded III-36 (0.296 g, 72% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -10^\circ$ (c 1.1 CHCl_3); IR (CHCl_3) 3062 (w), 3031 (w), 3009 (w), 2939 (m), 2864 (m), 1777 (w), 1712 (s), 1610 (w), 1469 (w), 1449 (m), 1396 (s), 1378 (s), 1181 (m), 1171 (s), 1120 (s), 1090 (s), 990 (w), 910 (s), 692 (w), 595 (m) cm^{-1} ; ^1H NMR (500 MHz,

- 142 -

CDCl₃) δ 7.96 (d, J = 7.5 Hz, 1 H), 7.84 (m, 2 H), 7.79 (m, 2 H), 7.66 (m, 2 H), 7.20-7.41 (m, 15 H), 4.91 (d, J = 11.0 Hz, 1 H), 4.60 (m, 2 H), 4.66 (d, J = 11.7 Hz, 1 H), 4.41 (dd, J = 9.7, 1.8 Hz, 1 H), 4.15 (dt, J = 9.5, 6.6 Hz, 1 H),
5 3.59-3.71 (m, 6 H), 3.47 (m, 2 H), 3.40 (m, 1 H), 2.94 (t, J = 6.6 Hz, 2 H), 2.57 (ddd, J = 14.2, 5.0, 3.2 Hz, 1 H), 1.57-1.68 (m, 5 H), 1.38 (m, 2 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.5, 25.5, 28.4, 29.2, 36.7, 37.9, 68.1, 70.0, 71.4, 75.0, 75.2, 78.2, 79.3, 99.9, 113.6, 119.6, 123.1, 123.5, 124.7,
10 126.7, 127.7, 128.0, 128.4, 129.2, 131.1, 132.1, 133.6, 133.8, 135.1, 138.3, 138.5, 168.4; high resolution mass spectrum (Cl, NH₃) m/z 814.3287 (M⁺; calcd for C₄₄H₅₀SO₈N₂: 814.3289).

AJ. 2 - (1 H - I n d o l - 3 - y l) e t h y l
2 - D e o x y - 3 , 4 - d i - O - b e n z y l - 6 - O -
15 (5-aminopentyl)- β -D-glucopyranoside (III-6).

A solution of hydrazine (0.2 M in MeOH, 6 ml) was added to III-36 (0.034 g, 0.043 mmol). After stirring for 16 h, the reaction mixture was concentrated in vacuo, the residue dissolved in ethanol (4 ml), and 5N NaOH (0.90 ml) added. The
20 mixture was heated at reflux for 4 h, cooled, and extracted with dichloromethane (3 x 10ml). The combined extracts were washed with brine, dried over magnesium sulfate, and concentrated in vacuo to an oil. Flash chromatography (11% methanol/dichloromethane) afforded 6 (11 mg, 44%) as a pale
25 yellow oil: $[\alpha]_D^{25}$ -15° (c 0.62, CHCl₃); IR (CHCl₃) 3490 (m), 3345 (br, m), 3020 (m), 2945 (s), 2882 (s), 1625 (w), 1500 (w), 1459 (m), 1370 (m), 1230 (w), 1100 (s), 695 (w) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.80 (br s, 1 H), 7.49 (d, J = 7.9 Hz, 1 H), 7.19-7.31 (m, 11 H), 7.10 (t, J = 7.1 Hz, 1 H), 7.00 (t, J = 8.0 Hz, 1 H), 6.97 (s, 1 H), 4.83 (d, J = 11.1 Hz, 1 H),
30 4.59 (d, J = 11.7 Hz, 1 H), 4.51 (d, J = 11.0 Hz), 4.50 (d, J = 11.7, 1 H), 4.39 (d, J = 9.7 Hz, 1 H), 4.00 (apparent q, J = 7.3 Hz, 1 H), 3.67 (apparent q, J = 7.3 Hz, 1 H), 3.60 (d, J = 9.0 Hz, 1 H), 3.56 (m, 1 H), 3.46 (dd, J = 10.8, 5.3 Hz),
35 3.31 (m, 4 H), 2.98 (t, J = 7.2 Hz, 2 H), 2.50 (t, J = 7.3 Hz, 2 H), 2.28 (m, 2 H), 1.57 (q, J = 10 Hz, 1 H), 1.42 (m, 4 H), 1.19 (m, 2 H); ¹³C NMR (62.9 MHz, CDCl₃) δ 138.3, 138.2, 136.2,

- 143 -

128.4, 128.0, 127.7, 127.5, 122.3, 121.8, 119.1, 118.7, 112.0, 111.4, 99.9, 79.3, 78.2, 74.9, 71.4, 71.0, 69.9, 69.8, 39.7, 36.7, 28.8, 27.6, 25.7, 23.1; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 573.3371 [(M + H)⁺; calcd for C₃₅H₄₄N₂O₅: 573.3328].

AK. Methyl 2,3,6-Tri-O-benzoyl-4-deoxy- α -D-glucopyranoside (III-38).

A solution of III-37 (5.00 g, 9.87 mmol) in THF (100 ml) was cooled to -78°C and treated with carbon disulfide (0.45 ml, 7.48 mmol) followed by sodium bis(trimethylsilyl)amide (1.0 M in THF, 10.5 ml, 51.8 mmol). After 20 min, methyl iodide (2.10 ml, 33.7 mmol) was added and the solution was stirred for 5 min at -78°C and then at room temperature for 45 min. The reaction mixture was quenched with water (5 ml) and extracted with ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording the crude xanthate as a pale yellow oil (5.70g, 97% yield) which was used without purification in the next step. Purification of an analytical sample by flash chromatography (20% ethyl acetate/petroleum ether) gave white crystals: mp 72-73°C; $[\alpha]_D^{25}$ +140° (c 0.13, acetonitrile); ¹³C NMR (125 MHz, CDCl₃) δ 166.10, 165.73, 165.53, 133.37, 133.13, 129.90, 129.75, 129.70, 129.21, 128.90, 128.37, 128.23, 96.94, 76.25, 71.83, 70.45, 67.36, 62.58, 55.60, 19.18; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 597.1286 [(M+H)⁺; calcd for C₃₀H₂₈O₉S₂: 597.1253].

Tributyltin hydride (6.68 ml, 24.8 mmol) was added to a solution of the crude xanthate (5.70 g, 9.55 mmol) and AIBN (50 mg) in toluene (120 ml), and the reaction mixture was then heated to reflux for 4 h, cooled, and concentrated in vacuo. The residue was taken up in acetonitrile (200 ml) and extracted with petroleum ether (5 x 100 ml). The acetonitrile solution was dried over sodium sulfate, filtered, and concentrated in vacuo, affording a clear, colorless oil which solidified on standing. Flash chromatography (20% ethyl acetate/petroleum ether) yielded III-38 (3.60 g, 82% yield)

- 144 -

as a white solid: mp 119-120°C; $[\alpha]_D^{25} +121^\circ$ (c 0.17, acetonitrile); IR (CHCl₃) 3010 (m), 1730 (s), 1600 (w), 1580 (w), 1460 (m), 1270 (s), 1220 (s), 1110 (s), 1080 (m), 1060 (m), 1040 (m), 750 (s), 710 (s), 660 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.08 (dd, *J* = 8.3, 1.2 Hz, 2 H), 8.05 (dd, *J* = 8.3, 1.2 Hz, 2 H), 8.00 (dd, *J* = 8.4, 1.3 Hz, 2 H), 7.51-7.35 (m, 9 H), 5.78 (m, 1 H), 5.31 (dd, *J* = 10.2, 3.6 Hz, 1 H), 5.15 (d, *J* = 3.6 Hz, 1 H), 4.45-4.43 (m, 3 H), 3.44 (s, 3 H), 2.47 (ddd, *J* = 12.5, 5.2, 2.1, 1 H), 1.89 (q, *J* = 12 Hz, 1 H); ¹³C NMR (125 MHz, CDCl₃) δ 166.23, 166.09, 165.81, 133.22, 133.16, 133.09, 129.84, 129.67, 129.62, 129.41, 128.42, 128.35, 128.32, 97.82, 72.57, 68.38, 66.05, 65.33, 55.32, 33.16; high resolution mass spectrum (Cl, NH₃) *m/z* 536.1902 [(M+NH₄)⁺; calcd for C₂₈H₃₀N₁O₈: 536.1919].

15 **AL. Acetyl 2,3,6-Tri-O-benzoyl-4-deoxy-α-D-glucopyranoside (III-39).**

A solution of glycoside III-38 (0.50 g, 1.1 mmol) in acetic anhydride (3.0 ml, 32 mmol) was cooled to 0°C and treated with boron trifluoride etherate (0.1 ml). The reaction mixture was then stirred at room temperature for 4 h, diluted with ethyl acetate, and poured into ice-cold saturated sodium bicarbonate. The layers were separated and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording III-39 (0.45 g, 85% yield) as a colorless oil which crystallized upon standing as white needles: mp 123-124°C; $[\alpha]_D^{25} +123^\circ$ (c 0.19, acetonitrile); IR (CHCl₃) 3020 (s), 2400 (w), 1760 (m), 1730 (s), 1460 (w), 1280 (s), 1220 (s), 1110 (s), 930 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.07-8.05 (m, 2 H), 7.97-7.92 (m, 5 H), 7.51-7.36 (m, 8 H), 6.58 (d, *J* = 3.7 Hz, 1 H), 5.78 (m, 1 H), 5.52 (m, 1 H), 4.46 (m, 3 H), 2.52 (ddd, *J* = 12.5, 5.2, 2.1 Hz, 1 H), 2.17 (s, 3 H), 2.03 (m, 1 H); ¹³C NMR (125 MHz, CDCl₃) δ 168.87, 166.16, 165.54, 133.35, 133.29, 133.22, 129.71, 129.66, 129.58, 129.35, 129.02, 128.42, 128.39, 90.32, 71.59, 71.36, 70.78, 68.12, 68.05, 65.57, 32.76, 20.86, 20.80;

- 145 -

high resolution mass spectrum (Cl, NH₃) m/z 536.1902 [(M+NH₄)⁺; calcd for C₂₉H₂₆O₉: 536.1919].

AM. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3,6-Tri-O-benzoyl-4-deoxy- β -D-glucopyranoside
(III-40).

5 A stirred solution of acetate III-39 (0.137 g, 0.29 mmol) in dichloromethane (3 ml) was cooled to 0°C and treated with 30% hydrogen bromide in acetic acid (0.07 ml, 0.33 mmol). The reaction mixture was stirred at room temperature for 4 h,
10 diluted with ethyl acetate, washed with saturated aqueous sodium bicarbonate and brine, dried over sodium sulfate, filtered, and concentrated in vacuo, furnishing a colorless oil which solidified upon standing. Recrystallization (ether/petroleum ether) yielded the bromide (0.15 g, 100%
15 yield) as white crystals: mp 134-135°C; [α]D²⁵ +114° (c 0.10, acetonitrile); ¹³C NMR (125 MHz, CDCl₃) δ 166.11, 165.64, 165.53, 133.65, 133.35, 133.32, 130.01, 129.78, 129.69, 129.49, 129.31, 128.75, 128.48, 128.42, 88.85, 71.54, 70.78, 68.63, 65.05, 32.16; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 539.0661 [(M+H)⁺; calcd for C₂₇H₂₃O₇Br: 539.0705].
20

A solution of the above bromide (0.40 g, 0.814 mmol) in hexane and benzene (2:3, 17 ml) was added to a mixture of activated, powdered 4 Angstrom molecular sieves (0.83 g),
25 protected tryptophol III-12 (0.37 g, 1.23 mmol) and silver(I) oxide (0.83 g, 3.58 mmol) in a flask wrapped with aluminum foil. The mixture was stirred at room temperature for two days, filtered through Celite, and concentrated in vacuo to furnish a colorless oil. Flash chromatography (50%
30 ether/petroleum ether) then yielded III-40 (0.50 g, 81% yield) as a colorless solid: mp 76-78°C; [α]D²⁵ +28° (c 0.12, acetonitrile); UV (9.21 x 10⁻⁵ M, acetonitrile) λ_{max} 237.6 (ϵ 4.47 x 10³), 198.8 (4.10 x 10³) nm; IR (CHCl₃) 3010 (s), 1730 (s), 1455 (m), 1380 (m), 1320 (m), 1280 (s), 1220 (s), 1180 (s), 1120 (s), 1100 (m), 1075 (m), 1030 (m), 770 (s), 710 (s),
35 670 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.03-7.10 (m, 2 H), 5.42 (m, 2 H), 4.74 (d, J = 7.5 Hz, 1 H), 4.47 (m, 2 H),

- 146 -

4.16-4.05 (m, 2 H), 3.82 (m, 1 H), 2.91 (m, 2 H), 2.47 (ddd, $J = 12.5, 4.6, 1.9$ Hz, 1 H), 1.90 (q, $J = 13.0$ Hz, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 166.20, 165.89, 165.42, 135.03, 133.55, 133.24, 133.22, 133.06, 130.86, 129.72, 129.67, 129.62, 129.49, 129.32, 129.12, 128.42, 128.37, 128.31, 126.67, 124.58, 123.43, 123.06, 119.42, 119.35, 113.56, 101.42, 72.53, 71.56, 69.75, 68.80, 65.81, 33.00, 25.60; high resolution mass spectrum (FAB, *m*-nitrobenzyl alcohol) m/z 759.2108 (M^+ ; calcd for $\text{C}_{43}\text{H}_{37}\text{NO}_{10}\text{S}$: 759.2138).

10 **AN. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
4-Deoxy- β -D-glucopyranoside (III-41).**

A solution of tribenzoate III-40 (120 mg, 0.158 mmol) in methanol (20 ml) was treated with sodium methoxide (0.027 g, 0.507 mmol) and then stirred for 16 h. The mixture was
15 neutralized with Amberlyst® 15 ion exchange resin, filtered, and the filtrate was concentrated in vacuo to yield a tan solid. Flash chromatography (10% methanol/dichloromethane) yielded III-41 (65 mg, 91% yield) as a white solid: mp 64-65°C; $[\alpha]_{\text{D}}^{25} -29^\circ$ (c 0.15, acetonitrile); UV (9.21×10^{-5} M, acetonitrile) λ_{max} 253.2 (ϵ 1.55×10^3), 212.0 (2.58×10^4)
20 nm; IR (CHCl_3) 3420 (w), 3010 (m), 1455 (m), 1370 (m), 1280 (w), 1220 (s), 1180 (m), 1120 (m), 1075 (m), 760 (s), 690 (w), 670 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.99-7.22 (m, 10 H), 4.22 (m, 2 H), 3.82 (m, 1 H), 3.69 (m, 2 H), 3.61 (m, 2 H),
25 3.24 (m, 1 H), 2.97 (m, 2 H), 2.76 (br s, 1 H), 2.61 (br s, 1 H), 1.89 (ddd, $J = 13.1, 5.1, 1.7$ Hz, 1 H), 1.56 (q, $J = 11.5$ Hz, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.19, 135.14, 133.74, 131.04, 129.23, 126.70, 124.83, 123.68, 123.22, 119.74, 119.31, 113.76, 102.92, 76.09, 72.75, 70.72, 68.72,
30 65.04, 33.75, 25.40; high resolution mass spectrum (FAB, *m*-nitrobenzyl alcohol) m/z 447.1389 (M^+ ; calcd for $\text{C}_{22}\text{H}_{25}\text{NO}_7\text{S}$: 447.1352).

35 **AO. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
4-Deoxy-6-O-tert-butyldimethylsilyl- β -D-glucopyranoside (III-42).**

A solution of triol III-41 (0.24 g, 0.536 mmol) in DMF (6 ml) was treated with imidazole (73 mg, 1.07 mmol) followed

- 147 -

by *tert*-butyldiphenylsilyl chloride (0.17 ml, 0.643 mmol). The reaction mixture was then heated at 70°C for 48 h, cooled, quenched with methanol (5 ml), and concentrated in vacuo. The residue was extracted with ethyl acetate and the extracts were
5 washed with brine, dried over sodium sulfate, filtered and concentrated in vacuo. The resultant pale yellow oil was purified by flash chromatography (3% methanol/dichloromethane) to give III-42 (0.36 g, 97% yield) as a colorless oil: $[\alpha]_D^{25}$ -24° (c 0.37, acetonitrile); UV (1.75×10^{-4} M, acetonitrile)
10 λ_{\max} 253.2 (ϵ 1.53×10^3), 212.0 (2.58×10^3) nm; IR (CHCl₃) 3440 (br), 3010 (m), 2960 (w), 2940 (m), 2870 (m), 1455 (m), 1430 (m), 1380 (m), 1280 (w), 1220 (s), 1180 (s), 1120 (s), 1070 (s), 1020 (w), 760 (s), 705 (m), 690 (m), 670 (m) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.98 (d, J = 8.4 Hz, 1 H), 7.82 (m,
15 2 H), 7.65 (m, 4 H), 7.39-7.17 (m, 13 H), 4.17 (d, J = 7.7 Hz, 1 H), 4.17-4.13 (m, 2 H), 3.81-3.62 (m, 3 H), 3.32 (t, J = 8.0 Hz, 2 H), 2.99-2.96 (m, 2 H), 2.76 (br s, 1 H), 2.59 (br s, 1 H), 2.15-2.08 (ddd, J = 13.1, 5.1, 1.7 Hz, 1 H), 1.45 (q, J = 12.7 Hz, 1 H), 1.04 (s, 9 H); ¹³C NMR (125 MHz, CDCl₃) δ
20 138.22, 135.55, 135.52, 133.64, 133.33, 129.69, 129.66, 129.16, 127.65, 126.65, 124.76, 123.49, 123.15, 119.75, 119.41, 113.71, 102.80, 76.24, 72.66, 70.82, 68.64, 66.09, 34.75, 26.75, 25.48, 19.20; high resolution mass spectrum (Cl, NH₃) m/z 703.2929 [(M+NH₄)⁺; calcd for C₃₈H₄₇N₂O₇SSi:
25 703.2873].

**AP. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3-Di-O-benzyl-4-deoxy-6-O-*tert*-butyldimethyl-
silyl- β -D-glucopyranoside (III-43).**

A solution of diol III-42 (0.50 g, 0.729 mmol) in THF
30 (7 ml) was added to a stirred suspension of sodium hydride (73.0 mg, 3.04 mmol, 60% oil dispersion) in THF (3 ml) at 0°C, and the reaction was stirred at room temperature for 30 min. The mixture was recooled to 0°C and benzyl bromide (0.26 ml, 2.2 mmol) was added dropwise. After 3 days at room
35 temperature, the reaction mixture was quenched with saturated aqueous ammonium chloride (10 ml) and extracted with ether. The extracts were washed with brine, dried over sodium

- 148 -

sulfate, filtered, and concentrated in vacuo. Flash chromatography (33% ether/petroleum ether) afforded III-43 (0.73 g, 76% yield) as a colorless oil: $[\alpha]_D^{25} -5.6^\circ$ (c 0.16, acetonitrile); UV (1.44×10^{-4} M, acetonitrile) λ_{\max} 252.8 (ϵ 2.27×10^3), 222.0 (2.63×10^3) nm; IR (CHCl_3) 3080 (w), 3010 (m), 2900 (m), 2850 (m), 1450 (m), 1430 (m), 1380 (m), 1220 (m), 1180 (m), 1100 (s), 750 (s), 700 (s), 660 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.98 (d, $J = 8.3$ Hz, 1 H), 7.80 (dd, $J = 8.1, 0.83$ Hz, 2 H), 7.64 (m, 4 H), 7.32 (m, 23 H), 4.67 (m, 4 H), 4.33 (d, $J = 7.7$ Hz, 1 H), 4.14 (m, 1 H), 3.81-3.77 (m, 2 H), 3.62 (m, 1 H), 3.57-3.48 (m, 2 H), 3.47-3.29 (m, 1 H), 3.29 (t, $J = 7.8$ Hz, 1 H), 2.99 (t, $J = 7.1$ Hz, 1 H), 2.13 (ddd, $J = 12.8, 5.2, 1.6$ Hz, 1 H), 1.40 (q, $J = 11.7$ Hz, 1 H), 1.08 (s, 9 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.64, 138.31, 135.58, 135.54, 135.19, 133.54, 133.46, 130.99, 129.70, 129.67, 129.10, 128.33, 128.20, 127.95, 127.66, 127.62, 127.54, 127.44, 126.62, 124.70, 123.42, 123.11, 119.74, 119.51, 113.69, 103.84, 82.95, 76.74, 74.89, 72.24, 68.55, 66.22, 33.66, 26.80, 25.80, 19.23; high resolution mass spectrum (Cl , NH_3) m/z 883.3898 [$(\text{M}+\text{NH}_4)^+$]; calcd for $\text{C}_{52}\text{H}_{59}\text{N}_2\text{O}_7\text{SSi}$: 883.3812].

**AQ. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3-Di-O-benzyl-4-deoxy- β -D-glucopyranoside (III-44).**

A solution of silyl ether III-43 (0.37 g, 0.427 mmol) in THF (11 ml) was treated with tetrabutylammonium fluoride (1.33 ml, 1.0 M in THF, 1.33 mmol) and stirred at room temperature for 3 h. The solution was then diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (33% petroleum ether/ethyl acetate) yielded III-44 (0.43 g, 85% yield) as a colorless oil: $[\alpha]_D^{25} -4.4^\circ$ (c 0.32, acetonitrile); IR (CHCl_3) 3600 (w), 3480 (br), 3010 (m), 2920 (m), 2890 (m), 1450 (m), 1380 (m), 1220 (s), 1180 (m), 1120 (m), 1100 (m), 760 (s), 700 (m), 690 (m), 670 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.98 (d, $J = 8.3$ Hz, 1 H), 7.83 (m, 2 H), 7.52-7.04 (m, 12 H), 4.74-4.66 (m, 5 H), 4.41 (d,

- 149 -

$J = 6.9$ Hz, 1 H), 4.19 (m, 1 H), 3.88 (m, 1 H), 3.67-3.50 (m, 4 H), 3.31-3.27 (m, 1 H), 2.99 (m, 2 H), 2.08 (t, $J = 5.9$ Hz, 1 H), 1.98 (ddd, $J = 12.8, 5.2, 1.9$ Hz, 1 H), 1.56 (q, $J = 11.7$ Hz, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.54, 138.19, 135.14, 133.61, 130.99, 129.12, 128.32, 128.20, 127.92, 127.60, 127.56, 127.49, 126.63, 124.72, 123.58, 123.13, 119.69, 119.37, 113.69, 103.84, 82.74, 78.11, 74.93, 72.29, 72.19, 68.65, 65.12, 32.61, 25.65; high resolution mass spectrum (Cl, CH_4) m/z 645.2675 [$(\text{M}+\text{NH}_4)^+$; calcd for $\text{C}_{36}\text{H}_{41}\text{N}_2\text{O}_7\text{S}$: 645.2634].

AR. 2 - (N-Phenylsulfonylindol-3-yl)ethyl
2,3-Di-O-benzyl-4-deoxy-6-O-(5-phthalimido-
pentyl)- β -D-glucopyranoside (III-45).

5-Phthalimidopentyl triflate was prepared as follows:

A stirred solution of 5-phthalimido-1-pentanol (III-33) (39.1 mg, 0.168 mmol) and 2,6-di-tert-butyl-4-methylpyridine (34.5 mg, 0.168 mmol) in dry dichloromethane (1.5 ml) was treated with triflic anhydride (28.3 ml, 0.168 mmol). After 10 min at room temperature, the mixture was diluted with water (25 ml) and extracted with dichloromethane (2 x 50 ml). The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo, affording a yellow solid which was used without purification in the next reaction. Sodium hydride (60% dispersion in oil, 51 mg, 1.3 mmol) was added to a solution of alcohol III-44 (150 mg, 0.240 mmol), 5-phthalimidopentyl triflate (1.37 mmol), and 2,6-di-tert-butyl-4-methylpyridine (282 mg, 1.39 mmol), in dichloromethane (1.5 ml) at 0°C. The reaction mixture was stirred for 48 h at room temperature, quenched with saturated aqueous ammonium chloride, and extracted with dichloromethane, and the organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (20% ethyl acetate/petroleum ether) gave III-45 (158 mg, 78% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -2.5^\circ$ (c 0.36, acetonitrile); UV (2.14×10^{-4} M, acetonitrile) λ_{max} 283.6 (ϵ 710), 242.4 (808) nm; IR (CHCl_3) 2940 (m), 2860 (m), 1775 (m), 1715 (s), 1450 (m), 1400 (s), 1370 (s), 1175 (m), 1120 (s),

- 150 -

1090 (s), 1050 (s), 745 (m), 720 (s), 700 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.96 (d, J = 8.3 Hz, 1 H), 7.84-7.80 (m, 4 H), 7.69-7.64 (m, 2 H), 7.50-7.17 (m, 12 H), 4.69 (d, J = 11.0 Hz, 1 H), 4.67 (s, 2 H), 4.64 (d, J = 11.0 Hz, 1 H), 4.36 (d, J = 7.7 Hz, 1 H), 4.21-4.17 (m, 1 H), 3.86-3.81 (m, 1 H), 3.66 (t, J = 7.3 Hz, 2 H), 3.60-3.39 (m, 6 H), 3.28 (dd, J = 7.8, 8.8 Hz, 1 H), 3.00 (t, J = 6.7 Hz, 2 H), 2.12 (dd, J = 5.4, 12.2 Hz, 1 H), 1.71-1.58 (m, 5 H), 1.47-1.36 (m, 3 H); ^{13}C NMR (125 MHz, CDCl_3) δ 168.37, 138.61, 138.31, 135.19, 133.83, 133.56, 132.13, 131.03, 129.11, 128.31, 128.19, 127.96, 127.63, 127.51, 127.44, 126.65, 124.68, 123.51, 123.12, 119.78, 119.49, 113.70, 103.85, 82.83, 78.23, 74.90, 73.10, 72.16, 71.39, 70.95, 68.68, 37.86, 33.94, 29.67, 29.11, 28.36, 25.75, 23.41; high resolution mass spectrum (FAB, *m*-nitrobenzyl alcohol) m/z 865.3201 [($\text{M}+\text{Na}$) $^+$]; calcd for $\text{C}_{49}\text{H}_{50}\text{N}_2\text{O}_9\text{SNa}$: 865.3134].

AS. 2-(1*H*-Indol-3-yl)ethyl 2,3-Di-O-benzyl-6-O-(5-aminopentyl)- β -D-glucopyranoside (III-7).

Sodium methoxide (40 mg, 0.740 mmol) was added to a solution of III-45 (150 mg, 0.178 mmol) in methanol (8 ml) and the reaction mixture was then heated at reflux for 24 h, cooled, poured into water (100 ml), and extracted with dichloromethane (2 x 100 ml). The combined organic layers were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) afforded III-7 (72.0 mg, 71% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} +3.9^\circ$ (c 1.8, acetonitrile); UV (1.57×10^{-4} M, acetonitrile) λ_{max} 280.0 (ϵ 1.41×10^3), 224.8 (1.66×10^3) nm; IR (CHCl_3) 3350 (br), 3060 (w), 2930 (m), 2860 (m), 1630 (m), 1590 (m), 1560 (m), 1450 (m), 1400 (m), 1270 (m), 1100 (s), 740 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.74 (br m, 1 H), 7.48 (d, J = 7.8 Hz, 1 H), 7.36-6.93 (m, 15 H), 4.62-4.49 (m, 4 H), 4.32 (d, J = 7.7 Hz, 1 H), 4.11 (dt, J = 9.4, 6.7 Hz, 1 H), 3.78 (dt, J = 9.2, 7.4 Hz, 1 H), 3.52 (m, 4 H), 3.26 (m, 2 H), 3.22 (t, J = 7.2 Hz, 1 H), 3.13 (t, J = 7.8 Hz, 1 H), 3.00 (t, J = 7.0 Hz, 2 H), 2.00 (ddd, J = 6.7, 5.2, 1.4 Hz, 1 H), 1.29 (m, 9 H); ^{13}C NMR

- 151 -

(125 MHz, CDCl₃) δ 140.11, 138.10, 130.75, 130.59, 129.31, 128.92, 128.84, 128.57, 128.44, 123.70, 122.24, 119.40, 112.82, 112.31, 105.01, 84.13, 79.55, 75.76, 74.12, 73.12, 72.53, 72.18, 71.29, 41.05, 34.54, 30.38, 29.90, 27.07, 24.72;
5 high resolution mass spectrum (Cl, NH₃) m/z 573.3301 [(M+H)⁺; calcd for C₃₅H₄₅N₂O₅: 573.3328].

**AT. Methyl 2,3,4-Tri-O-benzyl-6-O-(5-azidopentyl)-
 β -D-glucopyranoside (III-47a).**

At room temperature a solution of 5-azido-1-pentanol
10 (0.18 g, 1.40 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.3 g, 1.46 mmol) in dichloromethane (10 ml) was treated dropwise with triflic anhydride (0.240 ml, 1.43 mmol). After 15 min the mixture was diluted with dichloromethane (40 ml) and poured into saturated aqueous sodium bicarbonate. The
15 organic phase was washed with brine (2 x 20 ml), dried over magnesium sulfate, filtered, and concentrated, affording a light yellow solid which was used without purification. The alcohol III-46 (0.2 g, 0.429 mmol) and the crude triflate were dissolved in dichloromethane (2 ml) and treated with sodium
20 hydride (0.025 g, 0.625 mmol, 60% dispersion in oil). The mixture was stirred for 48 h, diluted with dichloromethane (40 ml), and poured into saturated aqueous ammonium chloride (40 ml). The aqueous phase was extracted with dichloromethane (3 x 20 ml) and the combined organic solutions were washed with
25 brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (15% ethyl acetate/hexane) provided III-47a (0.126 g, 51% yield) as a white solid: $[\alpha]_D^{25} +7.7^\circ$ (c 0.75, CHCl₃); IR 3028 (m), 2921 (m), 2863 (m), 2110 (s), 1497 (w), 1462 (m), 1421 (m), 1356 (m), 1280 (s), 1070 (s), 732 (br) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.23 (m, 15 H), 4.92 (d, J = 10.9 Hz, 1 H), 4.91 (d, J = 11.0 Hz, 1 H), 4.86 (d, J = 10.9 Hz, 1 H), 4.78 (d, J = 7.8 Hz, 1 H), 3.70-3.50 (m, 6 H), 3.56 (s, 3 H), 3.44-3.40 (m, 3 H), 3.23 (t, J = 6.9 Hz, 2 H), 1.63-1.40 (m, 6 H); ¹³C NMR
35 (125 MHz, CDCl₃) δ 138.59, 138.53, 138.27, 128.42, 128.35, 128.33, 128.07, 127.88, 127.83, 127.76, 127.60, 127.50, 104.73, 84.63, 82.32, 77.96, 75.67, 74.97, 74.84, 74.72,

- 152 -

71.41, 69.70, 57.08, 51.35, 29.22, 28.69, 23.44; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 598.2880 [(M+Na)⁺; calcd for C₃₃H₃₉N₃O₆: 598.2893].

AU. Methyl 2,3,4-Tri-O-benzyl-6-O-(5-aminopentyl)- β -D-glucopyranoside (III-8a).

5 Azide III-47a (0.126 g, 0.219 mmol) was dissolved in THF (12 ml) and treated with water (0.096 ml, 5.33 mmol) followed by triphenylphosphine (0.114 g, 0.44 mmol). The mixture was then heated at 60°C for 12 h, cooled, and
10 concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) afforded III-8a (87.3 mg, 73% yield) as a white solid: $[\alpha]_D^{25} +6.8^\circ$ (c 1.85, CHCl₃); IR (CH₂Cl₂) 3700 (w), 3040 (s), 2980 (s), 2920 (s), 2860 (m), 1420 (s), 1350 (m), 1310 (m), 1260 (s), 1140 (m), 1060 (s), 890 (s), 700
15 (br) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.25 (m, 15 H), 4.92 (d, J = 10.9 Hz, 1 H), 4.91 (d, J = 11.0 Hz, 1 H), 4.85 (d, J = 10.9 Hz, 1 H), 4.78 (d, J = 11.0 Hz, 1 H), 4.70 (d, J = 10.9 Hz, 1 H), 4.61 (d, J = 10.9 Hz, 1 H), 4.29 (d, J = 7.8 Hz, 1 H), 3.70-3.40 (m, 8 H), 3.56 (s, 3 H), 2.66 (t, J = 6.9
20 Hz, 2 H), 1.61-1.56 (m, 4 H), 1.46-1.35 (m, 4 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.58, 138.52, 138.25, 128.39, 128.31, 128.05, 127.85, 127.84, 127.73, 127.58, 127.55, 104.71, 84.61, 82.30, 77.94, 75.65, 74.95, 74.83, 74.70, 71.63, 69.61, 57.07, 42.02, 33.47, 29.48, 23.45; high resolution mass spectrum (FAB, m-nitrobenzyl alcohol) m/z 572.2997 [(M +Na)⁺; calcd for C₃₃H₄₃O₆N: 572.2988].
25

AV. Methyl 2,3,4-Tri-O-benzyl-6-amino-6-deoxy-6-N-(5-hydroxypentyl)- β -D-glucopyranoside (III-8b).

A stirred solution of III-46 (800 mg, 1.71 mmol) and
30 2,6-di-tert-butyl-4-methyl pyridine (632 mg, 3.08 mmol) in dichloromethane (9 ml) was cooled to -78°C and treated with triflic anhydride (0.345 ml, 2.05 mmol). After 15 min the mixture was warmed to room temperature over 20 min, poured into saturated aqueous sodium bicarbonate (20 ml), and
35 extracted with ethyl acetate (50 ml). The organic layer was washed with additional bicarbonate solution and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo,

- 153 -

affording crude triflate which was used in the next step without further purification.

A solution of 5-trifluoroacetamido-1-pentanol (III-18a) (1.7 g, 8.6 mmol) in THF (35 ml) was added to a stirred suspension of sodium hydride (855 mg, 21.4 mmol, 60% oil dispersion) in THF (60 ml) at 0°C. After 10 min the suspension was warmed to room temperature, stirred for 1 h, and recooled to 0°C. A solution of the crude triflate (1.71 mmol) in dichloromethane (60 ml) was then added and stirring continued at 0°C for 30 min and at room temperature for 24 h. The reaction mixture was quenched at 0°C with saturated aqueous ammonium chloride and extracted with ethyl acetate, and the combined organic extracts were washed with water and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Purification through a small plug of silica gel (30% ethyl acetate/petroleum ether) gave crude III-47b which was used immediately in the next step.

A stirred solution of the above crude III-47b in ethanol (10 ml) was treated with 5 N NaOH (3 ml, 15 mmol) at room temperature and then heated at reflux for 2 h, cooled, and concentrated in vacuo. The residue was diluted with dichloromethane and washed with 2 N HCl. The aqueous layer was extracted with dichloromethane (3 x 50 ml), and the combined organic solutions were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Recrystallization (ethyl acetate/petroleum ether) furnished pure III-8b (675 mg, 72% yield from 46) as a white solid: mp 95-95.5°C; $[\alpha]_D^{25} +9.3^\circ$ (c 0.15, acetonitrile); IR (film) 3280 (m), 3095 (w), 3065 (w), 3035 (w), 2935 (s), 2915 (s), 2860 (s), 1496 (w), 1454 (m), 1404 (w), 1393 (w), 1358 (m), 1214 (m), 1115 (s), 1072 (s), 1037 (m), 1027 (m), 1009 (m), 911 (w), 826 (w), 747 (s), 696 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.35-7.24 (m, 1.5 H), 4.92 (d, $J = 7.5$ Hz, 1 H), 4.90 (d, $J = 7.6$ Hz, 1 H), 4.85 (d, $J = 11.0$ Hz, 1 H), 4.78 (d, $J = 11.0$ Hz, 1 H), 4.70 (d, $J = 11.0$ Hz, 1 H), 4.60 (d, $J = 11.0$ Hz, 1 H), 4.32 (d, $J = 7.8$ Hz, 1 H), 3.66-3.59 (m, 3 H), 3.56 (s, 3 H), 3.48-3.36 (m, 3 H), 2.94 (dd, $J = 12.5, 2.1$ Hz, 1 H),

- 154 -

2.68 (dd, $J = 12.0, 6.8$ Hz, 1 H), 2.64-2.53 (m, 2 H), 1.71 (s, 2 H), 1.59-1.53 (m, 2 H), 1.51-1.45 (m, 2 H), 1.42-1.36 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.55, 138.47, 138.17, 128.39, 128.33, 128.03, 127.95, 127.85, 127.77, 127.60, 127.57, 104.72, 84.56, 82.45, 79.74, 75.66, 75.02, 74.74, 74.16, 62.62, 57.20, 50.69, 49.72, 32.49, 29.65, 23.37; high resolution mass spectrum (Cl , NH_3) m/z 550.3179 $[(\text{M}+\text{H})^+]$; calcd for $\text{C}_{33}\text{H}_{43}\text{O}_6\text{N}$: 550.3168].

AW. 2-(1H-Indol-3-yl)ethyl 2,3,4-Tri-O-benzyl- β -D-glucopyranoside (III-9).

10

A stirred solution of III-17 (100 mg, 0.136 mmol) in ethanol (3 ml) was treated with 5 N NaOH (1 ml) and then heated at reflux for 2 h, cooled, and concentrated in vacuo. The residue was diluted with dichloromethane and washed with 2 N HCl, and the aqueous layer was extracted with dichloromethane. The combined organic solutions were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography (25% ethyl acetate/petroleum ether) furnished III-9 (68 mg, 85% yield) as a colorless oil: $[\alpha]_{\text{D}}^{25} -2.5^\circ$ (c 1.37, acetonitrile); UV (2.89 $\times 10^{-4}$ M, acetonitrile) λ_{max} 289.6 (ϵ 3.56 $\times 10^3$), 281.2 (4.24 $\times 10^3$), 222.4 (1.01 $\times 10^4$) nm; IR (film) 3575 (sh), 3435 (m), 3085 (sh), 3065 (w), 3035 (w), 2925 (m), 2880 (m), 1500 (w), 1455 (m), 1360 (w), 1310 (w), 1150 (sh), 1085 (s), 1030 (s), 920 (w), 810 (w), 740 (s), 700 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.83 (s, 1 H), 7.59 (d, $J = 7.8$ Hz, 1 H), 7.33-7.24 (m, 1.5 H), 7.20-7.17 (m, 2 H), 7.11 (t, $J = 7.8$ Hz, 1 H), 7.01 (d, $J = 1.8$ Hz, 1 H), 4.91 (d, $J = 10.9$ Hz, 1 H), 4.85 (d, $J = 10.9$ Hz, 1 H), 4.80 (d, $J = 10.9$ Hz, 1 H), 4.79 (d, $J = 11.0$ Hz, 1 H), 4.64 (d, $J = 11.0$ Hz, 1 H), 4.63 (d, $J = 11.0$ Hz, 1 H), 4.49 (d, $J = 7.8$ Hz, 1 H), 4.22 (ddd, $J = 9.4, 6.7, 6.7$ Hz, 1 H), 3.90-3.82 (m, 2 H), 3.72-3.67 (m, 1 H), 3.65 (apparent t, $J = 9.1$ Hz, 1 H), 3.56 (apparent t, $J = 9.3$ Hz, 1 H), 3.42 (apparent t, $J = 8.1$ Hz, 1 H), 3.35 (ddd, $J = 9.5, 4.3, 2.8$ Hz, 1 H), 3.11 (t, $J = 7.0$ Hz, 2 H), 1.87 (dd, $J = 7.6, 5.9$ Hz, 1 H); ^{13}C NMR (500 MHz, CDCl_3) δ 138.52, 138.44, 137.98, 136.17, 128.46, 128.36, 128.29, 128.05,

35

- 155 -

128.00, 127.89, 127.86, 127.60, 127.57, 127.45, 122.09, 122.01, 119.34, 118.68, 112.60, 111.13, 103.69, 84.49, 82.34, 77.57, 75.64, 75.04, 75.01, 74.75, 70.25, 62.04, 25.86; high resolution mass spectrum (Cl, NH₃) m/z 611.3043 [(M+NH₄)⁺; calcd for C₃₇H₃₉O₆N: 611.3121].

AX. Methyl 2,3-Di-O-benzyl-4,6-di-O-isopropylidene- β -D-glucopyranoside (III-50).

A solution of glucoside III-49 (2.5 g, 10.7 mmol) in THF (100 ml) was added to a suspension of sodium hydride (0.94 g, 23.5 mmol) in THF (50 ml) at 0°C. The reaction was stirred at room temperature for 1 h and cooled to 0°C, and benzyl bromide (2.8 ml, 24 mmol) was then added dropwise, followed by tetrabutylammonium iodide (100 mg). The mixture was stirred at room temperature for 24 h, quenched with saturated aqueous ammonium chloride, extracted with ether, and the extracts washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% ethyl acetate/petroleum ether) afforded III-50 as a colorless oil (4.02 g, 91% yield): $[\alpha]_D^{25}$ -2.0° (c 0.15, acetonitrile); UV (6.01 x 10⁻⁴ M, acetonitrile) λ_{max} 257.6 (ε 508) nm; IR (film) 3060 (m), 3000 (m), 2980 (m), 2900 (m), 1460 (m), 1390 (m), 1380 (m), 1310 (w), 1270 (s), 1210 (m), 1180 (m), 1100 (s), 1080 (s), 1050 (m), 1030 (m), 860 (m), 740 (s), 705 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.23 (m, 10 H), 4.84 (d, J = 11.3 Hz, 2 H), 4.74 (dd, J = 11.4, 9.2 Hz, 1 H), 4.36 (d, J = 7.6 Hz, 1 H), 3.93 (dd, J = 10.8, 5.4 Hz, 1 H), 3.76 (t, J = 10.5 Hz, 1 H), 3.69 (t, J = 9.3 Hz, 1 H), 3.57 (m, 4 H), 3.37 (t, J = 8.3 Hz, 1 H), 3.23 (m, 1 H), 1.48 (s, 3 H), 1.42 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.81, 138.54, 128.26, 128.17, 127.98, 127.85, 127.58, 127.46, 105.16, 99.24, 82.14, 81.27, 75.19, 74.77, 74.27, 69.79, 62.25, 57.32, 29.14, 19.09; high resolution mass spectrum (Cl, NH₃) m/z 415.2137 [(M+H)⁺; calcd for C₂₄H₃₁O₆: 415.2120].

AY. Methyl 2,3-Di-O-benzyl- β -D-glucopyranoside (III-51).

Amberlyst® 15 ion exchange resin (0.5 g) was added to a solution of III-50 (1.00 g, 2.4 mmol) in methanol (50 ml)

- 156 -

and the mixture was stirred at room temperature for 4 h, filtered, and concentrated in vacuo. Flash chromatography (6% methanol/dichloromethane) yielded III-51 (0.75 g, 83% yield) as a white foam: $[\alpha]_D^{25} +16^\circ$ (c 0.15, acetonitrile); UV (2.00 x 10⁻⁴ M, acetonitrile) λ_{\max} 257.6 (ϵ 385.0) nm; IR (film) 3590 (w), 3410 (br), 3080 (m), 2910 (w), 2890 (w), 1500 (w), 1455 (m), 1270 (s), 1210 (w), 1065 (s), 1030 (s), 1000 (m), 900 (m), 740 (s), 700 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.34-7.23 (m, 10 H), 4.91 (dd, J = 15.6, 11.5 Hz, 2 H), 4.69 (dd, J = 11.5, 8.7 Hz, 2 H), 4.34 (d, J = 7.7 Hz, 1 H), 3.87-3.83 (m, 1 H), 3.77-3.72 (m, 1 H), 3.58-3.54 (m, 4 H), 3.44 (t, J = 9.1 Hz, 1 H), 3.37 (t, J = 7.6 Hz, 1 H), 3.31-3.27 (m, 1 H), 2.84 (br s, 1 H), 2.48 (br s, 1 H); ¹³C NMR (125 MHz, CDCl₃) δ 138.48, 138.34, 128.49, 128.46, 128.30, 127.99, 127.83, 127.76, 127.62, 104.85, 83.82, 81.87, 75.13, 74.90, 74.57, 70.18, 62.30, 57.20; high resolution mass spectrum (Cl, NH₃) m/z 392.2043 [(M+NH₄)⁺; calcd for C₂₁H₃₀NO₆: 392.2072].

20 **BA. Methyl 2,3-Di-O-benzyl-6-O-tert-butyldiphenylsilyl- β -D-glucopyranoside (III-52).**

A solution of III-51 (3.30 g, 8.81 mmol) and imidazole (0.84 g, 12.3 mmol) in a mixture of THF (150 ml) and DMF (25 ml) was treated with tert-butyldiphenylsilyl chloride (2.80 ml, 10.6 mmol) and heated at 50°C for 24 h. The reaction mixture was quenched with methanol (5 ml) and concentrated in vacuo. The resultant oil was taken up in ethyl acetate and the solution was washed with water and brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% ethyl acetate/petroleum ether) furnished III-52 (5.40 g, 100% yield) as a colorless oil: $[\alpha]_D^{25} +7.3^\circ$ (c 0.22, acetonitrile); UV (1.79 x 10⁻⁴ M, acetonitrile) λ_{\max} 258.8 (ϵ 836) nm; IR (film) 3500 (br), 3080 (w), 3030 (w), 2940 (m), 2860 (m), 1450 (w), 1430 (m), 1390 (w), 1360 (w), 1310 (w), 1270 (w), 1220 (w), 1190 (w), 1120 (s), 1070 (s), 830 (m), 805 (w), 740 (s), 700 (s) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.71-7.69 (m, 4 H), 7.42-7.25 (m, 16 H), 4.93 (d, J = 11.5 Hz, 2 H), 4.76 (d, J = 11.4 Hz, 1 H), 4.71 (d, J = 11.1

- 157 -

Hz, 1 H), 4.32 (d, $J = 7.6$ Hz, 1 H), 3.94-3.88 (m, 2 H), 3.69-3.64 (m, 1 H), 3.66 (s, 3 H), 3.47 (t, $J = 9.1$ Hz, 1 H), 3.41-3.34 (m, 2 H), 2.57 (br s, 1 H), 1.06 (s, 9 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.71, 138.62, 135.69, 135.61, 129.73, 128.50, 128.34, 128.03, 127.99, 127.78, 127.72, 127.69, 127.62, 104.68, 84.22, 81.93, 75.30, 74.89, 74.67, 71.62, 64.44, 56.86, 26.79, 19.25; high resolution mass spectrum (Cl_2 , NH_3) m/z 630.3296 [$(\text{M}+\text{NH}_4)^+$; calcd for $\text{C}_{37}\text{H}_{48}\text{NO}_6\text{Si}$: 630.3251].

10 **BB. Methyl 2,3-Di-O-benzyl-4-deoxy-6-O-tert-butyl**
diphenylsilyl- β -D-glucopyranoside (III-53).

A solution of III-52 (0.33 g, 0.54 mmol) in THF (20 ml) was cooled to -78°C and treated with sodium bis(trimethylsilyl)amide (0.66 ml, 1.0 M in THF, 0.66 mmol) followed by carbon disulfide (46 ml, 0.77 mmol). After 15 min, methyl iodide (137 ml, 2.20 mmol) was added, and the solution was stirred 15 min further at -78°C and then at room temperature for 45 min. The reaction mixture was quenched with water (2 ml) and extracted with ether. The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo affording the crude xanthate as a yellow oil which was used without purification.

A solution of crude xanthate (6.06 g, 8.62 mmol) and a catalytic amount of AIBN (ca. 50 mg) in toluene (350 ml) was treated with tributyltin hydride (7.0 ml, 26 mmol) and then heated at reflux for 3 h, cooled, and concentrated in vacuo. The residue was taken up in acetonitrile and extracted with petroleum ether (5 x 100 ml). The acetonitrile layer was dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (8% ethyl acetate/petroleum ether) yielded III-53 (3.60 g, 78% yield for two steps) as a colorless oil: $[\alpha]_{\text{D}}^{25} +2.7^\circ$ (c 0.15, acetonitrile); UV (1.26×10^{-4} M, acetonitrile) λ_{max} 258.4 (ϵ 976) nm; IR (film) 3080 (m), 2990 (w), 2880 (w), 1430 (w), 1270 (s), 1110 (m), 900 (w), 740 (s), 710 (s) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.68-7.66 (m, 4 H), 7.43-7.21 (m, 16 H), 4.89 (d, $J = 11.1$ Hz, 1 H), 4.75 (d, $J = 10.2$ Hz, 1 H), 4.67 (dd, $J = 18.2, 11.9$ Hz, 2 H), 4.22 (d, $J = 7.6$ Hz, 1 H), 3.80 (dd, $J = 10.5, 5.7$ Hz, 1 H),

- 158 -

3.63 (dd, $J = 10.4, 5.4$ Hz, 1 H), 3.59-3.44 (m, 5 H), 3.29 (t, $J = 8.9$ Hz, 1 H), 2.11 (ddd, $J = 12.8, 5.1, 1.5$ Hz, 1 H), 1.41 (q, $J = 11.8$ Hz, 1 H), 1.06 (s, 9 H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.91, 138.65, 135.60, 135.55, 133.48, 133.44, 129.66, 128.29, 128.22, 127.95, 127.63, 127.60, 127.49, 127.43, 104.81, 82.99, 78.32, 74.82, 72.20, 72.15, 66.22, 56.73, 33.62, 26.78, 19.22; high resolution mass spectrum (Cl, NH_3) m/z 614.3256 [(M+ NH_4) $^+$; calcd for $\text{C}_{17}\text{H}_{18}\text{NO}_5\text{Si}$: 614.3301].

10 **BC. Methyl 2,3-Di-O-benzyl-4-deoxy- β -D-glucopyranoside (III-54).**

A solution of III-53 (3.60 g, 6.02 mmol) in THF (125 ml) was treated with tetrabutylammonium fluoride (1.0 M in THF, 6.1 mmol, 6.1 ml) at room temperature, stirred for 4 h, poured into water, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (50% ethyl acetate/petroleum ether) afforded III-54 (2.03 g, 94% yield) as a colorless oil: $[\alpha]_D^{25} +8.0^\circ$ (c 0.15, acetonitrile); UV (2.09 $\times 10^{-4}$ M, acetonitrile) λ_{max} 257.6 (ϵ 177) nm; IR (film) 3450 (br), 3095 (w), 3060 (w), 3030 (w), 2920 (m), 2880 (m), 1500 (w), 1450 (m), 1380 (m), 1360 (m), 1300 (w), 1260 (w), 1210 (m), 1180 (w), 1070 (br), 910 (m), 740 (m), 700 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.38-7.25 (m, 10 H), 4.89 (d, $J = 11.1$ Hz, 1 H), 4.76 (d, $J = 11.1$ Hz, 1 H), 4.67 (m, 2 H), 4.28 (d, $J = 7.7$ Hz, 1 H), 3.73-3.49 (m, 7 H), 3.29 (t, $J = 7.9$ Hz, 1 H), 2.08 (br s, 1 H), 1.97 (ddd, $J = 12.9, 5.3, 1.9$ Hz, 1 H), 1.49 (dd, $J = 24.4, 11.7$ Hz, 1 H); ^{13}C NMR (500 MHz, CDCl_3) δ 138.71, 138.47, 128.33, 128.28, 128.00, 127.62, 127.58, 127.55, 104.95, 82.81, 78.07, 74.92, 72.26, 72.13, 65.20, 57.19, 32.65; high resolution mass spectrum (Cl, NH_3) m/z 359.1827 [(M+H) $^+$; calcd for $\text{C}_{21}\text{H}_{27}\text{O}_5$: 359.1858].

BD. Methyl 2,3-Di-O-benzyl-4-deoxy-6-O-(5-phthalimidopentyl)- β -D-glucopyranoside (III-55).

35 A solution of 5-phthalimido-1-pentanol (0.66 g, 2.83 mmol) and 2,6-di-tert-butyl-4-methylpyridine (0.58 g, 2.83 mmol) in dry dichloromethane (21 ml) was treated with triflic

- 159 -

anhydride (0.48 ml, 2.83 mmol) at room temperature, stirred for 10 min, poured into water, and extracted with dichloromethane. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. The freshly
5 generated triflate was then dissolved in dry dichloromethane (21 ml), 2,6-di-tert-butyl-4-methylpyridine (0.58 g, 2.83 mmol) was added, and the solution was cooled to 0°C. A solution of III-54 (1.0 g, 2.79 mmol) in dichloromethane (21 ml) was introduced, followed after 20 min by sodium hydride
10 (60% oil dispersion, 0.25 g, 6.25 mmol). The reaction mixture was stirred at room temperature for 24 h, quenched with saturated aqueous ammonium chloride, extracted with dichloromethane, and the combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated
15 in vacuo. Flash chromatography (50% ethyl acetate/petroleum ether) yielded III-55 (1.42 g, 89% yield) as a colorless oil: $[\alpha]_D^{25} +11^\circ$ (c 0.11, acetonitrile); UV (9.60×10^{-5} M, acetonitrile) λ_{\max} 290.8 (ϵ 3.20×10^3), 257.6 (2.20×10^3), 241.2 (1.69×10^4) nm; IR (film) 3480 (br), 3090 (w), 3040 (w), 3010 (w), 2940 (s), 2860 (s), 2250 (m), 1770 (m), 1715 (s), 1500 (w), 1470 (m), 1450 (m), 1430 (m), 1400 (s), 1370 (m), 1340 (w), 1300 (w), 1260 (w), 1210 (m), 1190 (m), 1170 (w), 1100 (br), 1000 (w), 910 (s), 730 (s), 720 (s), 700 (s),
20 650 (m) cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.84-7.68 (m, 4 H), 7.38-7.24 (m, 10 H), 4.88 (d, J = 11.1 Hz, 1 H), 4.75 (d, J = 11.1 Hz, 1 H), 4.67 (s, 2 H), 4.24 (d, J = 7.6 Hz, 1 H), 3.68 (t, J = 7.3 Hz, 2 H), 3.61-3.41 (m, 9 H), 3.28 (t, J = 8.5 Hz, 1 H), 2.10 (dd, J = 12.6, 5.3 Hz, 1 H), 1.73-1.59 (m, 5 H), 1.45-1.37 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 168.38,
25 138.87, 138.64, 133.84, 132.17, 128.30, 128.25, 128.02, 127.63, 127.48, 123.14, 104.85, 82.91, 78.24, 74.86, 73.15, 72.19, 71.42, 70.92, 56.97, 37.90, 33.94, 29.15, 28.38, 23.44; high resolution mass spectrum (Cl , NH_3) m/z 591.3014 [$(\text{M}+\text{NH}_4)^+$; calcd for $\text{C}_{34}\text{H}_{43}\text{O}_7\text{N}_2$: 591.3070].
30

- 160 -

BE. Methyl 2,3-Di-O-benzyl-4-deoxy-6-O-(5-amino-pentyl)- β -D-glucopyranoside (III-10).

A solution of phthalimide III-55 (0.79 g, 1.38 mmol) in methanol (100 ml) was treated with sodium methoxide (0.23 g, 4.26 mmol), heated at reflux for 4h, cooled, and concentrated in vacuo. The residue was taken up in water and extracted with dichloromethane, and the organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Flash chromatography (10% methanol/dichloromethane) furnished III-10 (0.46 g, 75% yield) as a white foam: $[\alpha]_D^{25} +8.9^\circ$ (c 0.18, acetonitrile); UV (2.03 x 10⁻⁴ M, acetonitrile) λ_{max} 276.4 (ϵ 1.54 x 10³), 257.6 (2.26 x 10³) nm; IR (film) 3330 (br), 3080 (w), 3020 (w), 2930 (s), 2870 (s), 1650 (s), 1550 (m), 1450 (m), 1370 (m), 1300 (s), 1210 (m), 1185 (m), 1100 (br), 1000 (w), 900 (w), 740 (s), 700 (s), 670 (w), 640 (w) cm⁻¹; ¹H NMR (500 MHz, CD₃OD) δ 7.76-7.74 (m, 1 H), 7.35-7.13 (m, 9 H), 4.74-4.49 (m, 4 H), 4.14 (d, J = 7.7 Hz, 1 H), 3.53-3.36 (m, 9 H), 3.20 (m, 2 H), 3.07 (t, J = 7.8 Hz, 1 H), 1.99 (ddd, J = 2.8, 5.3, 1.7 Hz, 1 H), 1.50-1.18 (m, 9 H); ¹³C NMR (125 MHz, CD₃OD) δ 140.18, 139.99, 138.83, 131.94, 130.83, 130.49, 129.28, 129.18, 128.99, 128.83, 128.69, 128.55, 128.49, 105.98, 84.04, 79.56, 75.73, 74.05, 73.04, 72.49, 72.12, 57.24, 40.94, 34.53, 30.30, 29.81, 24.64; high resolution mass spectrum (CI, NH₃) m/z 444.2783 [(M+H)⁺; calcd for C₂₅H₃₈NO₅: 444.2749].

EXAMPLE 12

Synthesis of H-Phe-Thr(t-Bu)-Xaa-D-Trp-Phe-Pro-2-chlorotrityl-Resin.

Assembly of multiple peptides on a single solid support was carried out using an Applied Biosystems, Inc. Model 431A automated peptide synthesizer. N- α -Fmoc amino acids were employed throughout, with appropriately protected side chain, from Bachem, Inc. Starting from 0.25 mmol of Fmoc-L-Pro-2-chlorotrityl polystyrene resin (0.44g, 0.57 mmol/g), the H-Phe-Thr(t-Bu)-Xaa-D-Trp-Phe-Pro-2-chlorotrityl-Resin was assembled generally according to standard

- 161 -

procedures. At the fixed positions, a four molar excess (1.0 mmol) individual amino acid along with 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethylammonium hexafluorophosphate (HBTU) was used in the coupling step, and the coupling reaction was carried out at room temperature for 2.0 h. At the mixed position, a total combined 1.0 mmol Fmoc-Xaa-OH mixtures [Xaa: Ala (0.13 mmol), Leu (0.28 mmol), Phe (0.12 mmol), Tyr(t-Bu) (0.48 mmol)] with molar ratio adjusted to compensate the reactivity difference according to Houghten's procedure was used. (See, Eichler, et al., *Biochemistry* 1993, 32, 11035 Thus, incorporated in order, were: Fmoc-Phe-OH, Fmoc-Thr(tBu)-OH, Fmoc-Xaa-OH (Xaa = Ala, Leu, Phe, Tyr(tBu), Fmoc-D-Trp-OH, Fmoc-Phe-OH. After each coupling, a Kaiser test was performed to monitor the coupling reaction and, if necessary, a double coupling reaction was performed. The N- α -Fmoc group was removed at the end of the synthesis. Methylene chloride washing was avoided through out the whole washing step. After the completion of synthesis, resin was dried under vacuum to afford 686.0 mg peptide resin.

20 EXAMPLE 13

Synthesis of Cyclo-(pro-Phe-Thr-Xaa-D-trp-Phe) (Xi: Ala, Leu, Phe, Tyr)

Peptide resin H-Phe-D-Trp-Xaa-Thr(tBu)-Phe-Pro-Resin (665 mg) [Xaa: Ala, Leu, Phe, Tyr(tBu)] was treated with 15 ml of 0.25% trifluoroacetic acid (TFA) in CH₂Cl₂ at room temperature. After 30 min, the slurry was filtered, washed with 0.25 % TFA in CH₂Cl₂ solution, the filtrate was evaporated at room temperature and the residue was titrated with ice-cold dry diethyl ether, filtered and washed with ether to provide a 247 mg white powder which was subjected to the next cyclization without further purification.

To a suspension of 200 mg H-Phe-D-Trp-Nleu-Thr(tBu)-Phe-Pro-OH and 251 mg of solid NaHCO₃ in 33 ml dry dimethylformamide (DMF), 65 l diphenylphosphoryl azide (DPPA) was added dropwise at 0°C. The reaction mixture was then stirred at 4°C. The cyclization was completed after 21 h as

- 162 -

indicated by analytical RP-HPLC. The reaction mixture was concentrated under reduced pressure to remove DMF, the residue was redissolved in 50% CH₃CN in water and lyophilized to afford 212.0 mg white powder.

5 To half of the above material (106 mg) dissolved in 3.4 mL CH₂Cl₂, 225 mL ethanedithiol (EDT) and 150 mL H₂O. TFA (3.75 mL) was added at room temperature dropwise. After stirring at room temperature for 50 min., the reaction mixture was concentrated to half its volume and flushed with dry
10 benzene (3 x 10 mL). The residue was precipitated with dry ether, filtered and washed extensively with ether and purified by RP-HPLC (C18 Dynamax 300 (21.4 x 250 mm) column, gradient 35-25'-95%B, flow rate = 12 mL/min. to afford the pure compounds (17a) (23 mg), (17b) (14 mg), (17c) (13 mg) and
15 (17d) (14.4 mg) in a combined yield of 82% from Fmoc-Pro-2-chlorotrityl polystyrene resin.

Cyclo (Phe-D-Trp-Leu-Thr-Phe-Pro) (17a) [a]²⁵_D = -121.78 (C = 0.28, CH₃OH); ¹H NMR (500 MHz, CD₃OD) δ 0.56 (d, J=6.35 Hz, 3H), 0.66 (d, J=6.44 Hz, 3H), 0.75-0.76 (m, 1H), 0.88-0.99 (m, 1H), 1.01-1.09 (m, 1H), 1.16 (d, J=6.37 Hz, 3H), 1.30-1.42 (m, 2H), 1.47-1.53 (m, 1H), 1.79-1.96 (m, 1H), 2.85 (dd, J=5.12, 13.5 Hz, 1H), 2.90-3.11 (m, 5H), 3.17-3.21 (m, 1H), 3.26-3.30 (m, 1H), 3.65 (d, J=7.78 Hz, 1H), 3.87-3.92 (m, 1H), 4.11-4.16 (m, 1H), 4.37-4.41 (m, 2H), 4.60-4.67 (m, 1H), 4.72-4.75 (m, 1H), 7.0-7.36 (m, 15H), 7.58 (d, J=7.78 Hz, 1H); ¹³C NMR (125 MHz, CD₃OD) δ 17.56, 19.69, 20.88, 22.01, 23.97, 27.13, 30.26, 37.31, 38.14, 39.66, 46.03, 53.46, 53.99, 54.74, 54.77, 56.21, 61.22, 67.14, 108.92, 111.02, 117.99, 118.52, 121.14, 123.26, 126.47, 127.05, 127.25, 128.09, 128.62, 129.10, 129.28, 135.51, 136.85, 170.30, 170.45, 171.16, 171.77, 173.19, 173.85; HR-FAB-MS m/z 814.3903 (M+Na calcd for C₄₄H₅₃N₇O₇, 814.3904).

Cyclo (Phe-D-Trp-Phe-Thr-Phe-Pro) (17b) [a]²⁵_D = -67.09 (C = 0.31, CH₃OH); ¹H NMR (500 MHz, CD₃OD, 315K) δ 0.78-0.86 (m 1), 0.96-1.05 (m, 1H), 1.13 (d, J=6.39 Hz, 3H), 1.43-1.47 (m, 1H), 1.75 (dd, J=6.36, 12.26 Hz, 1H), 2.79 (dd, J=6.31, 13.87 Hz, 1H), 2.85 (d, J=6.93 Hz, 2H), 2.86-2.95 (m,

- 163 -

3H), 3.02 (dd, J=6.62, 13.55 Hz, 1H), 3.07 (dd, J=5.59, 12.78 Hz, 1H), 3.11-3.14 (m, 1H), 3.19-3.25 (m, 1H), 3.63 (d, J=7.76 Hz, 1H), 4.07-4.12 (m, 1H), 4.26-4.29 (m, 1H), 4.38-4.41 (m, 2H), 4.48 (dd, J=6.44, 8.99 Hz, 1H), 4.64-4.67 (m, 1H),
 5 6.85-6.86 (m, 2H), 6.89 (s, 1H), 6.98-7.00 (m, 2H), 7.03-7.06 (m, 1H), 7.12-7.15 (m, 7H), 7.22-7.23 (m, 2H), 7.25-7.28 (m, 2H), 7.30-7.36 (m, 3H), 7.49 (d, J=7.88 Hz, 1H); ¹³C NMR (125 MHz, CD₃OD) δ 18.89, 22.23, 28.40, 31.59, 37.53, 38.73, 39.45, 47.37, 55.30, 55.97, 57.19, 57.87, 57.95, 62.60, 68.49,
 10 110.46, 112.57, 119.30, 119.87, 122.54, 124.51, 127.81, 128.48, 128.62, 129.39, 129.57, 129.84, 130.01, 130.42, 130.64, 136.93, 137.83, 138.07, 138.13, 171.50, 172.03, 172.49, 173.19, 173.68, 174.44. HR-FAB-MS m/z 848.3731 (M+Na calcd for C₄₇H₅₁N₇O₇, 848.3747)

15 Cyclo (Phe-D-Trp-Tyr-Thr-Phe-Pro) (17c) [α]^{Fo}(25,D) = -71.53 (C = 0.26, CH₃OH); ¹H NMR (500 MHz, CD₃OD) δ 0.80-0.90 (m, 1H), 1.0-1.13 (m, 1H), 1.12 (d, J=6.08 Hz, 3H), 1.44-1.48 (m, 1H), 1.74-1.75 (m, 1H), 2.64-2.67 (m, 1H), 2.77-2.88 (m, 2H), 2.91-3.08 (m, 6H), 3.12-3.16 (m, 1H), 3.22-3.26 (m, 1H),
 20 3.63-3.65 (m, 1H), 4.09-4.10 (m, 1H), 4.15-4.22 (m, 1H), 4.32-4.38 (m, 2H), 4.47-4.49 (m, 1H), 6.58-6.60 (m, 1H), 6.65-6.69 (m, 1H), 6.92 (m, 1H), 7.01-7.08 (m, 3H), 7.10-7.20 (m, 3H), 7.21-7.27 (m, 2H), 7.28-7.32 (m, 3H), 7.37-7.39 (m, 1H), 7.52 (d, J=7.64 Hz, 1H); ¹³C NMR (125 MHz, CD₃OD) δ
 25 18.83, 22.23, 28.40, 31.63, 36.71, 38.72, 39.48, 47.37, 55.27, 56.07, 57.36, 57.82, 57.90, 62.60, 68.48, 110.52, 112.59, 116.45, 119.29, 119.93, 122.59, 124.53, 127.79, 128.36, 128.46, 128.61, 129.38, 129.99, 130.43, 130.63, 130.93, 136.91, 138.04, 138.14, 157.27, 171.46, 172.02, 172.57,
 30 173.18, 173.79, 174.38. HR-FAB-MS m/z 864.3713 (M+Na calcd for C₄₇H₅₁N₇O₈, 864.3696).

Cyclo (Phe-D-Trp-Ala-Thr-Phe-Pro) (17d) [α]^{Fo}(25,D) = -78.5 (C = 0.475, CH₃OH); ¹H NMR (500 MHz, CD₃OD) δ 0.82-0.85 (m, 1H), 0.90-0.96 (m, 1H), 1.12 (d, J=7.34 Hz, 3H),
 35 1.16 (d, J=6.34 Hz, 3H), 1.41-1.42 (m, 1H), 1.74-1.78 (m, 1H), 2.88-2.96 (m, 3H), 3.05-3.13 (m, 4H), 3.20-3.26 (m, 2H), 3.63-3.64 (m, 1H), 3.91-3.94 (m, 1H), 4.11-4.13 (m, 1H),

- 164 -

4.37-4.40 (m, 2H), 4.42-4.50 (m, 1H), 7.02-7.09 (m, 4H),
7.11-7.19 (m, 4H), 7.23-7.35 (m, 7H), 7.58 (d, J=7.84 Hz, 1H);
13C NMR (125 MHz, CD₃OD) d 16.81, 19.06, 22.18, 28.42, 31.47,
38.72, 39.19, 47.35, 52.36, 55.37, 55.97, 56.21, 57.76, 62.61,
5 68.47, 110.72, 112.34, 119.38, 119.82, 122.51, 124.62, 127.78,
128.63, 128.65, 129.42, 130.00, 130.39, 130.67, 136.91,
138.12, 138.24, 171.73, 172.04, 172.53, 173.13, 174.59,
175.48. HR-FAB-MS m/z 772.3405 (M+Na calcd for C₄₁H₄₇N₇O₇,
772.3434).

10 EXAMPLE 14

Synthesis of Cyclo (Phe-D-Trp-p-MeO-Phe-Thr-Phe-Pro) (17e)

To a suspension of 3.3 mg (17c) and 1.6 mg K₂CO₃ in 0.1 mL DMF, 3.6 mL CH₃I was added. After 1.0 h, the reaction mixture was filtered and dissolved in 50% CH₃CN in water and
15 lyophilized to afford a solid which was purified by RP-HPLC to afford 3.1 mg 6e in 65%. 1H NMR (500 MHz, CD₃OD) d 0.79-0.84 (m, 1H), 0.95-1.05 (m, 1H), 1.14 (d, J=6.41 Hz, 3H), 1.40-1.46 (m, 1H), 1.76 (dd, J=5.96, 12.15 Hz, 1H), 2.71 (dd, J=4.9, 14.034 Hz, 1H), 2.78-2.84 (m, 2H), 2.87-2.98 (m, 3H),
20 3.01-3.14 (m, 3H), 3.20-3.26 (m, 1H), 3.64 (d, J=7.95 Hz, 1H), 4.07-4.09 (m, 1H), 4.11-4.22 (m, 1H), 4.38-4.40 (m, 2H), 4.41-4.48 (m, 1H), 4.64-4.67 (m, 1H), 6.6 (d, J=8.71 Hz, 2H), 6.72 (d, J=8.63 Hz, 2H), 6.95 (s, 1H), 7.01-7.08 (m, 3H), 7.13-7.18 (m, 4H), 7.23 (d, J=6.97 Hz, 2H), 7.26-7.38 (m, 4H),
25 7.51 (d, J=7.85 Hz, 1H), 7.99 (d, J=6.81 Hz, 1H), 8.19 (d, J=6.89 Hz, 1H), 8.39 (bs, 1H). HR-FAB-MS m/z 878.3831 (M+Na calcd for C₄₈H₅₃N₇O₈, 878.3853, 2.5 ppm err).

EXAMPLE 15

Synthesis of Cyclo (Phe-D-Trp-Xaa-Thr-Phe-Pro) (Xaa = p-F-Phe,
30 Homo-Phe, Cha, Trp, D-Phe).

Compounds (17f-k) were prepared using the same general procedure as for the synthesis of compounds (17a-d), except that a modified coupling protocol was used at the mixed position. At mixed position, a total combined one equivalent
35 (0.25 mmol) of eq-molar Fmoc-Xaa-OH per one equivalent peptide

- 165 -

resin (0.25 mmol) was used in the coupling reaction along with 2-(1H-aminobenzotriazol-1-yl)-1,1,3,3-tetramethylammonium hexafluorophosphate (HATU) as the coupling reagent. The final crude mixtures were separated with RP-HPLC (C18 Dynamax 300
5 (21.4 x 250 mm) column, gradient 35-25'-95%B, flow rate = 12 mL/min. to afford compounds (17f-k), and a mixture containing two compounds.

Cyclo (Phe-D-Trp-p-F-Phe-Thr-Phe-Pro) (17f)
[a]^{Fo}(25,D) = -74.44 (c = 0.53, CH₃CN); 1H NMR (500 MHz,
10 CD₃OD) δ 0.85-0.91 (m, 1H), 1.04-1.08 (m, 1H), 1.14 (d, J=6.38 Hz, 3H), 1.44-1.47 (m, 1H), 1.75-1.79 (m, 1H), 2.77-2.85 (m, 3H), 2.90-2.99 (m, 3H), 3.02-3.09 (m, 2H), 3.12-3.15 (m, 1H), 3.21-3.27 (m, 1H), 3.33-3.37 (m, 1H), 3.64-d (7.68, 1H), 4.09-4.13 (m, 1H), 4.24 (dd, J=5.35, 8.25 Hz, 1H), 4.37-4.41
15 (m, 2H), 4.49 (dd, J=5.94, 9.50 Hz, 1H), 6.79 (s, 2H), 6.80 (d, J=1.85 Hz, 2H), 6.93 (s, 1H), 7.04-7.10 (m, 4H), 7.13-7.20 (m, 4H), 7.22-7.24 (m, 2H), 7.29-7.38 (m, 4H), 7.50 (d, J=7.88 Hz, 1H); 13C NMR (125 MHz, CD₃OD) δ 18.95, 22.23, 28.47, 31.58, 36.69, 38.73, 39.50, 47.38, 55.31, 56.02, 57.25, 57.88,
20 57.96, 62.96, 68.52, 110.40, 112.60, 116.12 (d, J = 21.7Hz), 119.32, 119.89, 122.54, 124.60, 127.83, 128.47, 128.63, 129.43, 130.01, 130.43, 130.64, 131.39, 131.46, 133.68, 136.91, 138.12, 163.15 (d, J = 243.4 Hz), 171.51, 172.04, 172.50, 173.18, 173.55, 174.40. HR-FAB-MS m/z 866.3631 (M+Na
25 caclcd for C₄₇H₅₀FN₅O₇, 866.3654, 2.5 ppm err).

Cyclo (Phe-D-Trp-Homo-Phe-Thr-Phe-Pro) (17g)
[a]^{Fo}(25,D) = -75.67 (C = 0.6, CH₃CN); 1H NMR (500 MHz,
CD₃OD) δ 0.85-0.91 (m, 1H), 1.0-1.06 (m, 1H), 1.14 (d, J=6.4 Hz, 3H), 1.43-1.50 (m, 1H), 1.58-1.66 (m, 1H), 1.79 (dd,
30 J=6.26, 12.25 Hz, 1H), 1.87-1.98 (m, 3H), 2.86-2.99 (m, 3H), 3.03-3.18 (m, 4H), 3.23-3.29 (m, 1H), 3.63 (d, J=7.75 Hz, 1H), 3.82 (dd, J=3.50, 11.12 Hz, 1H), 4.09-4.14 (m, 1H), 4.36-4.39 (m, 2H), 4.56 (dd, J=5.40, 10.65 Hz, 1H), 4.71-4.76 (m, 1H), 6.86 (d, J=7.08 Hz, 2H), 7.05-7.20 (m, 9H), 7.22-7.26 (m, 4H),
35 7.27-7.37 (m, 4H), 7.60 (d, J=6.92 Hz, 1H); 13C NMR (125 MHz, CD₃OD, 315K) δ 19.04, 22.24, 28.49, 31.60, 32.94, 33.91, 38.69, 39.48, 47.40, 55.39, 55.99, 56.33, 57.59, 57.67, 62.61,

- 166 -

68.50, 110.53, 112.59, 119.46, 119.95, 122.59, 124.70, 126.94, 127.84, 128.60, 128.63, 129.29, 129.39, 129.47, 130.00, 130.47, 130.67, 136.89, 138.21, 138.24, 142.16, 171.66, 171.89, 172.46, 173.14, 174.54, 174.76. (M+Na calcd for

5 $C_{48}H_{61}N_7O_7$, 970.4530, 5 ppm err).

Cyclo (Phe-D-Trp-Cha-Thr-Phe-Pro) (17h) $[\alpha]^{25}_D$

= (c = 0.425, DMSO); 1H NMR (500 MHz, CD_3OD) δ 0.64-0.69

(m, 1H), 0.75 (bs, 2H), 0.83-0.89 (m, 1H), 0.97-1.13 (m, 4H),

1.15 (d, J=6.42 Hz, 3H), 1.30-1.49 (m, 5H), 1.57-1.59 (m, 3H),

10 1.79 (dd, J=6.26, 12.21 Hz, 1H), 2.85 (dd, J=5.28, 13.78 Hz,

1H), 2.90-2.99 (m, 2H), 3.03-3.14 (m, 4H), 3.21-3.27 (m, 1H),

3.64 (d, J=7.66 Hz, 1H), 3.89 (dd, J=3.76, 11.06 Hz, 1H),

4.09-4.13 (m, 1H), 4.40 (dd, J=5.48, 12.38 Hz, 2H), 4.58 (dd,

J=5.40, 10.35 Hz, 1H), 4.64-4.76 (m, 1H), 7.02-7.09 (m, 2H),

15 7.11-7.13 (m, 2H), 7.16-7.19 (m, 1H), 7.21-7.28 (m, 4H),

7.28-7.29 (m, 1H), 7.31-7.35 (m, 4H), 7.58 (d, J=7.83 Hz, 1H);

^{13}C NMR (125 MHz, CD_3OD , major) δ 19.04, 22.22, 26.82, 26.97,

27.48, 28.67, 31.51, 32.62, 34.79, 38.72, 39.34, 39.72, 47.39,

54.43, 55.39, 56.01, 56.17, 57.59, 62.61, 68.48, 110.39,

20 112.56, 119.38, 119.86, 122.49, 124.60, 127.84, 128.53,

128.64, 129.49, 130.00, 130.42, 130.67, 136.89, 138.12,

138.29, 171.79, 171.98, 172.53, 173.12, 174.48, 175.46.

HR-FAB-MS m/z 870.4491 (M+Na calcd for $C_{48}H_{61}N_7O_7$, 970.4530, 5 ppm err).

25 EXAMPLE 16

Synthesis of Cyclo (Yaa-D-Trp-Phe-Thr-Phe-Pro) (18a-i: Yaa = Ser, Ala, Asp, D-Pro, D-Homo-Phe, Cha, Trp, D-Phe, Nal).

Compounds (18a-i) were prepared using the same general procedure as for the synthesis of compounds (17f-k). The

30 final crude mixtures were separated with RP-HPLC (C18 Dynamax 300 (21.4 x 250 mm) column, gradient 35-25'-95% buffer B, flow rate = 12 mL/min. to afford compounds (18a-h), and a mixture containing two compounds which was further resolved by using a C8 Vydac column (10 x 250 mm), gradient 35-25'-95% buffer
35 B, to afford compounds (18i-j).

- 167 -

Cyclo (Ser-D-Trp-Phe-Thr-Phe-Pro) (18a) $[a]^{25}_D$
= -46.05 (C = 0.31, CH₃CN); ¹H NMR (500 MHz, CD₃OD) δ
0.89-0.94 (m, 1H), 1.11 (d, J=6.42 Hz, 3H), 1.56-1.63 (m, 2H),
1.90 (dd, J=6.2, 12.31 Hz, 1H), 2.71 (dd, J=4.98, 14.31 Hz,
5 1H), 2.86 (dd, J=8.53, 14.29 Hz, 1H), 2.92-2.99 (m, 1H),
3.05-3.12 (m, 3H), 3.34-3.43 (m, 2H), 3.65-3.74 (m, 3H),
4.06-4.07 (m, 1H), 4.29 (dd, J=4.94, 8.41 Hz, 1H), 4.37-4.40
(m, 1H), 4.44-4.48 (m, 2H), 4.57 (t, J=7.75 Hz, 1H), 6.77-6.79
(m, 2H), 6.98 (s, 1H), 7.02-7.05 (m, 1H), 7.10-7.16 (m, 4H),
10 7.24-7.38 (m, 6H), 7.54 (d, J=7.91 Hz, 1H);

Cyclo (Asp-D-Trp-Phe-Thr-Phe-Pro) (18b) $[a]^{25}_D$
= -54.19 (C= 0.26, CH₃CN); ¹H NMR (500 MHz, CD₃OD) δ
0.83-0.87 (m, 1H), 1.13 (d, J=6.33 Hz, 3H), 1.54-1.58 (m, 2H),
1.92-1.96 (dd, J=6.07, 12.33 Hz, 1H), 2.64 (dd, J=7.35, 15.93
15 Hz, 1H), 2.77 (dd, J=5.94, 16.00 Hz, 1H), 2.81-2.88 (m, 2H),
2.94 (t, J=11.67 Hz, 1H), 3.06 (d, J=7.44 Hz, 2H), 3.11 (dd,
J=5.13, 12.55 Hz, 1H), 3.32-3.39 (m, 2H), 3.71 (d, J=7.61 Hz,
1H), 4.05-4.09 (m, 1H), 4.23-4.27 (m, 1H), 4.38 (d, J=4.62 Hz,
1H), 4.43 (dd, J=4.87, 10.82 Hz, 1H), 4.52-4.55 (m, 1H),
20 4.75-4.81 (m, 1H), 6.85-6.86 (m, 2H), 6.96 (s, 1H), 7.01-7.04
(m, 1H), 7.11-7.15 (m, 4H), 7.26-7.31 (m, 3H), 7.33-7.37 (m,
3H), 7.51 (d, J=7.92 Hz, 1H), 7.96 (d, J=6.08 Hz, 1H), 8.02
(d, J=6.10 Hz, 1H), 8.08 (d, J=7.90 Hz, 1H), 8.35 (s, 1H).
HR-FAB-MS m/z 816.3310 (M+Na calcd for C₄₂H₄₇N₇O₉, 816.3333, 3
25 ppm err).

Cyclo (D-Pro-D-Trp-Phe-Thr-Phe-Pro) (18c) $[a]^{25}_D$
= +28.74 (C = 0.32, CH₃CN); HR-FAB-MS m/z 798.3536 (M+Na
calcd for C₄₃H₄₉N₇O₇, 798.3591, 4 ppm err).

Cyclo (Ala-D-Trp-Phe-Thr-Phe-Pro) (18d) $[a]^{25}_D$
30 = -65.55 (C = 0.56, CH₃CN); ¹H NMR (500 MHz, CD₃OD) δ
1.02-1.10 (m, 1H), 1.10 (d, J=6.40 Hz, 3H), 1.18 (d, J=6.71
Hz, 3H), 1.53-1.65 (m, 1H), 1.62-1.70 (m, 1H), 1.88 (dd,
J=5.95, 12.03 Hz, 1H), 2.78 (dd, J=4.75, 14.16 Hz, 1H),
2.85-2.96 (m, 2H), 3.0-3.07 (m, 3H), 3.35-3.44 (m, 1H), 3.66
35 (dd, J=7.98 Hz, 1H), 4.10-4.12 (m, 1H), 4.33-4.40 (m, 3H),
4.56-4.75 (m, 3H), 6.84-6.86 (m, 2H), 6.98 (s, 1H), 7.02-7.05
(m, 1H), 7.10-7.14 (m, 1H), 7.15-7.19 (m, 2H), 7.22 (d, J=7.0

- 168 -

Hz, 2H), 7.27-7.36 (m, 5H), 7.51 (d, J=7.87 Hz, 1H), 7.79 (d, J=5.61 Hz, 1H), 7.99 (d, J=7.10 Hz, 1H), 8.18 (d, J=3.60 Hz, 1H), 8.34 (d, J=7.68 Hz, 1H); HR-FAB-MS m/z 772.3437 (M+Na calcd for $C_{41}H_{47}N_7O_7$, 772.3435, <1 ppm err)

5 Cyclo (Trp-D-Trp-Phe-Thr-Phe-Pro) (18e) $[a]^{25}_D$ = -56.41 (C = 0.56, CH_3CN); 1H NMR (500 MHz, CD_3OD) d 0.85-0.89 (m, 2H), 1.15 (d, J=6.35 Hz, 3H), 1.34-1.40 (m, 1H), 1.69-1.73 (m, 1H), 2.59 (dd, J=5.45, 13.8 Hz, 1H), 2.71 (dd, J=4.89, 14.37 Hz, 1H), 2.83-2.88 (m, 2H), 2.90-2.95 (m, 2H),
10 3.05-3.08 (dd, J=5.69, 12.69 Hz, 1H), 3.12-3.21 (m, 3H), 3.35 (d, J=1.0 Hz, 1H), 3.61 (d, J=6.80 Hz, 1H), 4.09-4.13 (m, 1H), 4.23 (dd, J=4.89, 8.32 Hz, 1H), 4.36-4.39 (m, 2H), 4.43 (dd, J=5.61, 9.81 Hz, 1H), 6.76 (d, J=7.21 Hz, 2H), 6.83 (s, 1H), 6.96-6.99 (m, 2H), 7.02-7.15 (m, 7H), 7.21-7.23 (m, 2H),
15 7.25-7.36 (m, 7H), 7.44 (d, J=7.77 Hz, 1H), 7.53 (d, J=7.58 Hz, 1H); ^{13}C NMR (125 MHz, CD_3OD , major) d 18.92, 22.06, 28.32, 29.35, 31.75, 37.42, 38.71, 47.32, 55.33, 56.07, 56.23, 57.13, 57.98, 62.57, 68.61, 110.49, 110.89, 112.29, 112.51, 119.44, 119.49, 119.83, 119.99, 122.49, 124.49, 124.69,
20 127.80, 128.48, 128.61, 129.03, 129.56, 129.79, 130.01, 130.63, 136.94, 137.60, 137.99, 138.05, 171.46, 172.50, 172.74, 173.18, 173.67, 174.33; HR-FAB-MS m/z 887.3879 (M+Na calcd for $C_{49}H_{52}N_8O_7$, 887.3857, 2.5 ppm err).

Cyclo (D-Phe-D-Trp-Phe-Thr-Phe-Pro) (18f) HR-FAB-MS m/z
25 848.3721 (M+Na calcd for $C_{47}H_{51}N_7O_7$, 848.3748, 4 ppm err)

Cyclo (D-Homo-Phe-D-Trp-Phe-Thr-Phe-Pro) (18g)
HR-FAB-MS m/z 862.3911 (M+Na calcd for $C_{48}H_{53}N_7O_7$, 862.3904, 1 ppm err)

Cyclo (Cha-D-Trp-Phe-Thr-Phe-Pro) (18h) $[a]^{25}_D$
30 = -63.18 (C = 0.53, CH_3CN); 1H NMR (500 MHz, CD_3OD) d 0.78-0.90 (m, 3H), 0.98-1.12 (m, 5H), 1.14 (d, J=6.39 Hz, 3H), 1.42-1.53 (m, 2H), 1.56-1.62 (m, 5H), 1.63-1.71 (m, 1H), 1.91 (dd, J=6.33, 12.49 Hz, 1H), 2.86-2.88 (m, 2H), 2.92-2.96 (m, 1H), 2.98-3.02 (m, 2H), 3.09 (dd, J=5.36, 12.69 Hz, 1H),
35 3.36-3.39 (m, 2H), 3.73 (d, J=7.71 Hz, 1H), 4.06-4.08 (m, 1H), 4.30 (dd, J=5.45, 8.49 Hz, 1H), 4.40 (d, J=4.49 Hz, 1H), 4.43-4.51 (m, 2H), 4.55 (t, J=7.78 Hz, 1H), 6.91-6.92 (m, 2H),

- 169 -

6.98 (s, 1H), 7.02-7.05 (m, 1H), 7.10-7.15 (m, 1H), 7.15-7.17 (m, 3H), 7.25-7.26 (m, 2H), 7.28-7.30 (m, 1H), 7.35 (t, J=8.12 Hz, 3H), 7.52 (d, J=7.90 Hz, 1H); ¹³C NMR (125 MHz, CD₃OD, major) δ 17.61, 21.14, 25.79, 25.86, 26.00, 27.12, 30.31, 32.54, 33.12, 33.98, 36.10, 37.35, 39.96, 45.97, 51.24, 53.88, 54.75, 55.86, 56.43, 61.39, 67.10, 109.16, 111.15, 117.87, 118.49, 121.14, 123.10, 126.44, 127.13, 127.26, 128.19, 128.44, 128.65, 129.28, 135.56, 136.54, 136.75, 170.17, 170.99, 171.80, 171.93, 172.35, 173.23. HR-FAB-MS m/z 854.4203 (M+Na calcd for C₄₇H₅₇N₇O₇, 854.4218, 5 ppm err).

Cyclo (Nal-D-Trp-Phe-Thr-Phe-Pro) (18i) [α]_D²⁵ = -52.36 (c = 0.73, CH₃CN); ¹H NMR (500 MHz, CD₃OD) δ 0.82-0.90 (m, 2H), 1.16 (d, J=6.30 Hz, 3H), 1.38-1.41 (m, 1H), 1.72-1.74 (m, 1H), 2.50 (dd, J=5.35, 13.6 Hz, 1H), 2.72-2.90 (m, 3H), 2.93-2.99 (m, 2H), 3.05-3.09 (dd, J=5.57, 12.8 Hz, 1H), 3.16-3.22 (m, 1H), 3.42 (dd, J=7.48, 13.65 Hz, 1H), 3.52 (dd, J=7.98, 13.88 Hz, 1H), 3.62 (d, J=6.99 Hz, 1H), 4.12-4.16 (m, 1H), 4.22 (dd, J=4.63, 8.03 Hz, 1H), 4.35-4.40 (m, 2H), 4.43 (dd, J=5.57, 9.58 Hz, 1H), 4.81 (t, J=7.49 Hz, 1H), 6.78 (t, J=5.22 Hz, 3H), 7.05 (t, J=7.85 Hz, 1H), 7.09-7.15 (m, 4H), 7.22 (d, J=7.53 Hz, 2H), 7.27 (d, J=6.67 Hz, 2H), 7.31 (t, J=6.40 Hz, 3H), 7.34-7.37 (m, 1H), 7.43 (d, J=7.86 Hz, 1H), 7.45-7.52 (m, 2H), 7.73 (d, J=8.11 Hz, 1H), 7.83 (d, J=7.98 Hz, 1H), 8.22 (8.36, 1H); ¹³C NMR (125 MHz, CD₃OD, 315K) δ 18.98, 22.19, 28.41, 31.63, 36.35, 37.46, 38.67, 47.34, 55.39, 55.86, 56.26, 57.23, 57.98, 62.53, 68.64, 110.31, 112.57, 119.35, 119.84, 122.52, 124.42, 124.94, 126.41, 126.79, 127.44, 127.80, 128.42, 128.64, 128.72, 128.85, 129.57, 129.75, 129.79, 130.2, 130.64, 133.65, 134.35, 135.40, 136.90, 137.64, 138.07, 171.54, 172.07, 172.63, 173.14, 173.70, 174.28; HR-FAB-MS m/z 898.3926 (M+Na calcd for C₅₁H₅₃N₇O₇, 898.3904, 5 ppm err).

EXAMPLE 17**Synthesis of Cyclo (Nal-D-Trp-p-F-Phe-Thr-Phe-Pro) (19)**

Compound (19) was synthesized as a single compound using the same general procedure as for the synthesis of

- 170 -

compounds (17a-d). The final product was purified by using a RP-HPLC (C18 Dynamax 300 (21.4 x 250 mm) column, gradient 35-25'-95% buffer B, flow rate = 12 mL/min. $[\alpha]^{25}_D = -64.25$ (C = 0.73 CH₃CN); ¹H NMR (CD₃OD MHz, 500 (315K)) d 0.77-0.80 (m, 1H), 0.87-0.88 (m, 1H), 1.17 (d, J=6.39 Hz, 3H), 1.37-1.40 (m, 1H), 1.74 (dd, J=5.52, 11.62 Hz, 1H), 2.46 (dd, J=5.11, 13.74 Hz, 1H), 2.71 (dd, J=4.90, 14.42 Hz, 1H), 2.80 (dd, J=9.94, 14.17 Hz, 2H), 2.92 (t, J=11.1 Hz, 1H), 2.98 (t, J=9.93 Hz, 1H), 3.05 (dd, J=5.53, 12.74 Hz, 1H), 3.15-3.19 (m, 1H), 3.43 (dd, J=7.6, 13.6 Hz, 1H), 3.53 (dd, J=7.70, 13.63 Hz, 1H), 3.61 (d, J=7.44 Hz, 1H), 4.11-4.16 (m, 2H), 4.36 (dd, J=5.50, 10.91 Hz, 1H), 4.39 (d, J=4.82 Hz, 1H), 4.43 (dd, J=5.19, 10, 37 Hz, 1H), 6.69-6.72 (m, 2H), 6.74-6.78 (m, 2H), 6.82 (s, 1H), 7.05 (t, J=7.84 Hz, 1H), 7.14 (t, J=7.88 Hz, 1H), 7.20-7.22 (m, 2H), 7.25-7.37 (m, 7H), 7.42 (d, J=7.91 Hz, 1H), 7.46-7.49 (m, 1H), 7.51-7.54 (m, 1H), 7.73 (d, J=7.99 Hz, 1H), 7.84 (d, J=8.07 Hz, 1H), 8.24 (d, J=8.38 Hz, 1H); ¹³C NMR (125 MHz, CD₃OD (315K)) d 18.99, 22.20, 28.44, 31.66, 36.38, 36.55, 38.65, 47.35, 55.38, 55.86, 56.31, 57.27, 57.96, 62.52, 68.66, 110.23, 112.58, 116.10 (d, J = 21.50 Hz), 119.36, 119.84, 122.51, 124.52, 124.95, 126.41, 126.80, 127.44, 128.39, 128.63, 128.73, 128.85, 129.79, 130.01, 130.62, 131.35, 133.50, 133.64, 134.36, 135.39, 136.86, 138.04, 163.11 (d, J = 243.3 Hz), 171.52, 172.03, 172.62, 173.12, 173.52, 174.21; HR-FAB-MS m/z 916.3810 (M+Na calcd for C₅₁H₅₂N₇O₇F, 916.3810, <1 ppm err).

EXAMPLE 18

The affinity of a variety of compounds for the substance P receptor was determined employing the following procedure.

A. Receptor Expression in COS

To express the cloned human neurokinin-1 receptor (NK1R) transiently in COS, the cDNA for the human NK1R was cloned into the expression vector pCDM9 which was derived from pCDM8 (Invitrogen) by inserting the ampicillin resistance gene (nucleotide 1973 to 2964 from Bluescript SK+) into the Sac II

- 171 -

site. Transfection of 20 μ g of the plasmid DNA into 10 million COS cells was achieved by electroporation in 800 μ l of the transfection buffer (135 mM CaCl_2 , 1.2 mM MgCl_2 , 2.4 mM K_2HPO_4 , 0.6 mM KH_2PO_4 , 10 nM glucose, 10 mM HEPES pH 7.4) at 260 V and 950 μ F using the IBI Genezapper (IBI, New Haven, CT). The cells were incubated in 10% fetal calf serum, 2 mM glutamine, 100 U/ml penicillin-streptomycin, and 90% DMEM media (Gibco, Grand Island, NY) in 5% CO_2 at 37°C for three days before the binding assay.

10 **B. Assay Protocol using COS**

The binding assay of human NK1R expressed in COS cells is based on the use of ^{125}I -substance P (^{125}I -SP, from DuPont, Boston, MA) as a radioactively labeled ligand which competes with unlabeled substance P or any other ligand for binding to the human NK1R. Monolayer cell cultures of COS were dissociated by the non-enzymatic solution (Specialty Media, Lavallette, NJ) and resuspended in appropriate volume of the bind buffer (50 mM Tris pH 7.5, 5 mM MnCl_2 , 150 mM NaCl, 0.04 mg/ml bacitracin, 0.004 mg/ml leupeptin, 0.2 mg/ml BSA, 0.01 mM phosphoramidon) such that 200 μ l of the cell suspension would give rise to about 10,000 cpm of specific ^{125}I -SP binding (approximately 50,000 to 200,000 cells). In the binding assay, 200 μ l of cells were added to a tube containing 20 μ l of 1.5 to 2.5 nM of ^{125}I -SP and 20 μ l of unlabeled substance P or any other test compound. The tubes were incubated at 4°C or at room temperature for 1 hour with gentle shaking. The bound radioactivity was separated from unbound radioactivity by GF/C filter (Brandel, Gaithersburg, MD) which was pre-wetted with 0.1 polyethylenimine. The filter was washed with 3 ml of wash buffer (50 mM Tris pH 7.5, 5 mM MnCl_2 , 150 mM NaCl) three times and its radioactivity was determined by gamma counter.

A variety of compounds were tested according to the COS cell procedure. The concentration of compound required to inhibit the binding of substance P to the human neurokinin-1 receptor by 50% was measured. The following data were obtained:

- 172 -

<u>Compound</u>	<u>IC₅₀</u>
1	120 nM
2	180 nM
7	56 nM
8	840 nM
9	400 nM
11	400 nM
13	1000 nM

EXAMPLE 19

10 The affinity of a variety of compounds for the SRIF
receptor was determined by studying the displacement of ¹²⁵I-
CGP-23996 from AtT-20 cells using the method generally in
accordance with Raynor and Reisine, *Journal of Pharmacology*
and *Experimental Therapeutics*, 1989, 251;2, 510. The
15 following data were obtained:

<u>Compound</u>	<u>IC₅₀</u>
1	9500 nM
2	1300 nM
8	40000 nM
13	does not bind
SRIF	9.3 nM
MK 678	60 nM
L-363,301	18.7 nM

EXAMPLE 20

25 The affinity of a 2-(1-phenylsulfonyl-indol-3yl)ethyl-
6-O-(5-aminopentyl)-2,3,4-tri-O-benzyl-β-D-glucopyranoside,
structure (1) and methyl 6-O-(5-aminopentyl)-2,3,4-Tri-O-
benzyl-β-D-glucopyranoside, structure (8) for a variety of G-
protein-linked receptors was determined by studying the
30 displacement of a variety of radioligands from AtT-20 and
brain cells using the method disclosed by Reisine, et al.,
Brain Research, 1979, 117, 241. The following data was
obtained (¹²⁵I-CYP = ¹²⁵I-cyanopindolol; ³H-QNB = quinuclidinyl
benzilate):

- 173 -

	<u>Receptor</u>	<u>Radioligand</u>	<u>Compound</u>	<u>Binding Inhibition</u>	<u>Tissue</u>
5	β -Adrenergic	¹²⁵ I-CYP (0.1 nM)	1	70%	AtT-20
			1	45	Brain
			8	0	AtT-20
	Opiate Receptor	³ H-naloxone (0.5 nM)	1	55	Brain
	Dopamine Receptor	³ H-spiperone (0.1 nM)	1	82	Brain
10	Muscarinic cholinergic	³ H-QNB 0.1 (nM)	1	20	AtT-20
			1	83	Brain

As can be seen from these Examples, the peptide analogs of the present invention are selectively bound by certain receptors. For example, structure (1) exhibits approximately 14-fold greater selectivity than structure (8) for the substance P receptor, while structure (8) is bound by the substance P and SRIF receptors but does not bind the β -adrenergic receptor.

EXAMPLE 21

The extent to which the cyclic hexapeptides of the invention inhibit HIV-1 protease was determined generally according to the methods disclosed by Berridge, et al., *Biochemistry Journal* 1982, 206, 587, and Cascieri, et al., *J. Pharmacol. Toxicol. Meth.*, 1995, 33, 179. The following data were obtained:

- 174 -


	Compound	IC ₅₀ (nM)
5	(17a)	3420
	(17b)	330
	(17c)	2150
	(17d)	9000
	(17e)	130
	(17f)	37
10	(18a)	2210
	(18b)	4020
	(18c)	51
	(18d)	2200
	(18e)	73
15	(18f)	1120
	(18g)	161
	(18h)	700
	(18i)	22
	(19)	2.8

20 Those skilled in the art will appreciate that numerous
changes and modifications may be made to the preferred
embodiments of the invention and that such changes and
modifications may be made without departing from the spirit
of the invention. It is therefore intended that the appended
25 claims cover all such equivalent variations as fall within the
true spirit and scope of the invention.

- 175 -

WHAT IS CLAIMED IS:

1. A compound having the structure:



wherein:

R_1 is $-(CH_2)_n R_A$ or $-C(O)(CH_2)_n R_A$ where R_A is $-H$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms, and n is an integer from 0 to about 12;

at least two of R_2 , R_3 , and R_4 , independently, are $-(CH_2)_m R_B$ or $-C(O)(CH_2)_m R_B$ where R_B is aryl, and m is an integer from 0 to about 5;

R_5 is $-(CH_2)_p NHR_C$, $-C(O)(CH_2)_p NHR_C$, $-(CH_2)_p R_D$, $-(CH_2)_p R_E$ or $-C(O)(CH_2)_p R_D$;

p is an integer from 0 to about 10;

R_C is $-R_E$ or $-C(O)R_E$;

R_D is $-H$, $-OR_E$, or $-C(O)R_E$;

R_E is $-H$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms; or a pharmaceutically acceptable salt thereof, provided that R_1 is not $-(CH_2)_2(3\text{-indole})$ where R_2 and R_4 are benzyl, R_3 is benzyl or H , and R_5 is $-(CH_2)_5 NH_2$ or $-(CH_2)_5 NHC(O)CH_3$.

2. The compound of claim 1 wherein R_A is aryl having at least one nitrogen atom.

3. The compound of claim 1 wherein R_A is indolyl.

- 176 -

4. The compound of claim 1 wherein R_A is 3-indolyl.
5. The compound of claim 1 wherein n is 2.
6. The compound of claim 1 wherein R_B is phenyl.
7. The compound of claim 1 wherein m is 1.
8. The compound of claim 1 wherein m is 0.
9. The compound of claim 1 wherein p is from about 2 to about 8.
10. The compound of claim 1 wherein p is from about 3 to about 6.
11. The compound of claim 1 wherein p is 5.
12. The compound of claim 1 wherein R_5 is $-(CH_2)_pNH_2$.
13. The compound of claim 1 wherein R_5 is $-(CH_2)_pNHR_C$.
14. The compound of claim 6 wherein R_C is $-CH_3$.
15. The compound of claim 1 wherein R_B is fluorophenyl or naphthyl.
16. The compound of claim 1 wherein R_2 is CH_2 -fluorophenyl.
17. The compound of claim 1 wherein R_2 is CH_2 -naphthyl.
18. The compound of claim 1 wherein R_E is fluorophenyl.
19. The compound of claim 1 wherein R_E is naphthyl.

- 177 -


20. The compound of claim 1 wherein R_5 is CH_2 -fluorophenyl.

21. The compound of claim 1 wherein R_5 is CH_2 -naphthyl.

22. The compound of claim 1 wherein R_1 is $(\text{CH}_2)_2$ -indolyl, R_2 is CH_2 -fluorophenyl, R_3 and R_4 are benzyl, and R_5 is CH_2 -naphthyl.

23. The compound of claim 1 wherein R_1 is $(\text{CH}_2)_2$ -indolyl, R_2 is CH_2 -naphthyl, R_3 and R_4 are benzyl, and R_5 is CH_2 -fluorophenyl.

24. A compound having the structure:



wherein:

R_1 is $-(\text{CH}_2)_n R_A$ or $-\text{C}(\text{O})(\text{CH}_2)_n R_A$ where R_A is $-\text{H}$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms, and n is an integer from 0 to about 12;

at least two of R_2 , R_3 , and R_4 , independently, are $-(\text{CH}_2)_m R_B$ or $-\text{C}(\text{O})(\text{CH}_2)_m R_B$ where R_B is aryl, and m is an integer from 0 to about 5;

R_5 is $-(\text{CH}_2)_p \text{NHR}_C$, $-\text{C}(\text{O})(\text{CH}_2)_p \text{NHR}_C$, $-(\text{CH}_2)_p R_D$, $-(\text{CH}_2)_p R_E$ or $-\text{C}(\text{O})(\text{CH}_2)_p R_D$;

p is an integer from 0 to about 10;

R_C is $-\text{R}_8$ or $-\text{C}(\text{O})\text{R}_8$;


R_D is $-\text{H}$, $-\text{OR}_8$, or $-\text{C}(\text{O})\text{R}_8$;

R_8 is $-\text{H}$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl

- 178 -

having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms; or a pharmaceutically acceptable salt thereof, provided that R_1 is not $-(CH_2)_2(3\text{-indole})$ where R_2 and R_4 are benzyl, R_3 is benzyl or H, and R_5 is $-(CH_2)_5NH_2$ or $-(CH_2)_5NHC(O)CH_3$.

25. A compound having the structure:



wherein:

R_1 is $-O(CH_2)_nR_A$, $-OC(O)(CH_2)_nR_A$, $-(CH_2)_nR_A$, or $-C(O)(CH_2)_nR_A$ where R_A is $-H$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms, and n is an integer from 0 to about 12;

at least one of R_2 , R_3 , and R_4 , independently, is $-O(CH_2)_mR_B$, $-OC(O)(CH_2)_mR_B$, $-(CH_2)_mR_B$ or $-C(O)(CH_2)_mR_B$ where R_B is $-H$ or aryl, and m is an integer from 0 to about 5; and

R_5 is $-O(CH_2)_pNHR_C$, $-OC(O)(CH_2)_pNHR_C$, $-O(CH_2)_pR_D$, $-OC(O)(CH_2)_pR_D$, $-(CH_2)_pNHR_C$, $-C(O)(CH_2)_pNHR_C$, $-(CH_2)_pR_D$ or $-C(O)(CH_2)_pR_D$, where:

p is an integer from 0 to about 10;

R_C is $-R_E$ or $-C(O)R_E$;

R_D is $-H$, $-OR_E$, or $-C(O)R_E$;

R_E is $-H$, alkyl or alkenyl having from about 1 to about 14 carbon atoms and up to about 4 nitrogen atoms, or aryl having from about 6 to about 14 carbon atoms and up to about 4 nitrogen atoms; or a pharmaceutically acceptable salt thereof.

- 179 -

26. A method for producing a prophylactic or therapeutic response in a mammal, comprising administering to the mammal an effective amount of a compound of claim 25.

27. The method of claim 26 wherein the prophylactic or therapeutic response is a modulation of inflammation in the mammal.

28. The method of claim 26 wherein the prophylactic or therapeutic response is a modulation of pain in the mammal.

29. A method for modulating the activity of at least one mammalian G-protein-linked receptor, comprising administering to a mammal an effective amount of a compound of claim 25.


30. The method of claim 29 wherein at least one G-protein-linked receptor is a somatostatin receptor.

31. A method for modulating the activity of at least one mammalian Substance P receptor, comprising administering to a mammal an effective amount of a compound of claim 25.

32. A method for mimicking or inhibiting the chemical activity of a peptide, comprising providing in place of the peptide at least one chemical compound of claim 25.

- 180 -

33. A compound having the structure:



wherein:

R_{10} is indolyl;

R_{11} is H, isopropyl, phenyl, 4-hydroxyphenyl, 4-methoxyphenyl, or fluorophenyl;

R_{12} is phenyl; and

R_{13} is -OH, -C(O)OH, -H, -indolyl, -phenyl, -CH₂-phenyl, -cyclohexyl, or -naphthyl.

34. The compound of claim 33 wherein R_{11} is fluorophenyl.

35. The compound of claim 34 wherein R_{13} is CH₂-naphthyl.

36. A method for producing a prophylactic or therapeutic response in a mammal, comprising administering to the mammal an effective amount of a compound of claim 33.


37. A method for modulating the activity of at least one mammalian G-protein-linked receptor, comprising administering to a mammal an effective amount of a compound of claim 33.

38. A method for modulating the activity of at least one mammalian Substance P receptor, comprising administering to a mammal an effective amount of a compound of claim 33.


- 181 -

39. A method for mimicking or inhibiting the chemical activity of a peptide, comprising providing in place of the peptide at least one chemical compound of claim 33.


FIGURE 1 SHEET 1



SCHEME 1




SCHEME 2




SCHEME 5


FIGURE 2 SHEET 1



SCHEME 6




SCHEME 7




SCHEME 8


FIGURE 2 SHEET 3.



SCHEME 11




SCHEME 12




SCHEME 13

FIGURE 2 SHEET 4




SCHEME 14




SCHEME 15

FIGURE 3 SHEET 2




SCHEME 16

FIGURE 3 SHEET 3




SCHEME 17

FIGURE 3 SHEET 4




SCHEME 18

..GURE 3 SHEET 5




SCHEME 19

FIGURE 3 SHEET 6



SCHEME 20



SCHEME 21

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/01097

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07H 15/00, 17/00; A61K 38/08; C07K 7/66

US CL : 536/17.4, 4.1, 17.2, 17.3; 514/11; 530/311, 317, 329

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/17.4, 4.1, 17.2, 17.3; 514/11; 530/311, 317, 329

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, APS

search terms: somatostatin, hexapeptide, d-glucose

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95/11686 A1 (TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA) 04 May 1995 (04/05/95), see entire document.	1-39
A	US 4,310,518 A (FREIDINGER et al.) 12 January 1982 (12/01/82), see entire document.	33-39
A	US 4,659,691 A (VEBER et al.) 21 April 1987 (21/04/87), see entire document.	33-39
A	US 4,360,516 A (FREIDINGER et al.) 23 November 1982 (23/11/82), see entire document.	33-39

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

20 MAY 1997

Date of mailing of the international search report

11 JUL 1997

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Authorized officer

ANISH GUPTA

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/01097

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/01097

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-32, drawn to a heterocyclic compound and its method of use.

Group II, claim(s) 33-39, drawn to a cyclopeptide compound and its method of use.

The inventions listed as Groups I and Group II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The compounds of Group I are structurally distinct and dissimilar to the cyclopeptide of Group II.